



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 03:33 am GMT

PDB ID : 3S0T
Title : Crystal structure of the CofA Type IV pilin subunit from enterotoxigenic E. coli
Authors : Kolappan, S.; Craig, L.
Deposited on : 2011-05-13
Resolution : 1.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1690 (1.30-1.22)
Clashscore	112137	1781 (1.30-1.22)
Ramachandran outliers	110173	1716 (1.30-1.22)
Sidechain outliers	110143	1714 (1.30-1.22)
RSRZ outliers	101464	1693 (1.30-1.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	184	<div> <div>3%</div> <div> <div></div> <div>80%</div> <div>14%</div> <div>• •</div> </div> </div>
1	B	184	<div> <div>5%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div>• • •</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	1[A]	-	-	-	X
2	GOL	A	1[B]	-	-	-	X
2	GOL	B	2	-	-	-	X
3	NA	B	3	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3173 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CFA/III pilin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	180	Total	C	N	O	S	0	17	0
			1374	831	243	295	5			
1	B	181	Total	C	N	O	S	0	19	0
			1395	847	245	295	8			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	GLY	-	EXPRESSION TAG	UNP Q59393
A	26	SER	-	EXPRESSION TAG	UNP Q59393
A	27	HIS	-	EXPRESSION TAG	UNP Q59393
A	28	MET	-	EXPRESSION TAG	UNP Q59393
B	25	GLY	-	EXPRESSION TAG	UNP Q59393
B	26	SER	-	EXPRESSION TAG	UNP Q59393
B	27	HIS	-	EXPRESSION TAG	UNP Q59393
B	28	MET	-	EXPRESSION TAG	UNP Q59393

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	1
			12	6	6		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	184	Total	O	0	0
			184	184		
4	B	201	Total	O	0	0
			201	201		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CFA/III pilin



- Molecule 1: CFA/III pilin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	29.50Å 34.15Å 64.23Å 95.75° 97.38° 94.00°	Depositor
Resolution (Å)	25.00 – 1.26 24.51 – 1.26	Depositor EDS
% Data completeness (in resolution range)	96.0 (25.00-1.26) 92.9 (24.51-1.26)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.28 (at 1.26Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.139 , 0.181 0.144 , 0.188	Depositor DCC
R_{free} test set	3219 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	10.7	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 52.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	3173	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 85.47 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.2486e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.11	3/1452 (0.2%)	1.06	6/1970 (0.3%)
1	B	1.09	0/1475	1.08	7/1997 (0.4%)
All	All	1.10	3/2927 (0.1%)	1.07	13/3967 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	34[A]	SER	CB-OG	-5.35	1.35	1.42
1	A	34[B]	SER	CB-OG	-5.35	1.35	1.42
1	A	144	GLU	CD-OE1	-5.02	1.20	1.25

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	187	LEU	CB-CG-CD1	10.21	128.37	111.00
1	A	80	VAL	CG1-CB-CG2	8.89	125.12	110.90
1	B	187	LEU	CA-CB-CG	-8.20	96.44	115.30
1	B	116	ILE	CG1-CB-CG2	7.49	127.89	111.40
1	B	196[A]	CYS	CA-CB-SG	7.22	127.00	114.00
1	B	196[B]	CYS	CA-CB-SG	7.22	127.00	114.00
1	B	176	LEU	CB-CG-CD1	-6.02	100.77	111.00
1	B	53	ARG	NE-CZ-NH1	-5.62	117.49	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	112[A]	SER	C-N-CA	-5.55	110.65	122.30
1	A	112[B]	SER	C-N-CA	-5.55	110.65	122.30
1	A	95[A]	ARG	NE-CZ-NH2	5.37	122.99	120.30
1	A	95[B]	ARG	NE-CZ-NH2	5.37	122.99	120.30
1	A	193	LEU	CB-CG-CD2	-5.07	102.38	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	44[A]	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1374	0	1348	43	0
1	B	1395	0	1383	23	0
2	A	12	0	16	1	0
2	B	6	0	8	0	0
3	B	1	0	0	0	0
4	A	184	0	0	16	0
4	B	201	0	0	5	0
All	All	3173	0	2755	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177[B]:ARG:HD2	1:A:187:LEU:HD21	1.07	1.07
1:A:187:LEU:HD23	4:A:372:HOH:O	1.59	1.01
1:A:199[B]:THR:HG21	4:A:285:HOH:O	1.65	0.97
1:A:177[B]:ARG:HD2	1:A:187:LEU:CD2	1.95	0.95

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:LEU:HD22	1:A:187:LEU:N	1.90	0.85
1:B:177[B]:ARG:HD2	1:B:187:LEU:HD22	1.56	0.85
1:A:177[B]:ARG:CD	1:A:187:LEU:HD21	2.01	0.84
1:A:187:LEU:HD22	1:A:187:LEU:H	1.51	0.74
1:A:171[A]:ASP:HB2	4:A:374:HOH:O	1.90	0.72
1:B:176:LEU:N	1:B:176:LEU:HD12	2.05	0.71
1:A:130:GLU:CG	4:A:272:HOH:O	2.39	0.70
1:A:187:LEU:CD2	4:A:372:HOH:O	2.28	0.69
1:A:36:LEU:O	1:A:40:THR:HG23	1.93	0.69
1:B:53:ARG:NH1	4:B:284:HOH:O	2.26	0.69
1:A:177[B]:ARG:HG3	1:A:187:LEU:HD11	1.75	0.69
1:A:195[B]:THR:HG22	1:A:196:CYS:N	2.09	0.68
1:A:171[B]:ASP:HB2	4:A:374:HOH:O	1.95	0.66
1:A:41:ASN:HB2	4:A:373:HOH:O	1.96	0.64
1:B:114:SER:OG	1:B:115:THR:N	2.30	0.64
1:A:199[B]:THR:CG2	4:A:285:HOH:O	2.35	0.63
1:A:130:GLU:HG3	4:A:272:HOH:O	1.95	0.63
1:B:177[B]:ARG:CD	1:B:187:LEU:HD22	2.27	0.62
1:B:116:ILE:HD13	1:B:118:LYS:HE3	1.82	0.61
1:B:177[A]:ARG:HD2	1:B:187:LEU:HD22	1.81	0.61
1:B:122[B]:MET:HE3	1:B:143:TRP:CH2	2.38	0.58
1:A:130:GLU:HG2	4:A:272:HOH:O	2.00	0.58
1:A:53:ARG:HH21	2:A:1[B]:GOL:H31	1.69	0.58
1:A:187:LEU:H	1:A:187:LEU:CD2	2.17	0.56
1:A:187:LEU:N	1:A:187:LEU:CD2	2.63	0.56
1:A:173:THR:O	1:A:195[B]:THR:HG23	2.06	0.56
1:B:177[B]:ARG:CD	1:B:187:LEU:CD2	2.84	0.55
1:B:129[B]:GLN:HG2	1:B:196[B]:CYS:O	2.07	0.55
1:B:28:MET:N	4:B:311:HOH:O	2.39	0.55
1:B:177[B]:ARG:HD2	1:B:187:LEU:CD2	2.34	0.53
1:A:177[B]:ARG:CG	1:A:187:LEU:HD11	2.39	0.53
1:A:177[B]:ARG:CD	1:A:187:LEU:HD11	2.40	0.52
1:B:116:ILE:CD1	1:B:118:LYS:HE3	2.38	0.52
1:B:132[A]:CYS:SG	1:B:196[A]:CYS:SG	3.03	0.52
1:A:176:LEU:HD21	1:A:195[A]:THR:HG21	1.92	0.51
1:A:143:TRP:O	1:A:177[A]:ARG:NH1	2.44	0.50
1:B:36:LEU:O	1:B:40:THR:HG23	2.12	0.49
1:B:52:GLN:HG3	4:B:282:HOH:O	2.13	0.49
1:B:129[A]:GLN:HG3	4:B:250:HOH:O	2.12	0.49
1:A:37:VAL:O	1:A:40:THR:OG1	2.27	0.48
1:A:144:GLU:HG3	4:A:245:HOH:O	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129[A]:GLN:HG2	1:A:196:CYS:O	2.14	0.48
1:A:31:ARG:HH11	1:A:31:ARG:HG3	1.79	0.47
1:A:174[A]:THR:HG22	4:A:318:HOH:O	2.14	0.47
1:A:95[B]:ARG:NH2	4:A:333:HOH:O	2.46	0.45
1:A:145:TYR:HB3	1:A:206:ALA:HB3	1.98	0.45
1:B:122[B]:MET:HE3	1:B:143:TRP:HH2	1.81	0.45
1:B:112:SER:HA	1:B:113:GLY:HA2	1.74	0.45
1:B:176:LEU:N	1:B:176:LEU:CD1	2.78	0.45
1:A:115:THR:HG23	4:A:255:HOH:O	2.17	0.45
1:B:187:LEU:HA	1:B:187:LEU:HD12	1.86	0.44
1:A:95[B]:ARG:NH1	4:A:333:HOH:O	2.30	0.43
1:A:141:ASP:HA	1:A:177[A]:ARG:HH21	1.84	0.42
1:A:31:ARG:HH11	1:A:31:ARG:CG	2.33	0.42
1:A:41:ASN:OD1	1:A:44[B]:ARG:NH1	2.53	0.42
1:A:141:ASP:HA	1:A:177[A]:ARG:NH2	2.35	0.41
1:B:61:GLN:HG2	4:B:372:HOH:O	2.21	0.41
1:A:192:ILE:HA	1:A:195[A]:THR:HG22	2.03	0.41
1:A:65:SER:HB2	4:A:328:HOH:O	2.20	0.41
1:B:145:TYR:HB3	1:B:206:ALA:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/184 (106%)	193 (99%)	2 (1%)	0	100	100
1	B	198/184 (108%)	195 (98%)	3 (2%)	0	100	100
All	All	393/368 (107%)	388 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/146 (108%)	152 (96%)	6 (4%)	38	4
1	B	160/146 (110%)	153 (96%)	7 (4%)	33	3
All	All	318/292 (109%)	305 (96%)	13 (4%)	37	3

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ARG
1	A	115	THR
1	A	130	GLU
1	A	172	ASN
1	A	176	LEU
1	A	187	LEU
1	B	30	SER
1	B	115	THR
1	B	130[A]	GLU
1	B	130[B]	GLU
1	B	187	LEU
1	B	196[A]	CYS
1	B	196[B]	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	1[A]	-	5,5,5	0.62	0	5,5,5	1.07	1 (20%)
2	GOL	A	1[B]	-	5,5,5	0.87	0	5,5,5	1.25	1 (20%)
2	GOL	B	2	-	5,5,5	0.99	0	5,5,5	1.60	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1[A]	-	-	0/4/4/4	0/0/0/0
2	GOL	A	1[B]	-	-	0/4/4/4	0/0/0/0
2	GOL	B	2	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	GOL	O2-C2-C3	-2.73	95.96	108.84
2	A	1[A]	GOL	C3-C2-C1	-2.09	103.22	111.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	GOL	O1-C1-C2	2.02	120.24	110.07
2	A	1[B]	GOL	O2-C2-C3	2.16	119.05	108.84

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1[B]	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	180/184 (97%)	-0.17	6 (3%)	47	40	7, 12, 21, 31	6 (3%)
1	B	181/184 (98%)	-0.11	10 (5%)	26	23	6, 12, 22, 34	4 (2%)
All	All	361/368 (98%)	-0.14	16 (4%)	35	29	6, 12, 22, 34	10 (2%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	116	ILE	4.6
1	B	113	GLY	4.5
1	B	114	SER	4.3
1	B	115	THR	3.5
1	B	28	MET	3.5
1	B	196[A]	CYS	3.3
1	B	29	ASP	3.0
1	A	196	CYS	2.7
1	A	172	ASN	2.7
1	A	31	ARG	2.7
1	B	31	ARG	2.5
1	A	173	THR	2.3
1	B	32	THR	2.2
1	A	29	ASP	2.2
1	B	30	SER	2.1
1	A	115	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NA	B	3	1/1	0.97	0.35	7.43	13,13,13,13	0
2	GOL	B	2	6/6	0.88	0.12	7.37	26,30,34,36	0
2	GOL	A	1[B]	6/6	0.91	0.19	6.82	17,20,25,27	6
2	GOL	A	1[A]	6/6	0.91	0.19	6.48	15,21,23,30	6

6.5 Other polymers [i](#)

There are no such residues in this entry.