



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2017 – 04:02 PM EDT

PDB ID : 3SHT  
Title : Crystal structure of human MCPH1 tandem BRCT domains  
Authors : Shao, Z.H.; Li, F.D.; Yan, W.  
Deposited on : unknown  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	206	<div> <div>2%</div> <div>87%</div> <div>9%</div> </div>
1	B	206	<div> <div>2%</div> <div>89%</div> <div>9%</div> </div>
1	C	206	<div> <div>2%</div> <div>85%</div> <div>5%</div> <div>10%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4796 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Microcephalin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	188	Total	C	N	O	S	0	0	0
			1457	939	248	260	10			
1	B	187	Total	C	N	O	S	0	0	0
			1450	934	247	259	10			
1	C	186	Total	C	N	O	S	0	0	0
			1442	928	246	258	10			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	630	MET	-	EXPRESSION TAG	UNP Q8NEM0
A	631	GLY	-	EXPRESSION TAG	UNP Q8NEM0
A	632	HIS	-	EXPRESSION TAG	UNP Q8NEM0
A	633	HIS	-	EXPRESSION TAG	UNP Q8NEM0
A	634	HIS	-	EXPRESSION TAG	UNP Q8NEM0
A	635	HIS	-	EXPRESSION TAG	UNP Q8NEM0
A	636	HIS	-	EXPRESSION TAG	UNP Q8NEM0
A	637	HIS	-	EXPRESSION TAG	UNP Q8NEM0
A	638	MET	-	EXPRESSION TAG	UNP Q8NEM0
A	761	ALA	VAL	SEE REMARK 999	UNP Q8NEM0
B	630	MET	-	EXPRESSION TAG	UNP Q8NEM0
B	631	GLY	-	EXPRESSION TAG	UNP Q8NEM0
B	632	HIS	-	EXPRESSION TAG	UNP Q8NEM0
B	633	HIS	-	EXPRESSION TAG	UNP Q8NEM0
B	634	HIS	-	EXPRESSION TAG	UNP Q8NEM0
B	635	HIS	-	EXPRESSION TAG	UNP Q8NEM0
B	636	HIS	-	EXPRESSION TAG	UNP Q8NEM0
B	637	HIS	-	EXPRESSION TAG	UNP Q8NEM0
B	638	MET	-	EXPRESSION TAG	UNP Q8NEM0
B	761	ALA	VAL	SEE REMARK 999	UNP Q8NEM0
C	630	MET	-	EXPRESSION TAG	UNP Q8NEM0
C	631	GLY	-	EXPRESSION TAG	UNP Q8NEM0
C	632	HIS	-	EXPRESSION TAG	UNP Q8NEM0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	633	HIS	-	EXPRESSION TAG	UNP Q8NEM0
C	634	HIS	-	EXPRESSION TAG	UNP Q8NEM0
C	635	HIS	-	EXPRESSION TAG	UNP Q8NEM0
C	636	HIS	-	EXPRESSION TAG	UNP Q8NEM0
C	637	HIS	-	EXPRESSION TAG	UNP Q8NEM0
C	638	MET	-	EXPRESSION TAG	UNP Q8NEM0
C	761	ALA	VAL	SEE REMARK 999	UNP Q8NEM0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	156	Total	O	0	0
			156	156		
2	B	158	Total	O	0	0
			158	158		
2	C	133	Total	O	0	0
			133	133		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



#### • Molecule 1: Microcephalin



#### • Molecule 1: Microcephalin



#### • Molecule 1: Microcephalin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.47Å 113.67Å 74.58Å 90.00° 91.04° 90.00°	Depositor
Resolution (Å)	32.58 – 1.95 32.58 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.2 (32.58-1.95) 97.2 (32.58-1.95)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.40 (at 1.95Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.202 , 0.242 0.202 , 0.242	Depositor DCC
$R_{free}$ test set	2767 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.6	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 51.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.032 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4796	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.67	0/1498	0.66	0/2043
1	B	0.68	0/1490	0.67	0/2032
1	C	0.61	0/1482	0.66	0/2021
All	All	0.66	0/4470	0.66	0/6096

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1457	0	1479	5	2
1	B	1450	0	1471	2	1
1	C	1442	0	1460	8	0
2	A	156	0	0	2	1
2	B	158	0	0	1	0
2	C	133	0	0	1	0
All	All	4796	0	4410	15	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:762:MET:HE2	1:C:797:ILE:HG13	1.76	0.66
1:A:738:PRO:HG2	2:A:354:HOH:O	1.96	0.65
1:C:693:ARG:HD3	1:C:821:THR:HG21	1.82	0.61
1:A:709:SER:HB3	1:A:730:LEU:HD21	1.85	0.58
1:C:693:ARG:CD	1:C:821:THR:HG21	2.35	0.57
1:C:647:THR:HA	2:C:424:HOH:O	2.09	0.51
1:C:763:PHE:CE2	1:C:790:PRO:HG3	2.47	0.49
1:C:693:ARG:HD2	1:C:698:LEU:HD21	1.95	0.48
1:A:769:SER:HA	1:A:770:PRO:C	2.35	0.47
1:B:768:SER:HB2	1:B:813:GLU:HG2	1.97	0.45
1:C:762:MET:CE	1:C:815:TRP:HZ3	2.31	0.44
1:B:758:ASP:HB2	2:B:117:HOH:O	2.17	0.44
1:A:774:LYS:HD3	1:A:774:LYS:HA	1.83	0.43
1:A:681:GLU:HG3	2:A:284:HOH:O	2.19	0.42
1:C:762:MET:HE2	1:C:797:ILE:CG1	2.45	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:803:LYS:NZ	2:A:354:HOH:O[1_455]	2.18	0.02
1:A:794:SER:OG	1:B:794:SER:OG[1_455]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/206 (90%)	184 (99%)	2 (1%)	0	100	100
1	B	185/206 (90%)	183 (99%)	2 (1%)	0	100	100
1	C	184/206 (89%)	181 (98%)	2 (1%)	1 (0%)	32	19
All	All	555/618 (90%)	548 (99%)	6 (1%)	1 (0%)	51	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	691	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/180 (92%)	165 (100%)	0	100	100
1	B	164/180 (91%)	164 (100%)	0	100	100
1	C	163/180 (91%)	163 (100%)	0	100	100
All	All	492/540 (91%)	492 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	660	GLN
1	A	665	GLN
1	B	721	HIS
1	B	733	HIS
1	C	745	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	188/206 (91%)	-0.08	4 (2%) 64 73	12, 23, 36, 42	2 (1%)
1	B	187/206 (90%)	0.14	4 (2%) 64 73	13, 24, 38, 50	2 (1%)
1	C	186/206 (90%)	0.17	5 (2%) 55 65	15, 27, 43, 51	6 (3%)
All	All	561/618 (90%)	0.08	13 (2%) 61 71	12, 25, 39, 51	10 (1%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	792	GLN	3.5
1	C	789	VAL	3.2
1	B	833	LEU	3.1
1	B	830	ASN	2.4
1	B	750	PRO	2.4
1	C	766	PRO	2.4
1	B	780	HIS	2.3
1	A	646	PRO	2.2
1	A	829	GLU	2.2
1	C	791	ARG	2.1
1	A	732	HIS	2.1
1	A	749	GLY	2.0
1	C	790	PRO	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.