



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:51 pm GMT

PDB ID : 1SMC
Title : Mycobacterium tuberculosis dUTPase complexed with dUTP in the absence of metal ion.
Authors : Sawaya, M.R.; Chan, S.; Segelke, B.; Lakin, T.; Krupka, H.; Cho, U.S.; Kim, M.-Y.; So, M.; Kim, C.-Y.; Naranjo, C.M.; Rogers, Y.C.; Park, M.S.; Waldo, G.S.; Pashkov, I.; Cascio, D.; Yeates, T.O.; Perry, J.L.; Terwilliger, T.C.; Eisenberg, D.; TB Structural Genomics Consortium (TBSGC)
Deposited on : 2004-03-09
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	174	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; bottom: 0; left: 72%; width: 28%; height: 10px; background-color: grey;"></div> <div style="position: absolute; bottom: 0; left: 72%; width: 28%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; bottom: 0; left: 25%; width: 3%; height: 10px; background-color: orange;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 10%; height: 10px; background-color: red;"></div> </div> <div>72% 25%</div> </div>
1	B	174	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; bottom: 0; left: 72%; width: 28%; height: 10px; background-color: grey;"></div> <div style="position: absolute; bottom: 0; left: 72%; width: 28%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; bottom: 0; left: 21%; width: 3%; height: 10px; background-color: orange;"></div> <div style="position: absolute; bottom: 0; left: 3%; width: 10%; height: 10px; background-color: red;"></div> </div> <div>3% 72% 7% 21%</div> </div>
1	C	174	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; bottom: 0; left: 70%; width: 30%; height: 10px; background-color: grey;"></div> <div style="position: absolute; bottom: 0; left: 70%; width: 30%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 10%; height: 10px; background-color: red;"></div> </div> <div>70% 26%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NO3	C	155	-	-	-	X
3	DUT	A	2170	-	-	-	X
3	DUT	C	3170[A]	-	-	-	X
3	DUT	C	3170[B]	-	-	-	X
4	TRS	A	2171	-	-	-	X
4	TRS	B	155	-	-	-	X
4	TRS	C	3171	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3288 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxyuridine 5'-triphosphate nucleotidohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	131	Total	C	N	O	S	0	0	1
			964	608	174	181	1			
1	B	138	Total	C	N	O	S	0	0	0
			1013	637	185	189	2			
1	C	129	Total	C	N	O	S	0	0	0
			951	601	173	176	1			

There are 60 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	CLONING ARTIFACT	UNP P0A552
A	-18	GLY	-	CLONING ARTIFACT	UNP P0A552
A	-17	SER	-	CLONING ARTIFACT	UNP P0A552
A	-16	SER	-	CLONING ARTIFACT	UNP P0A552
A	-15	HIS	-	EXPRESSION TAG	UNP P0A552
A	-14	HIS	-	EXPRESSION TAG	UNP P0A552
A	-13	HIS	-	EXPRESSION TAG	UNP P0A552
A	-12	HIS	-	EXPRESSION TAG	UNP P0A552
A	-11	HIS	-	EXPRESSION TAG	UNP P0A552
A	-10	HIS	-	EXPRESSION TAG	UNP P0A552
A	-9	SER	-	CLONING ARTIFACT	UNP P0A552
A	-8	SER	-	CLONING ARTIFACT	UNP P0A552
A	-7	GLY	-	CLONING ARTIFACT	UNP P0A552
A	-6	LEU	-	CLONING ARTIFACT	UNP P0A552
A	-5	VAL	-	CLONING ARTIFACT	UNP P0A552
A	-4	PRO	-	CLONING ARTIFACT	UNP P0A552
A	-3	ARG	-	CLONING ARTIFACT	UNP P0A552
A	-2	GLY	-	CLONING ARTIFACT	UNP P0A552
A	-1	SER	-	CLONING ARTIFACT	UNP P0A552
A	0	HIS	-	CLONING ARTIFACT	UNP P0A552
B	-19	MET	-	CLONING ARTIFACT	UNP P0A552
B	-18	GLY	-	CLONING ARTIFACT	UNP P0A552
B	-17	SER	-	CLONING ARTIFACT	UNP P0A552

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	SER	-	CLONING ARTIFACT	UNP P0A552
B	-15	HIS	-	EXPRESSION TAG	UNP P0A552
B	-14	HIS	-	EXPRESSION TAG	UNP P0A552
B	-13	HIS	-	EXPRESSION TAG	UNP P0A552
B	-12	HIS	-	EXPRESSION TAG	UNP P0A552
B	-11	HIS	-	EXPRESSION TAG	UNP P0A552
B	-10	HIS	-	EXPRESSION TAG	UNP P0A552
B	-9	SER	-	CLONING ARTIFACT	UNP P0A552
B	-8	SER	-	CLONING ARTIFACT	UNP P0A552
B	-7	GLY	-	CLONING ARTIFACT	UNP P0A552
B	-6	LEU	-	CLONING ARTIFACT	UNP P0A552
B	-5	VAL	-	CLONING ARTIFACT	UNP P0A552
B	-4	PRO	-	CLONING ARTIFACT	UNP P0A552
B	-3	ARG	-	CLONING ARTIFACT	UNP P0A552
B	-2	GLY	-	CLONING ARTIFACT	UNP P0A552
B	-1	SER	-	CLONING ARTIFACT	UNP P0A552
B	0	HIS	-	CLONING ARTIFACT	UNP P0A552
C	-19	MET	-	CLONING ARTIFACT	UNP P0A552
C	-18	GLY	-	CLONING ARTIFACT	UNP P0A552
C	-17	SER	-	CLONING ARTIFACT	UNP P0A552
C	-16	SER	-	CLONING ARTIFACT	UNP P0A552
C	-15	HIS	-	EXPRESSION TAG	UNP P0A552
C	-14	HIS	-	EXPRESSION TAG	UNP P0A552
C	-13	HIS	-	EXPRESSION TAG	UNP P0A552
C	-12	HIS	-	EXPRESSION TAG	UNP P0A552
C	-11	HIS	-	EXPRESSION TAG	UNP P0A552
C	-10	HIS	-	EXPRESSION TAG	UNP P0A552
C	-9	SER	-	CLONING ARTIFACT	UNP P0A552
C	-8	SER	-	CLONING ARTIFACT	UNP P0A552
C	-7	GLY	-	CLONING ARTIFACT	UNP P0A552
C	-6	LEU	-	CLONING ARTIFACT	UNP P0A552
C	-5	VAL	-	CLONING ARTIFACT	UNP P0A552
C	-4	PRO	-	CLONING ARTIFACT	UNP P0A552
C	-3	ARG	-	CLONING ARTIFACT	UNP P0A552
C	-2	GLY	-	CLONING ARTIFACT	UNP P0A552
C	-1	SER	-	CLONING ARTIFACT	UNP P0A552
C	0	HIS	-	CLONING ARTIFACT	UNP P0A552

- Molecule 2 is NITRATE ION (three-letter code: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	N	O		0	0
			4	1	3			

- Molecule 3 is DEOXYURIDINE-5'-TRIPHOSPHATE (three-letter code: DUT) (formula: C₉H₁₅N₂O₁₄P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	N	O	P	0	0
			28	9	2	14	3		
3	A	1	Total	C	N	O	P	0	0
			28	9	2	14	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	N	O	P	0	1
			56	18	4	28	6		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	N	O	0	0
			8	4	1	3		
4	A	1	Total	C	N	O	0	0
			8	4	1	3		
4	B	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	68	Total	O	0	0
			68	68		
5	B	73	Total	O	0	0
			73	73		
5	C	79	Total	O	0	0
			79	79		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase



- Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase



- Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.59Å 77.70Å 94.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.10 49.84 – 2.05	Depositor EDS
% Data completeness (in resolution range)	98.9 (50.00-2.10) 98.6 (49.84-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.99 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.165 , 0.207 0.180 , 0.216	Depositor DCC
R_{free} test set	2527 reflections (10.95%)	DCC
Wilson B-factor (Å ²)	25.0	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3288	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, DUT, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.91	0/979	1.11	5/1336 (0.4%)
1	B	0.90	0/1027	1.10	4/1398 (0.3%)
1	C	0.90	0/966	1.12	6/1319 (0.5%)
All	All	0.90	0/2972	1.11	15/4053 (0.4%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	46	ARG	NE-CZ-NH1	8.69	124.65	120.30
1	C	98	ASP	CB-CG-OD2	7.50	125.05	118.30
1	A	41	ARG	NE-CZ-NH1	6.77	123.69	120.30
1	B	70	ARG	NE-CZ-NH2	-6.66	116.97	120.30
1	B	22	ASP	CB-CG-OD1	-6.17	112.75	118.30
1	C	28	ASP	CB-CG-OD2	6.04	123.73	118.30
1	A	42	ARG	NE-CZ-NH1	5.83	123.22	120.30
1	C	19	ARG	NE-CZ-NH2	-5.70	117.45	120.30
1	C	19	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	B	9	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	A	118	ARG	NE-CZ-NH1	5.35	122.97	120.30
1	C	11	ASP	CB-CG-OD2	5.35	123.11	118.30
1	A	41	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	19	ARG	NE-CZ-NH1	5.23	122.91	120.30
1	B	19	ARG	NE-CZ-NH1	5.03	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	964	0	994	2	0
1	B	1013	0	1047	5	0
1	C	951	0	988	1	0
2	C	4	0	0	0	0
3	A	28	0	11	0	0
3	C	84	0	33	3	0
4	A	8	0	12	3	0
4	B	8	0	12	0	0
4	C	8	0	12	1	0
5	A	68	0	0	0	0
5	B	73	0	0	0	0
5	C	79	0	0	0	0
All	All	3288	0	3109	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:LEU:HD13	4:A:2171:TRS:H21	1.86	0.56
4:A:2171:TRS:O1	1:B:60:LEU:HD13	2.08	0.54
1:B:113:GLN:NE2	3:C:3170[B]:DUT:O2G	2.44	0.50
1:B:64:ARG:HA	3:C:3170[A]:DUT:O2B	2.15	0.47
1:C:74:SER:OG	4:C:3171:TRS:H31	2.16	0.46
1:B:28:ASP:OD2	1:B:110:ARG:NH1	2.48	0.42
1:A:60:LEU:HD13	4:A:2171:TRS:C2	2.48	0.41
3:C:3170[B]:DUT:H5'1	3:C:3170[B]:DUT:O3G	2.21	0.41
1:B:30:TYR:O	1:B:47:THR:HA	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/174 (74%)	127 (98%)	2 (2%)	0	100	100
1	B	135/174 (78%)	130 (96%)	5 (4%)	0	100	100
1	C	127/174 (73%)	127 (100%)	0	0	100	100
All	All	391/522 (75%)	384 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	103/134 (77%)	103 (100%)	0	100	100
1	B	107/134 (80%)	106 (99%)	1 (1%)	82	87
1	C	102/134 (76%)	101 (99%)	1 (1%)	80	85
All	All	312/402 (78%)	310 (99%)	2 (1%)	89	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	131	ASP
1	C	2	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	DUT	A	2170	-	22,29,29	1.26	3 (13%)	24,45,45	2.50	11 (45%)
4	TRS	A	2171	-	7,7,7	0.88	0	9,9,9	2.50	2 (22%)
4	TRS	B	155	-	7,7,7	0.81	0	9,9,9	2.76	5 (55%)
3	DUT	C	1170	-	22,29,29	1.08	2 (9%)	24,45,45	2.64	8 (33%)
2	NO3	C	155	-	1,3,3	2.99	1 (100%)	0,3,3	0.00	-
3	DUT	C	3170[A]	-	22,29,29	0.98	2 (9%)	24,45,45	2.47	7 (29%)
3	DUT	C	3170[B]	-	22,29,29	0.83	1 (4%)	24,45,45	2.86	8 (33%)
4	TRS	C	3171	-	7,7,7	1.08	0	9,9,9	2.92	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DUT	A	2170	-	-	0/18/34/34	0/2/2/2
4	TRS	A	2171	-	-	0/9/9/9	0/0/0/0
4	TRS	B	155	-	-	0/9/9/9	0/0/0/0
3	DUT	C	1170	-	-	0/18/34/34	0/2/2/2
2	NO3	C	155	-	-	0/0/0/0	0/0/0/0
3	DUT	C	3170[A]	-	-	0/18/34/34	0/2/2/2
3	DUT	C	3170[B]	-	-	0/18/34/34	0/2/2/2
4	TRS	C	3171	-	-	0/9/9/9	0/0/0/0

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	3170[B]	DUT	O4-C4	2.35	1.30	1.24
3	A	2170	DUT	PG-O3B	2.36	1.63	1.60
3	A	2170	DUT	C4-N3	2.38	1.37	1.33
3	C	1170	DUT	O4-C4	2.47	1.30	1.24
3	C	3170[A]	DUT	PG-O3B	2.48	1.64	1.60
2	C	155	NO3	O1-N	2.99	1.35	1.23
3	C	1170	DUT	PG-O3B	3.03	1.65	1.60
3	C	3170[A]	DUT	O4-C4	3.05	1.32	1.24
3	A	2170	DUT	O4-C4	3.68	1.33	1.24

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1170	DUT	C2'-C1'-N1	-4.62	103.31	114.23
4	C	3171	TRS	C2-C-N	-4.62	97.91	107.73
4	B	155	TRS	C2-C-N	-4.61	97.93	107.73
3	C	3170[B]	DUT	C2'-C1'-N1	-4.44	103.76	114.23
3	C	3170[A]	DUT	C2'-C1'-N1	-3.84	105.15	114.23
3	A	2170	DUT	C2'-C1'-N1	-3.70	105.49	114.23
4	A	2171	TRS	C3-C-C1	-2.95	102.69	111.06
4	C	3171	TRS	C3-C-N	-2.64	102.11	107.73
3	C	3170[B]	DUT	O4'-C1'-C2'	-2.36	101.73	106.25
3	A	2170	DUT	O4'-C1'-C2'	-2.33	101.78	106.25
3	C	3170[A]	DUT	C6-N1-C2	-2.29	117.57	121.28
3	A	2170	DUT	C6-N1-C2	-2.28	117.58	121.28
3	C	3170[A]	DUT	C4'-O4'-C1'	2.11	114.57	109.42
4	B	155	TRS	C3-C-N	2.13	112.25	107.73
3	C	1170	DUT	O3G-PG-O2G	2.14	116.23	107.61
3	A	2170	DUT	O3'-C3'-C4'	2.19	118.77	110.13
3	C	1170	DUT	O3'-C3'-C4'	2.20	118.81	110.13
3	A	2170	DUT	O3G-PG-O2G	2.28	116.80	107.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	155	TRS	O2-C2-C	2.33	117.22	110.47
3	A	2170	DUT	O5'-C5'-C4'	2.39	117.47	109.00
3	C	1170	DUT	C4'-O4'-C1'	2.41	115.33	109.42
3	C	3170[A]	DUT	O3'-C3'-C4'	2.49	119.94	110.13
3	C	3170[B]	DUT	O2A-PA-O1A	2.59	125.71	112.28
3	A	2170	DUT	C4'-O4'-C1'	2.61	115.80	109.42
3	A	2170	DUT	O3'-C3'-C2'	2.83	121.13	110.83
3	C	1170	DUT	C3'-C2'-C1'	3.00	110.17	102.48
3	C	3170[B]	DUT	O3'-C3'-C4'	3.19	122.73	110.13
3	C	3170[B]	DUT	O3'-C3'-C2'	3.20	122.47	110.83
3	C	3170[A]	DUT	C3'-C2'-C1'	3.25	110.81	102.48
3	C	1170	DUT	O3'-C3'-C2'	3.46	123.41	110.83
4	B	155	TRS	C1-C-N	3.81	115.83	107.73
3	A	2170	DUT	C3'-C2'-C1'	4.02	112.78	102.48
4	B	155	TRS	O1-C1-C	4.14	122.45	110.47
3	C	3170[B]	DUT	C3'-C2'-C1'	4.43	113.84	102.48
3	C	3170[A]	DUT	O4'-C1'-N1	4.95	116.13	107.78
3	C	1170	DUT	O4'-C1'-N1	5.06	116.30	107.78
3	A	2170	DUT	C4-N3-C2	5.57	118.92	114.13
4	A	2171	TRS	C1-C-N	5.96	120.39	107.73
3	C	3170[B]	DUT	C4-N3-C2	6.12	119.39	114.13
3	A	2170	DUT	O4'-C1'-N1	6.14	118.13	107.78
4	C	3171	TRS	C1-C-N	6.53	121.61	107.73
3	C	3170[B]	DUT	O4'-C1'-N1	7.51	120.44	107.78
3	C	1170	DUT	C4-N3-C2	7.77	120.81	114.13
3	C	3170[A]	DUT	C4-N3-C2	7.79	120.83	114.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2171	TRS	3	0
3	C	3170[A]	DUT	1	0
3	C	3170[B]	DUT	2	0
4	C	3171	TRS	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	131/174 (75%)	0.08	2 (1%) 74 77	13, 18, 31, 53	0
1	B	138/174 (79%)	0.27	5 (3%) 43 50	13, 19, 45, 60	0
1	C	129/174 (74%)	0.07	2 (1%) 72 76	11, 16, 29, 41	0
All	All	398/522 (76%)	0.14	9 (2%) 61 66	11, 18, 32, 60	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	132	GLU	4.9
1	B	0	HIS	3.7
1	B	140	ARG	3.2
1	A	20	ALA	3.0
1	C	2	SER	2.8
1	B	132	GLU	2.5
1	B	131	ASP	2.4
1	B	136	ALA	2.3
1	C	20	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	TRS	C	3171	8/8	0.63	0.45	17.99	46,47,48,48	0
4	TRS	B	155	8/8	0.70	0.34	14.99	47,50,51,51	0
4	TRS	A	2171	8/8	0.82	0.31	14.50	27,32,33,34	0
3	DUT	C	3170[A]	28/28	0.80	0.31	6.40	18,30,53,54	28
3	DUT	C	3170[B]	28/28	0.80	0.31	6.40	13,32,70,71	28
3	DUT	A	2170	28/28	0.81	0.20	4.37	21,37,66,68	0
2	NO3	C	155	4/4	0.97	0.19	3.95	29,29,31,33	0
3	DUT	C	1170	28/28	0.81	0.23	1.62	22,30,61,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.