



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 12, 2017 – 07:29 pm GMT

PDB ID : 1SN9
Title : An Oligomeric Domain-Swapped Beta-Beta-Alpha Mini-Protein
Authors : Ali, M.H.; Peisach, E.; Allen, K.N.; Imperiali, B.
Deposited on : 2004-03-10
Resolution : 1.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1201 (1.24-1.16)
Ramachandran outliers	110173	1148 (1.24-1.16)
Sidechain outliers	110143	1147 (1.24-1.16)
RSRZ outliers	101464	1132 (1.24-1.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	23	 13% 52% 43% .
1	B	23	 13% 70% 26% .
1	C	23	 4% 78% 17% .
1	D	23	 4% 70% 26% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 931 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tetrameric beta-beta-alpha mini-protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	23	Total	C	N	O	0	4	1
			202	130	37	35			
1	B	23	Total	C	N	O	0	5	1
			209	134	38	37			
1	C	23	Total	C	N	O	0	3	1
			209	133	40	36			
1	D	23	Total	C	N	O	0	1	1
			193	124	35	34			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	32	Total	O	0	0
			32	32		
2	B	30	Total	O	0	0
			30	30		
2	C	27	Total	O	0	0
			27	27		
2	D	29	Total	O	0	0
			29	29		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: tetrameric beta-beta-alpha mini-protein

Chain A: 





- Molecule 1: tetrameric beta-beta-alpha mini-protein

Chain B: 



- Molecule 1: tetrameric beta-beta-alpha mini-protein

Chain C: 



- Molecule 1: tetrameric beta-beta-alpha mini-protein

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	38.94Å 56.17Å 32.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	99.00 – 1.20 24.87 – 1.20	Depositor EDS
% Data completeness (in resolution range)	86.5 (99.00-1.20) 86.5 (24.87-1.20)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.78 (at 1.20Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.188 , 0.285 0.195 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	12.3	Xtriage
Anisotropy	0.233	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 94.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	931	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DAL, DBZ, DPR, ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/174	1.29	1/225 (0.4%)
1	B	4.00	2/183 (1.1%)	1.43	2/237 (0.8%)
1	C	0.52	0/183	1.33	0/238
1	D	0.49	0/163	1.39	4/212 (1.9%)
All	All	2.08	2/703 (0.3%)	1.36	7/912 (0.8%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	215[A]	LEU	CA-CB	37.96	2.41	1.53
1	B	215[B]	LEU	CA-CB	37.96	2.41	1.53

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	215[A]	LEU	N-CA-CB	-8.10	94.19	110.40
1	B	215[B]	LEU	N-CA-CB	-8.10	94.19	110.40
1	D	417[A]	ARG	NE-CZ-NH2	-6.97	116.81	120.30
1	D	417[B]	ARG	NE-CZ-NH2	-6.97	116.81	120.30
1	D	402	ARG	CD-NE-CZ	6.38	132.53	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	202	0	195	13	0
1	B	209	0	201	13	0
1	C	209	0	201	6	0
1	D	193	0	186	6	0
2	A	32	0	0	5	0
2	B	30	0	0	3	0
2	C	27	0	0	1	0
2	D	29	0	0	3	0
All	All	931	0	783	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 31 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112[B]:LEU:HD22	1:B:215[B]:LEU:HD22	1.66	0.77
1:D:413:ALA:O	1:D:417[B]:ARG:HG3	1.89	0.73
1:B:215[B]:LEU:HD12	2:B:4:HOH:O	1.97	0.64
2:A:150:HOH:O	1:B:215[B]:LEU:HD11	2.03	0.58
1:A:114:LYS:O	1:A:118:GLN:HG3	2.04	0.58

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	21/23 (91%)	21 (100%)	0	0	100	100
1	B	22/23 (96%)	20 (91%)	2 (9%)	0	100	100
1	C	21/23 (91%)	20 (95%)	1 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	19/23 (83%)	19 (100%)	0	0	100	100
All	All	83/92 (90%)	80 (96%)	3 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	17/15 (113%)	15 (88%)	2 (12%)	6	0
1	B	18/15 (120%)	16 (89%)	2 (11%)	7	0
1	C	18/15 (120%)	16 (89%)	2 (11%)	7	0
1	D	16/15 (107%)	16 (100%)	0	100	100
All	All	69/60 (115%)	63 (91%)	6 (9%)	19	0

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	215[A]	LEU
1	C	317[B]	ARG
1	B	215[B]	LEU
1	A	110	ASP
1	C	317[A]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	DPR	A	104	1	6,7,8	0.94	0	7,8,10	1.25	1 (14%)
1	DAL	A	109	1	4,4,5	0.90	0	1,4,6	0.10	0
1	DBZ	A	120	1	14,14,15	1.38	1 (7%)	14,17,19	1.27	1 (7%)
1	DPR	B	204	1	6,7,8	0.75	0	7,8,10	1.46	2 (28%)
1	DAL	B	209	1	4,4,5	1.24	1 (25%)	1,4,6	0.04	0
1	DBZ	B	220	1	14,14,15	1.38	1 (7%)	14,17,19	1.30	1 (7%)
1	DPR	C	304	1	6,7,8	1.30	1 (16%)	7,8,10	1.07	0
1	DAL	C	309	1	4,4,5	1.49	1 (25%)	1,4,6	1.12	0
1	DBZ	C	320	1	14,14,15	1.32	1 (7%)	14,17,19	1.56	2 (14%)
1	DPR	D	404	1	6,7,8	1.18	1 (16%)	7,8,10	1.04	1 (14%)
1	DAL	D	409	1	4,4,5	1.36	1 (25%)	1,4,6	0.18	0
1	DBZ	D	420	1	14,14,15	1.23	1 (7%)	14,17,19	1.51	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DPR	A	104	1	-	0/0/9/11	0/1/1/1
1	DAL	A	109	1	-	0/0/2/4	0/0/0/0
1	DBZ	A	120	1	-	0/11/11/13	0/1/1/1
1	DPR	B	204	1	-	0/0/9/11	0/1/1/1
1	DAL	B	209	1	-	0/0/2/4	0/0/0/0
1	DBZ	B	220	1	-	0/11/11/13	0/1/1/1
1	DPR	C	304	1	-	0/0/9/11	0/1/1/1
1	DAL	C	309	1	-	0/0/2/4	0/0/0/0
1	DBZ	C	320	1	-	0/11/11/13	0/1/1/1
1	DPR	D	404	1	-	0/0/9/11	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DAL	D	409	1	-	0/0/2/4	0/0/0/0
1	DBZ	D	420	1	-	0/11/11/13	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	220	DBZ	O-C	-4.64	1.22	1.42
1	A	120	DBZ	O-C	-4.61	1.22	1.42
1	C	320	DBZ	O-C	-4.37	1.24	1.42
1	D	420	DBZ	O-C	-4.03	1.25	1.42
1	B	209	DAL	CA-C	2.31	1.53	1.50

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	220	DBZ	OD2-CD2-C1	-2.75	116.07	120.94
1	D	420	DBZ	CA-CB-NG	-2.71	104.50	111.89
1	C	320	DBZ	OD2-CD2-C1	-2.54	116.43	120.94
1	A	104	DPR	O-C-CA	-2.38	119.59	125.15
1	B	204	DPR	O-C-CA	-2.24	119.93	125.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	304	DPR	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	18/23 (78%)	0.25	0 100 100	14, 22, 31, 32	0
1	B	18/23 (78%)	0.81	3 (16%) 2 3	15, 22, 30, 32	0
1	C	18/23 (78%)	0.34	1 (5%) 25 23	14, 20, 29, 44	0
1	D	18/23 (78%)	0.37	1 (5%) 25 23	13, 20, 37, 40	0
All	All	72/92 (78%)	0.44	5 (6%) 18 17	13, 22, 32, 44	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	221[A]	GLY	3.4
1	B	201	TYR	3.2
1	D	403	ILE	2.7
1	C	321	GLY	2.4
1	B	215[A]	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	DBZ	D	420	14/15	0.95	0.06	-	16,22,27,29	0
1	DAL	B	209	5/6	0.94	0.08	-	15,16,17,19	0
1	DPR	C	304	7/8	0.94	0.09	-	22,25,29,35	0
1	DPR	A	104	7/8	0.90	0.11	-	18,21,23,23	0
1	DAL	A	109	5/6	0.94	0.08	-	14,15,15,15	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	DBZ	A	120	14/15	0.94	0.07	-	18,20,22,26	0
1	DAL	C	309	5/6	0.97	0.06	-	14,14,17,18	0
1	DBZ	B	220	14/15	0.89	0.09	-	24,28,32,43	0
1	DPR	B	204	7/8	0.88	0.12	-	20,22,26,27	0
1	DAL	D	409	5/6	0.96	0.08	-	14,16,17,22	0
1	DPR	D	404	7/8	0.83	0.14	-	33,36,40,43	0
1	DBZ	C	320	14/15	0.90	0.10	-	16,19,28,36	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.