



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 2, 2017 – 06:38 AM EDT

PDB ID : 1SO0  
Title : Crystal structure of human galactose mutarotase complexed with galactose  
Authors : Thoden, J.B.; Timson, D.J.; Reece, R.J.; Holden, H.M.  
Deposited on : unknown  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	 65% 29% 5% .
1	B	344	 59% 32% 7% ..
1	C	344	 58% 33% 8% .
1	D	344	 59% 32% 8% .

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 11200 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called aldose 1-epimerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	344	Total 2698	C 1718	N 469	O 506	S 5	0	2	0
1	B	342	Total 2679	C 1707	N 465	O 502	S 5	0	1	0
1	C	344	Total 2688	C 1712	N 469	O 502	S 5	0	0	0
1	D	342	Total 2681	C 1708	N 468	O 500	S 5	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	CLONING ARTIFACT	UNP Q96C23
A	0	HIS	-	CLONING ARTIFACT	UNP Q96C23
B	-1	GLY	-	CLONING ARTIFACT	UNP Q96C23
B	0	HIS	-	CLONING ARTIFACT	UNP Q96C23
C	-1	GLY	-	CLONING ARTIFACT	UNP Q96C23
C	0	HIS	-	CLONING ARTIFACT	UNP Q96C23
D	-1	GLY	-	CLONING ARTIFACT	UNP Q96C23
D	0	HIS	-	CLONING ARTIFACT	UNP Q96C23

- Molecule 2 is BETA-D-GALACTOSE (three-letter code: GAL) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 12 6 6	0	0
2	B	1	Total C O 12 6 6	0	0
2	C	1	Total C O 12 6 6	0	0
2	D	1	Total C O 12 6 6	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	118	Total O 118 118	0	0
3	B	99	Total O 99 99	0	0
3	C	113	Total O 113 113	0	0
3	D	76	Total O 76 76	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: aldose 1-epimerase




- Molecule 1: aldose 1-epimerase




- Molecule 1: aldose 1-epimerase





- Molecule 1: aldose 1-epimerase

Chain D: 59% 32% 8%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.00Å 68.70Å 98.90Å 107.70° 98.40° 102.70°	Depositor
Resolution (Å)	30.00 – 2.30 57.92 – 2.24	Depositor EDS
% Data completeness (in resolution range)	94.7 (30.00-2.30) 81.1 (57.92-2.24)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.26 (at 2.25Å)	Xtrriage
Refinement program	TNT	Depositor
R, $R_{free}$	0.167 , 0.198 0.164 , 0.165	Depositor DCC
$R_{free}$ test set	6054 reflections (11.15%)	DCC
Wilson B-factor (Å <sup>2</sup> )	21.9	Xtrriage
Anisotropy	0.178	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 69.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11200	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.01	18/2778 (0.6%)	1.48	37/3775 (1.0%)
1	B	0.99	19/2754 (0.7%)	1.44	40/3742 (1.1%)
1	C	1.02	20/2760 (0.7%)	1.43	34/3750 (0.9%)
1	D	1.00	16/2756 (0.6%)	1.42	36/3744 (1.0%)
All	All	1.01	73/11048 (0.7%)	1.44	147/15011 (1.0%)

The worst 5 of 73 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	253	GLU	CD-OE2	9.76	1.36	1.25
1	D	61	GLU	CD-OE2	9.55	1.36	1.25
1	B	61	GLU	CD-OE2	9.53	1.36	1.25
1	C	90	VAL	C-O	8.33	1.39	1.23
1	D	102	GLU	CD-OE2	8.31	1.34	1.25

The worst 5 of 147 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	90	VAL	O-C-N	10.92	140.17	122.70
1	A	54	ASP	CB-CG-OD2	-10.68	108.69	118.30
1	C	313	ASP	CB-CG-OD2	-10.39	108.95	118.30
1	A	236	ASP	CB-CG-OD2	-9.39	109.85	118.30
1	A	47	ASP	CB-CG-OD2	-9.12	110.09	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2698	0	2614	75	0
1	B	2679	0	2602	90	0
1	C	2688	0	2610	110	0
1	D	2681	0	2609	98	0
2	A	12	0	12	0	0
2	B	12	0	12	0	0
2	C	12	0	12	0	0
2	D	12	0	12	0	0
3	A	118	0	0	3	0
3	B	99	0	0	2	0
3	C	113	0	0	7	0
3	D	76	0	0	5	0
All	All	11200	0	10483	368	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 368 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:76:ILE:HD11	1:A:176:HIS:HB2	1.19	1.14
1:C:1:MET:HE2	1:C:125:ASN:HD22	1.11	1.08
1:D:240:ASN:HD21	1:D:300:LYS:NZ	1.50	1.07
1:C:90:VAL:C	3:C:791:HOH:O	1.91	1.07
1:B:184:GLN:HE22	1:B:293:LYS:HB2	1.24	1.02

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	344/344 (100%)	323 (94%)	19 (6%)	2 (1%)	28	34
1	B	341/344 (99%)	318 (93%)	22 (6%)	1 (0%)	44	55
1	C	342/344 (99%)	319 (93%)	21 (6%)	2 (1%)	28	34
1	D	341/344 (99%)	315 (92%)	24 (7%)	2 (1%)	28	34
All	All	1368/1376 (99%)	1275 (93%)	86 (6%)	7 (0%)	32	39

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	73	GLY
1	D	73	GLY
1	A	81	ASN
1	D	243	ASP
1	B	70	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/285 (101%)	266 (93%)	21 (7%)	16	21
1	B	285/285 (100%)	260 (91%)	25 (9%)	12	14
1	C	285/285 (100%)	262 (92%)	23 (8%)	14	17
1	D	285/285 (100%)	262 (92%)	23 (8%)	14	17
All	All	1142/1140 (100%)	1050 (92%)	92 (8%)	14	17

5 of 92 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	301	HIS
1	C	76	ILE
1	D	252	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	315	VAL
1	C	29	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	275	GLN
1	B	309	GLN
1	C	301	HIS
1	B	245	ASN
1	B	261	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GAL	A	500	-	12,12,12	0.58	0	17,17,17	1.27	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GAL	B	600	-	12,12,12	0.49	0	17,17,17	1.40	3 (17%)
2	GAL	C	700	-	12,12,12	0.48	0	17,17,17	1.43	2 (11%)
2	GAL	D	800	-	12,12,12	0.44	0	17,17,17	1.48	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	A	500	-	-	0/2/22/22	0/1/1/1
2	GAL	B	600	-	-	0/2/22/22	0/1/1/1
2	GAL	C	700	-	-	0/2/22/22	0/1/1/1
2	GAL	D	800	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	700	GAL	C4-C3-C2	-3.98	103.81	110.84
2	D	800	GAL	C3-C4-C5	-3.06	104.82	110.22
2	B	600	GAL	C3-C4-C5	-2.53	105.77	110.22
2	B	600	GAL	C1-C2-C3	-2.51	106.12	110.65
2	A	500	GAL	C3-C4-C5	-2.39	106.01	110.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	344/344 (100%)	-0.55	1 (0%) 93   96	18, 29, 54, 87	0
1	B	342/344 (99%)	-0.48	5 (1%) 74   78	20, 32, 60, 86	0
1	C	344/344 (100%)	-0.50	2 (0%) 89   92	19, 32, 60, 89	0
1	D	342/344 (99%)	-0.48	0 100   100	20, 34, 63, 88	0
All	All	1372/1376 (99%)	-0.50	8 (0%) 89   92	18, 31, 60, 89	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	14	SER	2.8
1	B	251	SER	2.8
1	B	249	LYS	2.5
1	B	1	MET	2.4
1	A	249	LYS	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GAL	B	600	12/12	0.97	0.12	1.12	17,33,49,89	0
2	GAL	D	800	12/12	0.96	0.11	0.83	18,35,71,100	0
2	GAL	A	500	12/12	0.96	0.11	-0.06	17,33,45,77	0
2	GAL	C	700	12/12	0.97	0.10	-0.56	21,38,51,100	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.