



Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 07:55 pm GMT

PDB ID : 1SSF
Title : Solution structure of the mouse 53BP1 fragment (residues 1463-1617)
Authors : Charier, G.; Couprie, J.; Alpha-Bazin, B.; Meyer, V.; Quemeneur, E.; Guerois, R.; Callebaut, I.; Gilquin, B.; Zinn-Justin, S.
Deposited on : 2004-03-24

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 86%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	156	

2 Ensemble composition and analysis

This entry contains 10 models. Model 9 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy conformer*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:11-A:71, A:76-A:84, A:92-A:127 (106)	0.40	9

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 9, 10
Single-model clusters	8

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1953 atoms, of which 966 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Transformation related protein 53 binding protein 1.

Mol	Chain	Residues	Atoms						Trace
1	A	123	Total	C	H	N	O	S	0
			1953	631	966	164	189	3	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	CLONING ARTIFACT	UNP Q91YC9

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Transformation related protein 53 binding protein 1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Transformation related protein 53 binding protein 1




4.2.2 Score per residue for model 2

- Molecule 1: Transformation related protein 53 binding protein 1




4.2.3 Score per residue for model 3

- Molecule 1: Transformation related protein 53 binding protein 1



4.2.4 Score per residue for model 4


- Molecule 1: Transformation related protein 53 binding protein 1



4.2.5 Score per residue for model 5


- Molecule 1: Transformation related protein 53 binding protein 1






4.2.6 Score per residue for model 6

- Molecule 1: Transformation related protein 53 binding protein 1




4.2.7 Score per residue for model 7

- Molecule 1: Transformation related protein 53 binding protein 1




4.2.8 Score per residue for model 8

- Molecule 1: Transformation related protein 53 binding protein 1




4.2.9 Score per residue for model 9 (medoid)

- Molecule 1: Transformation related protein 53 binding protein 1



4.2.10 Score per residue for model 10

- Molecule 1: Transformation related protein 53 binding protein 1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry, simulated annealing*.

Of the 1000 calculated structures, 10 were deposited, based on the following criterion: *back calculated data agree with experimental NOESY spectrum, structures with acceptable covalent geometry, structures with favorable non-bond energy, structures with the least restraint violations, structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	1.0
CNS	structure solution	1.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5878
Number of chemical shift lists	2
Total number of shifts	1861
Number of shifts mapped to atoms	1861
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	86%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.07±0.01	0±0/859 (0.0±0.0%)	1.83±0.06	21±4/1158 (1.8±0.3%)
All	All	1.07	0/8590 (0.0%)	1.83	210/11580 (1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	4.5±1.1
All	All	0	45

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	92	TYR	CB-CG-CD2	-12.57	113.46	121.00	3	3
1	A	93	TYR	CB-CG-CD2	-12.03	113.78	121.00	7	2
1	A	92	TYR	CB-CG-CD1	11.71	128.02	121.00	3	1
1	A	14	VAL	CA-CB-CG2	9.51	125.16	110.90	3	5
1	A	113	LEU	N-CA-CB	9.42	129.25	110.40	6	9
1	A	41	LEU	CB-CA-C	9.09	127.47	110.20	5	10
1	A	93	TYR	CB-CG-CD1	8.50	126.10	121.00	7	3
1	A	42	PHE	CB-CG-CD1	8.34	126.64	120.80	2	7
1	A	62	PRO	N-CA-CB	-8.08	93.61	103.30	5	1
1	A	12	LEU	CB-CG-CD2	-7.80	97.73	111.00	9	6
1	A	56	LEU	CB-CG-CD2	7.76	124.19	111.00	10	1
1	A	76	PHE	CB-CG-CD1	7.74	126.22	120.80	2	7
1	A	26	SER	N-CA-CB	-7.74	98.89	110.50	6	3
1	A	92	TYR	CA-CB-CG	7.59	127.82	113.40	3	2
1	A	76	PHE	CB-CG-CD2	-7.35	115.66	120.80	2	6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	94	SER	N-CA-CB	7.30	121.46	110.50	1	1
1	A	62	PRO	C-N-CA	7.20	139.70	121.70	6	7
1	A	12	LEU	CA-CB-CG	7.13	131.71	115.30	9	7
1	A	57	LEU	CA-CB-CG	7.04	131.50	115.30	9	4
1	A	12	LEU	CB-CA-C	7.03	123.55	110.20	1	3
1	A	104	TYR	CB-CG-CD2	-7.02	116.79	121.00	2	2
1	A	24	PHE	CB-CG-CD2	6.88	125.62	120.80	6	4
1	A	42	PHE	CB-CG-CD2	-6.86	116.00	120.80	2	4
1	A	61	ILE	CB-CA-C	6.53	124.66	111.60	4	3
1	A	56	LEU	CB-CG-CD1	-6.52	99.92	111.00	9	3
1	A	70	LEU	C-N-CA	6.39	137.68	121.70	2	7
1	A	14	VAL	CA-CB-CG1	6.38	120.47	110.90	7	1
1	A	63	LEU	N-CA-C	6.38	128.21	111.00	3	4
1	A	104	TYR	CB-CG-CD1	6.38	124.83	121.00	2	1
1	A	18	TRP	CB-CG-CD2	-6.35	118.35	126.60	5	8
1	A	26	SER	C-N-CA	6.29	135.51	122.30	4	3
1	A	58	CYS	CB-CA-C	6.23	122.86	110.40	6	3
1	A	123	TYR	CB-CG-CD2	-6.23	117.26	121.00	4	3
1	A	40	LEU	CB-CG-CD1	6.23	121.59	111.00	4	2
1	A	118	ARG	C-N-CA	6.13	137.04	121.70	1	1
1	A	24	PHE	C-N-CA	6.13	137.03	121.70	5	4
1	A	64	ASP	N-CA-CB	-6.13	99.57	110.60	5	6
1	A	60	PRO	N-CA-CB	6.02	110.52	103.30	5	1
1	A	63	LEU	CB-CG-CD1	6.00	121.19	111.00	6	2
1	A	15	VAL	CA-CB-CG1	-5.96	101.95	110.90	2	2
1	A	53	LYS	C-N-CA	5.96	136.60	121.70	3	4
1	A	125	LEU	CD1-CG-CD2	5.94	128.31	110.50	5	1
1	A	68	THR	CA-CB-CG2	-5.92	104.11	112.40	9	4
1	A	49	ASP	N-CA-CB	5.89	121.21	110.60	5	3
1	A	67	VAL	N-CA-CB	5.87	124.41	111.50	3	2
1	A	63	LEU	CB-CG-CD2	-5.79	101.15	111.00	10	2
1	A	123	TYR	CB-CG-CD1	5.76	124.45	121.00	4	2
1	A	58	CYS	N-CA-CB	-5.73	100.28	110.60	7	2
1	A	63	LEU	C-N-CA	5.73	136.02	121.70	6	2
1	A	18	TRP	CB-CG-CD1	5.71	134.42	127.00	5	3
1	A	66	GLU	C-N-CA	5.70	135.96	121.70	1	3
1	A	24	PHE	N-CA-CB	5.68	120.83	110.60	4	3
1	A	16	ALA	N-CA-CB	-5.66	102.18	110.10	5	1
1	A	114	GLU	N-CA-CB	5.63	120.74	110.60	8	1
1	A	49	ASP	N-CA-C	-5.63	95.80	111.00	5	1
1	A	11	GLY	C-N-CA	5.61	135.72	121.70	3	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	81	VAL	CG1-CB-CG2	-5.60	101.94	110.90	3	1
1	A	71	SER	N-CA-C	-5.59	95.90	111.00	5	2
1	A	65	THR	N-CA-CB	5.47	120.69	110.30	1	1
1	A	12	LEU	CB-CG-CD1	5.39	120.17	111.00	3	2
1	A	106	ARG	N-CA-CB	-5.38	100.92	110.60	8	1
1	A	41	LEU	N-CA-CB	-5.32	99.76	110.40	3	2
1	A	24	PHE	CB-CG-CD1	-5.30	117.09	120.80	6	2
1	A	109	VAL	CA-CB-CG1	-5.27	102.99	110.90	7	1
1	A	115	GLN	C-N-CA	5.26	133.35	122.30	4	1
1	A	23	TYR	N-CA-CB	5.22	120.00	110.60	3	1
1	A	21	ASN	N-CA-CB	-5.22	101.20	110.60	7	1
1	A	56	LEU	C-N-CA	5.21	134.73	121.70	4	2
1	A	57	LEU	CB-CA-C	-5.19	100.34	110.20	2	1
1	A	111	LEU	CB-CG-CD2	5.16	119.77	111.00	8	1
1	A	125	LEU	CB-CA-C	5.13	119.95	110.20	5	1
1	A	123	TYR	CA-C-N	5.09	126.37	116.20	1	1
1	A	56	LEU	N-CA-C	5.06	124.66	111.00	8	1
1	A	65	THR	CA-CB-CG2	5.03	119.44	112.40	5	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	93	TYR	Sidechain	8
1	A	104	TYR	Sidechain	7
1	A	25	TYR	Sidechain	5
1	A	38	TYR	Sidechain	5
1	A	51	LEU	Peptide	4
1	A	92	TYR	Sidechain	4
1	A	24	PHE	Sidechain	4
1	A	46	TYR	Sidechain	4
1	A	123	TYR	Sidechain	3
1	A	23	TYR	Sidechain	1

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	840	840	840	19±3
All	All	8400	8400	8400	189

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:56:LEU:HD11	1:A:125:LEU:HD13	0.94	1.37	10	4
1:A:12:LEU:HD11	1:A:57:LEU:CB	0.82	2.05	6	2
1:A:12:LEU:HD11	1:A:57:LEU:HB2	0.81	1.50	2	4
1:A:12:LEU:HD22	1:A:57:LEU:HB2	0.80	1.52	7	1
1:A:24:PHE:CD2	1:A:111:LEU:HD13	0.78	2.13	10	7
1:A:12:LEU:HD22	1:A:13:ARG:H	0.76	1.39	1	2
1:A:12:LEU:HD11	1:A:57:LEU:HA	0.73	1.59	8	1
1:A:56:LEU:HD11	1:A:125:LEU:CD1	0.72	2.13	10	1
1:A:61:ILE:CG1	1:A:111:LEU:HD23	0.72	2.14	5	3
1:A:12:LEU:HD21	1:A:57:LEU:CB	0.71	2.15	9	2
1:A:24:PHE:CD2	1:A:111:LEU:HD11	0.70	2.21	8	1
1:A:56:LEU:HD21	1:A:125:LEU:HA	0.69	1.64	2	5
1:A:61:ILE:HG13	1:A:111:LEU:HD23	0.69	1.61	5	4
1:A:12:LEU:HD22	1:A:13:ARG:N	0.69	2.02	1	1
1:A:12:LEU:HD11	1:A:57:LEU:HB3	0.69	1.62	6	1
1:A:61:ILE:HD12	1:A:119:LEU:HD13	0.66	1.64	4	3
1:A:40:LEU:HD23	1:A:55:ILE:HG21	0.66	1.68	1	3
1:A:33:VAL:HG23	1:A:37:LYS:HB2	0.64	1.68	9	10
1:A:67:VAL:HA	1:A:115:GLN:HE22	0.62	1.53	10	2
1:A:12:LEU:HD23	1:A:29:ILE:HG21	0.61	1.71	9	1
1:A:40:LEU:CD2	1:A:55:ILE:HG21	0.61	2.26	8	3
1:A:12:LEU:HD21	1:A:57:LEU:HB3	0.60	1.73	5	3
1:A:104:TYR:HB3	1:A:108:ALA:HB3	0.60	1.73	3	1
1:A:56:LEU:HD22	1:A:125:LEU:HD13	0.59	1.74	9	2
1:A:84:HIS:CD2	1:A:93:TYR:CZ	0.59	2.90	5	2
1:A:12:LEU:HD21	1:A:57:LEU:HB2	0.58	1.74	9	1
1:A:12:LEU:HD13	1:A:29:ILE:HG13	0.57	1.75	6	1
1:A:56:LEU:CD1	1:A:125:LEU:HD13	0.57	2.24	10	1
1:A:84:HIS:CD2	1:A:93:TYR:CE2	0.56	2.92	3	4
1:A:24:PHE:CG	1:A:111:LEU:HD13	0.56	2.35	5	4
1:A:63:LEU:HA	1:A:81:VAL:HG23	0.55	1.76	5	1
1:A:61:ILE:CD1	1:A:111:LEU:HD23	0.55	2.32	3	3
1:A:52:GLY:O	1:A:56:LEU:HD12	0.55	2.02	10	1
1:A:14:VAL:HG11	1:A:29:ILE:CG2	0.55	2.32	3	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:12:LEU:HD11	1:A:57:LEU:CA	0.54	2.32	8	1
1:A:24:PHE:CD2	1:A:111:LEU:CD1	0.54	2.91	1	4
1:A:12:LEU:HD23	1:A:13:ARG:N	0.53	2.17	7	1
1:A:61:ILE:HD11	1:A:111:LEU:HD23	0.53	1.80	3	2
1:A:17:LYS:HG2	1:A:24:PHE:CG	0.51	2.40	4	7
1:A:61:ILE:HD12	1:A:115:GLN:O	0.50	2.06	5	1
1:A:24:PHE:CD2	1:A:111:LEU:HD21	0.50	2.41	7	2
1:A:41:LEU:HD21	1:A:45:GLY:HA2	0.50	1.82	4	3
1:A:24:PHE:CE2	1:A:116:GLY:HA3	0.49	2.42	2	1
1:A:61:ILE:HD13	1:A:119:LEU:CD1	0.49	2.38	3	3
1:A:95:ILE:HD12	1:A:109:VAL:HG11	0.49	1.85	1	1
1:A:12:LEU:HD22	1:A:57:LEU:CB	0.49	2.31	7	1
1:A:14:VAL:HG11	1:A:29:ILE:HG23	0.49	1.84	3	1
1:A:56:LEU:HD11	1:A:125:LEU:HA	0.48	1.84	9	2
1:A:61:ILE:HD13	1:A:119:LEU:HD13	0.48	1.85	3	3
1:A:60:PRO:HD2	1:A:93:TYR:CE2	0.48	2.44	9	1
1:A:14:VAL:HG23	1:A:27:GLY:HA3	0.48	1.86	10	1
1:A:62:PRO:CD	1:A:119:LEU:HD12	0.48	2.39	3	1
1:A:17:LYS:HG3	1:A:24:PHE:HA	0.47	1.85	8	1
1:A:41:LEU:HD23	1:A:42:PHE:O	0.47	2.08	8	2
1:A:12:LEU:HD23	1:A:29:ILE:HG13	0.47	1.87	5	1
1:A:63:LEU:CD1	1:A:63:LEU:H	0.47	2.21	6	2
1:A:14:VAL:HG11	1:A:29:ILE:CD1	0.47	2.39	6	1
1:A:15:VAL:HG21	1:A:125:LEU:HD21	0.47	1.85	1	1
1:A:61:ILE:CG2	1:A:65:THR:HG21	0.47	2.40	4	1
1:A:12:LEU:HD21	1:A:57:LEU:HA	0.47	1.85	1	1
1:A:56:LEU:HD11	1:A:125:LEU:HG	0.47	1.87	5	1
1:A:63:LEU:HD12	1:A:63:LEU:H	0.46	1.70	10	2
1:A:14:VAL:HG11	1:A:29:ILE:HG12	0.46	1.87	5	1
1:A:60:PRO:HD2	1:A:93:TYR:CE1	0.46	2.45	6	1
1:A:61:ILE:HD11	1:A:116:GLY:HA2	0.45	1.88	4	1
1:A:12:LEU:CD2	1:A:13:ARG:H	0.45	2.23	6	1
1:A:15:VAL:HG11	1:A:111:LEU:HD21	0.45	1.88	3	1
1:A:50:VAL:HG21	1:A:55:ILE:HB	0.45	1.89	8	3
1:A:95:ILE:N	1:A:95:ILE:HD13	0.45	2.26	6	1
1:A:93:TYR:HB3	1:A:109:VAL:HG11	0.45	1.89	5	1
1:A:37:LYS:HB3	1:A:51:LEU:HA	0.45	1.89	8	1
1:A:84:HIS:CG	1:A:93:TYR:CE1	0.45	3.05	6	2
1:A:95:ILE:HG13	1:A:109:VAL:HG11	0.45	1.88	7	1
1:A:61:ILE:HG12	1:A:119:LEU:HD13	0.44	1.89	6	1
1:A:93:TYR:CB	1:A:109:VAL:HG11	0.44	2.41	5	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:58:CYS:SG	1:A:125:LEU:CD2	0.44	3.06	7	1
1:A:24:PHE:CG	1:A:111:LEU:CD1	0.44	3.01	8	1
1:A:61:ILE:HD11	1:A:111:LEU:HD12	0.44	1.90	2	1
1:A:57:LEU:HD13	1:A:124:GLY:HA2	0.44	1.90	10	1
1:A:12:LEU:CD1	1:A:57:LEU:HB2	0.44	2.41	4	1
1:A:58:CYS:SG	1:A:125:LEU:HD22	0.43	2.53	2	1
1:A:12:LEU:HD13	1:A:29:ILE:CG1	0.43	2.42	6	1
1:A:58:CYS:HB3	1:A:60:PRO:HA	0.43	1.89	6	2
1:A:24:PHE:CG	1:A:111:LEU:HD11	0.43	2.47	8	1
1:A:24:PHE:CD1	1:A:24:PHE:N	0.43	2.87	7	4
1:A:65:THR:O	1:A:67:VAL:HG22	0.42	2.13	1	1
1:A:60:PRO:O	1:A:119:LEU:HD13	0.42	2.15	5	1
1:A:94:SER:HB3	1:A:103:TRP:CD2	0.42	2.49	1	1
1:A:61:ILE:CG1	1:A:111:LEU:HD13	0.42	2.44	2	1
1:A:12:LEU:HD21	1:A:57:LEU:CA	0.42	2.44	4	1
1:A:95:ILE:HG13	1:A:109:VAL:HG21	0.42	1.91	8	1
1:A:61:ILE:HD11	1:A:111:LEU:CD1	0.42	2.45	2	1
1:A:38:TYR:CE2	1:A:57:LEU:HD12	0.42	2.50	6	1
1:A:119:LEU:CD2	1:A:125:LEU:H	0.41	2.29	4	1
1:A:33:VAL:HG23	1:A:37:LYS:CB	0.41	2.44	6	1
1:A:62:PRO:HD3	1:A:119:LEU:HD12	0.41	1.91	2	1
1:A:14:VAL:HG11	1:A:29:ILE:HD11	0.41	1.92	6	1
1:A:61:ILE:HA	1:A:62:PRO:HD3	0.41	1.77	2	1
1:A:14:VAL:CG1	1:A:29:ILE:HG12	0.41	2.46	6	1
1:A:56:LEU:CD2	1:A:125:LEU:HA	0.41	2.43	2	1
1:A:61:ILE:HD11	1:A:111:LEU:CD2	0.41	2.45	3	1
1:A:17:LYS:HE3	1:A:24:PHE:CE2	0.41	2.50	8	1
1:A:24:PHE:N	1:A:24:PHE:CD1	0.41	2.89	4	1
1:A:61:ILE:HG13	1:A:111:LEU:HD13	0.40	1.91	2	1
1:A:17:LYS:HG2	1:A:24:PHE:CD1	0.40	2.51	10	1
1:A:12:LEU:H	1:A:29:ILE:CG2	0.40	2.29	8	1
1:A:81:VAL:HG21	1:A:93:TYR:CE2	0.40	2.51	6	1
1:A:95:ILE:HD13	1:A:95:ILE:N	0.40	2.31	10	1
1:A:59:ASP:O	1:A:84:HIS:CE1	0.40	2.75	8	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	106/156 (68%)	89±1 (84±1%)	11±1 (10±1%)	6±1 (6±1%)	4	23
All	All	1060/1560 (68%)	892 (84%)	109 (10%)	59 (6%)	4	23

All 11 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	60	PRO	10
1	A	63	LEU	10
1	A	125	LEU	9
1	A	57	LEU	7
1	A	106	ARG	6
1	A	34	GLY	5
1	A	53	LYS	4
1	A	124	GLY	3
1	A	126	GLY	3
1	A	66	GLU	1
1	A	12	LEU	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	88/131 (67%)	64±3 (72±3%)	24±3 (28±3%)	2	21
All	All	880/1310 (67%)	636 (72%)	244 (28%)	2	21

All 53 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	57	LEU	10
1	A	76	PHE	10
1	A	33	VAL	10
1	A	53	LYS	10
1	A	55	ILE	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	40	LEU	10
1	A	67	VAL	10
1	A	24	PHE	9
1	A	70	LEU	9
1	A	58	CYS	9
1	A	17	LYS	9
1	A	12	LEU	7
1	A	111	LEU	7
1	A	110	ILE	7
1	A	113	LEU	7
1	A	118	ARG	7
1	A	37	LYS	7
1	A	48	CYS	6
1	A	61	ILE	6
1	A	23	TYR	6
1	A	125	LEU	5
1	A	51	LEU	5
1	A	109	VAL	5
1	A	56	LEU	5
1	A	59	ASP	5
1	A	101	ARG	4
1	A	84	HIS	4
1	A	105	LYS	3
1	A	21	ASN	3
1	A	13	ARG	3
1	A	93	TYR	3
1	A	115	GLN	3
1	A	29	ILE	2
1	A	92	TYR	2
1	A	81	VAL	2
1	A	68	THR	2
1	A	49	ASP	2
1	A	120	ARG	2
1	A	95	ILE	2
1	A	63	LEU	2
1	A	15	VAL	2
1	A	32	ASP	1
1	A	97	LYS	1
1	A	28	LYS	1
1	A	26	SER	1
1	A	100	GLN	1
1	A	98	GLU	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	119	LEU	1
1	A	123	TYR	1
1	A	39	LYS	1
1	A	18	TRP	1
1	A	106	ARG	1
1	A	60	PRO	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 86% for the well-defined parts and 86% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 5878

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1769
Number of shifts mapped to atoms	1769
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	149	-0.06 ± 0.12	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	133	0.21 ± 0.18	None needed (< 0.5 ppm)
$^{13}\text{C}'$	143	0.19 ± 0.09	None needed (< 0.5 ppm)
^{15}N	138	0.55 ± 0.27	Should be applied

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 86%, i.e. 1126 atoms were assigned a chemical shift out of a possible 1312. 4 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	512/524 (98%)	204/209 (98%)	208/212 (98%)	100/103 (97%)
Sidechain	496/666 (74%)	308/391 (79%)	183/243 (75%)	5/32 (16%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	118/122 (97%)	61/63 (97%)	55/56 (98%)	2/3 (67%)
Overall	1126/1312 (86%)	573/663 (86%)	446/511 (87%)	107/138 (78%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 86%, i.e. 1309 atoms were assigned a chemical shift out of a possible 1529. 4 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	583/609 (96%)	232/243 (95%)	239/246 (97%)	112/120 (93%)
Sidechain	583/773 (75%)	364/454 (80%)	213/282 (76%)	6/37 (16%)
Aromatic	143/147 (97%)	74/76 (97%)	67/68 (99%)	2/3 (67%)
Overall	1309/1529 (86%)	670/773 (87%)	519/596 (87%)	120/160 (75%)

7.1.4 Statistically unusual chemical shifts ⓘ


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	94	SER	HB2	1.50	5.18 – 2.58	-9.2
1	A	103	TRP	CZ2	124.06	121.76 – 106.66	6.5

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: BMRB entry 5878

Chemical shift list name: *assigned_chem_shift_list_2*

7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	92
Number of shifts mapped to atoms	92
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.2.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 6%, i.e. 74 atoms were assigned a chemical shift out of a possible 1312. 0 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	64/524 (12%)	27/209 (13%)	23/212 (11%)	14/103 (14%)
Sidechain	10/666 (2%)	4/391 (1%)	6/243 (2%)	0/32 (0%)
Aromatic	0/122 (0%)	0/63 (0%)	0/56 (0%)	0/3 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Overall	74/1312 (6%)	31/663 (5%)	29/511 (6%)	14/138 (10%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 5%, i.e. 74 atoms were assigned a chemical shift out of a possible 1529. 0 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	64/609 (11%)	27/243 (11%)	23/246 (9%)	14/120 (12%)
Sidechain	10/773 (1%)	4/454 (1%)	6/282 (2%)	0/37 (0%)
Aromatic	0/147 (0%)	0/76 (0%)	0/68 (0%)	0/3 (0%)
Overall	74/1529 (5%)	31/773 (4%)	29/596 (5%)	14/160 (9%)

7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

