



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:41 am GMT

PDB ID : 3SV0
Title : Crystal structure of casein kinase-1 like protein in plant
Authors : Park, H.H.; Do, K.H.
Deposited on : 2011-07-12
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	<div> <div>5%</div> <div>47%</div> <div>11%</div> <div>39%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2606 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Casein kinase I-like.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	294	Total	C	N	O	S	0	0	0
			2424	1560	423	433	8			


There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	473	ALA	-	EXPRESSION TAG	UNP Q8LR51
A	474	ALA	-	EXPRESSION TAG	UNP Q8LR51
A	475	ALA	-	EXPRESSION TAG	UNP Q8LR51
A	476	LEU	-	EXPRESSION TAG	UNP Q8LR51
A	477	GLU	-	EXPRESSION TAG	UNP Q8LR51
A	478	HIS	-	EXPRESSION TAG	UNP Q8LR51
A	479	HIS	-	EXPRESSION TAG	UNP Q8LR51
A	480	HIS	-	EXPRESSION TAG	UNP Q8LR51
A	481	HIS	-	EXPRESSION TAG	UNP Q8LR51
A	482	HIS	-	EXPRESSION TAG	UNP Q8LR51
A	483	HIS	-	EXPRESSION TAG	UNP Q8LR51

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	182	Total	O	0	0
			182	182		

- Molecule 1: Casein kinase I-like



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	108.83Å 69.60Å 55.85Å 90.00° 109.47° 90.00°	Depositor
Resolution (Å)	30.83 – 2.00 30.83 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (30.83-2.00) 99.5 (30.83-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.197 , 0.250 0.198 , 0.254	Depositor DCC
R_{free} test set	1331 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	30.4	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2606	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	4/2481 (0.2%)	1.14	21/3338 (0.6%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	247	GLU	CB-CG	17.49	1.85	1.52
1	A	247	GLU	CG-CD	-7.97	1.40	1.51
1	A	286	TYR	N-CA	6.29	1.58	1.46
1	A	213	TRP	CE3-CZ3	5.22	1.47	1.38

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	192	ARG	NE-CZ-NH1	-13.26	113.67	120.30
1	A	256	ARG	NE-CZ-NH1	-12.16	114.22	120.30
1	A	256	ARG	NE-CZ-NH2	11.57	126.08	120.30
1	A	285	ASP	C-N-CA	-11.00	94.21	121.70
1	A	127	ARG	NE-CZ-NH1	-9.36	115.62	120.30
1	A	192	ARG	NE-CZ-NH2	8.97	124.78	120.30
1	A	247	GLU	CB-CG-CD	-7.52	93.89	114.20
1	A	128	ASP	CB-CG-OD1	7.43	124.99	118.30
1	A	279	ARG	NE-CZ-NH1	-6.92	116.84	120.30
1	A	11	LEU	CA-CB-CG	6.86	131.09	115.30
1	A	247	GLU	CG-CD-OE1	-6.67	104.97	118.30
1	A	271	ARG	NE-CZ-NH1	-6.57	117.01	120.30
1	A	285	ASP	O-C-N	-6.32	112.59	122.70
1	A	127	ARG	NE-CZ-NH2	6.24	123.42	120.30
1	A	271	ARG	NE-CZ-NH2	5.96	123.28	120.30
1	A	115	ARG	CG-CD-NE	-5.89	99.42	111.80
1	A	256	ARG	CD-NE-CZ	5.89	131.85	123.60
1	A	128	ASP	CB-CG-OD2	-5.57	113.29	118.30
1	A	91	ASP	CB-CG-OD2	-5.07	113.74	118.30
1	A	242	ARG	NE-CZ-NH2	-5.04	117.78	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	178	ARG	NE-CZ-NH1	-5.03	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2424	0	2426	80	0
2	A	182	0	0	15	0
All	All	2606	0	2426	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:LEU:HD13	1:A:225:TYR:CE2	1.14	1.60
1:A:173:LEU:CD1	1:A:225:TYR:CE2	1.88	1.55
1:A:247:GLU:CG	1:A:247:GLU:CB	1.85	1.54
1:A:173:LEU:HD13	1:A:225:TYR:CZ	1.62	1.32
1:A:247:GLU:CB	1:A:247:GLU:OE1	1.95	1.14
1:A:45:LYS:HD2	1:A:45:LYS:H	1.10	1.12
1:A:247:GLU:HB3	1:A:247:GLU:OE1	1.45	1.10
1:A:2:GLU:HB3	1:A:3:PRO:HD2	1.42	1.02
1:A:2:GLU:HB3	1:A:3:PRO:CD	1.95	0.97
1:A:173:LEU:CD2	1:A:184:THR:O	2.18	0.92
1:A:247:GLU:H	1:A:247:GLU:HG2	1.33	0.91
1:A:247:GLU:CB	1:A:247:GLU:CD	2.39	0.90
1:A:192:ARG:HD2	2:A:584:HOH:O	1.71	0.89
1:A:45:LYS:N	1:A:45:LYS:HD2	1.91	0.86
1:A:41:ASN:HD22	1:A:43:LYS:H	1.20	0.85
1:A:60:GLN:HG2	2:A:629:HOH:O	1.76	0.85
1:A:169:GLU:O	2:A:586:HOH:O	1.95	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:ARG:CD	2:A:584:HOH:O	2.25	0.84
1:A:247:GLU:CG	1:A:247:GLU:H	1.90	0.84
1:A:173:LEU:HD13	1:A:225:TYR:HE2	1.01	0.84
1:A:173:LEU:HD12	1:A:225:TYR:CE2	2.12	0.82
1:A:173:LEU:CD1	1:A:225:TYR:HE2	1.57	0.82
1:A:51:TYR:O	1:A:55:ILE:HD12	1.81	0.79
1:A:132:ASP:OD1	2:A:627:HOH:O	1.99	0.78
1:A:41:ASN:ND2	1:A:43:LYS:H	1.82	0.77
1:A:111:GLN:HE22	1:A:145:VAL:H	1.29	0.77
1:A:111:GLN:NE2	1:A:145:VAL:H	1.84	0.74
1:A:173:LEU:HD23	1:A:184:THR:O	1.86	0.74
1:A:238:GLU:OE2	1:A:242:ARG:NH2	2.22	0.72
1:A:173:LEU:HB3	1:A:225:TYR:OH	1.91	0.70
1:A:173:LEU:CD1	1:A:225:TYR:CZ	2.50	0.69
1:A:247:GLU:CG	1:A:247:GLU:N	2.56	0.68
1:A:295:TYR:C	1:A:295:TYR:CD2	2.69	0.66
1:A:60:GLN:CG	2:A:629:HOH:O	2.37	0.66
1:A:173:LEU:CD2	1:A:185:HIS:HA	2.25	0.66
1:A:173:LEU:HD11	1:A:225:TYR:CE2	2.22	0.65
1:A:54:LYS:HG3	2:A:637:HOH:O	1.98	0.64
1:A:144:GLN:HG3	2:A:620:HOH:O	1.98	0.64
1:A:247:GLU:CG	1:A:247:GLU:CA	2.76	0.63
1:A:295:TYR:C	1:A:295:TYR:HD2	2.00	0.63
1:A:41:ASN:HD22	1:A:43:LYS:N	1.97	0.62
1:A:173:LEU:HD11	1:A:225:TYR:HE2	1.57	0.61
1:A:247:GLU:HG3	1:A:276:LEU:HD22	1.82	0.60
1:A:287:VAL:HG12	1:A:292:ILE:HG12	1.82	0.60
1:A:173:LEU:HD22	1:A:185:HIS:HA	1.83	0.59
1:A:127:ARG:HD3	1:A:152:LEU:O	2.02	0.59
1:A:41:ASN:ND2	1:A:43:LYS:HB2	2.20	0.56
1:A:207:LEU:HD13	1:A:244:TYR:CD2	2.42	0.55
1:A:247:GLU:N	1:A:247:GLU:HG2	2.12	0.53
1:A:173:LEU:CD1	1:A:225:TYR:CD2	2.82	0.52
1:A:30:GLN:H	1:A:30:GLN:HE21	1.57	0.51
1:A:54:LYS:HE3	2:A:507:HOH:O	2.11	0.51
1:A:56:TYR:CE1	1:A:82:MET:HE1	2.46	0.50
1:A:173:LEU:HD21	1:A:185:HIS:HA	1.93	0.50
1:A:51:TYR:CZ	1:A:55:ILE:HD11	2.47	0.49
1:A:130:LYS:HE2	1:A:133:ASN:ND2	2.28	0.49
1:A:253:HIS:ND1	1:A:256:ARG:NH1	2.61	0.49
1:A:27:THR:HG23	2:A:601:HOH:O	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:LEU:HD21	1:A:185:HIS:C	2.34	0.48
1:A:246:THR:HG23	2:A:605:HOH:O	2.12	0.48
1:A:60:GLN:CD	2:A:629:HOH:O	2.52	0.47
1:A:173:LEU:HD21	1:A:185:HIS:CA	2.45	0.47
1:A:192:ARG:HD3	2:A:584:HOH:O	2.02	0.47
1:A:111:GLN:HE22	1:A:145:VAL:N	2.07	0.47
1:A:44:THR:HA	1:A:45:LYS:NZ	2.29	0.47
1:A:54:LYS:CE	2:A:507:HOH:O	2.62	0.47
1:A:45:LYS:H	1:A:45:LYS:CD	1.96	0.47
1:A:40:GLU:HB3	1:A:49:LEU:HD12	1.97	0.47
1:A:173:LEU:HD13	1:A:225:TYR:OH	2.09	0.47
1:A:130:LYS:HE2	1:A:133:ASN:HD21	1.80	0.46
1:A:111:GLN:HE22	1:A:144:GLN:HA	1.81	0.46
1:A:2:GLU:CB	1:A:3:PRO:CD	2.80	0.46
1:A:167:TYR:CE2	1:A:169:GLU:HG3	2.52	0.44
1:A:17:SER:HB3	2:A:519:HOH:O	2.19	0.43
1:A:111:GLN:HE21	1:A:115:ARG:NH1	2.18	0.42
1:A:45:LYS:N	1:A:45:LYS:CD	2.65	0.42
1:A:178:ARG:HD2	1:A:179:TYR:CZ	2.55	0.42
1:A:113:ILE:HG21	1:A:270:LYS:HG3	2.03	0.41
1:A:173:LEU:CD2	1:A:185:HIS:CA	2.97	0.41
1:A:171:LYS:HE2	1:A:171:LYS:HB3	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	292/483 (60%)	278 (95%)	13 (4%)	1 (0%)	44 40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	15	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/412 (63%)	242 (93%)	19 (7%)	16	11

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LYS
1	A	13	ARG
1	A	14	LYS
1	A	30	GLN
1	A	45	LYS
1	A	60	GLN
1	A	82	MET
1	A	140	ARG
1	A	160	SER
1	A	172	ASN
1	A	174	THR
1	A	192	ARG
1	A	207	LEU
1	A	224	LYS
1	A	229	SER
1	A	247	GLU
1	A	263	LYS
1	A	292	ILE
1	A	295	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	41	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	111	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/483 (60%)	0.12	23 (7%) 14 14	19, 31, 63, 78	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	173	LEU	9.7
1	A	19	SER	5.0
1	A	20	PHE	4.7
1	A	45	LYS	4.0
1	A	18	GLY	3.7
1	A	13	ARG	3.6
1	A	295	TYR	3.3
1	A	46	HIS	3.1
1	A	44	THR	3.1
1	A	2	GLU	3.0
1	A	29	ILE	3.0
1	A	225	TYR	2.9
1	A	30	GLN	2.8
1	A	16	GLY	2.6
1	A	4	ARG	2.6
1	A	17	SER	2.5
1	A	199	LEU	2.4
1	A	24	TYR	2.4
1	A	202	VAL	2.2
1	A	174	THR	2.2
1	A	170	ASN	2.2
1	A	14	LYS	2.1
1	A	160	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.