



Full wwPDB X-ray Structure Validation Report i

Nov 2, 2017 – 01:38 AM EDT

PDB ID : 3SVQ

Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with 6-((3R,4R)-4-(2-((2,2-Difluoro-2-(2,3-difluorophenyl)ethyl)amino)ethoxy)pyrrolidin-3-yl)methyl)-4-methylpyridin-2-amine

Authors : Li, H.; Poulos, T.L.

Deposited on : unknown

Resolution : 2.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.7.2 (RC1), CSD as538be (2017)

Xtriage (Phenix) : 1.9-1692

EDS : rb-20030345

Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)

Refmac : 5.8.0135

CCP4 : 6.5.0

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	100719	5526 (2.20-2.16)
Clashscore	112137	6386 (2.20-2.16)
Ramachandran outliers	110173	6282 (2.20-2.16)
Sidechain outliers	110143	6282 (2.20-2.16)
RSRZ outliers	101464	5562 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	JK4	A	800	-	-	-	X
4	JK4	B	800	-	-	-	X

2 Entry composition (i)


There are 7 unique types of molecules in this entry. The entry contains 6979 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	407	3324	2128	567	608	21	0	2	0
1	B	411	3351	2144	574	611	22	0	2	0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: C₉H₁₅N₅O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 17 9 5 3	0	0
3	B	1	Total C N O 17 9 5 3	0	0

- Molecule 4 is 6-{|[(3R,4R)-4-(2-{[2-(2,3-difluorophenyl)-2,2-difluoroethyl]amino}ethoxy)pyrrolidin-3-yl]methyl}-4-methylpyridin-2-amine (three-letter code: JK4) (formula: C₂₁H₂₆F₄N₄O).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C F N O 30 21 4 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
4	B	1	30	21	4	4	1	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	4	2	2	0	0
5	B	1	4	2	2	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	A	1	1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	51	51	51	0	0
7	B	64	64	64	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.75 Å 111.13 Å 164.71 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.82 – 2.18 43.82 – 2.18	Depositor EDS
% Data completeness (in resolution range)	97.2 (43.82-2.18) 97.2 (43.82-2.18)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$< I/\sigma(I) >$ ¹	1.23 (at 2.18 Å)	Xtriage
Refinement program	REFMAC	Depositor
R , R_{free}	0.209 , 0.262 0.214 , 0.267	Depositor DCC
R_{free} test set	2460 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	49.7	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.8	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6979	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: JK4, HEM, ZN, H4B, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.73	0/3423	0.74	0/4644
1	B	0.81	0/3450	0.77	0/4677
All	All	0.77	0/6873	0.75	0/9321

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3324	0	3233	56	0
1	B	3351	0	3269	42	0
2	A	43	0	30	3	0
2	B	43	0	30	8	0
3	A	17	0	15	0	0
3	B	17	0	15	1	0
4	A	30	0	26	5	0
4	B	30	0	26	5	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	51	0	0	1	0
7	B	64	0	0	2	0
All	All	6979	0	6650	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:750:HEM:HBA2	4:B:800:JK4:F15	1.53	0.99
1:B:592:GLU:OE1	4:B:800:JK4:H26	1.71	0.90
1:B:587:TRP:H	2:B:750:HEM:HAB	1.37	0.89
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.60	0.83
2:A:750:HEM:HMC2	2:A:750:HEM:HBC2	1.66	0.77
1:A:592:GLU:OE1	4:A:800:JK4:H26	1.88	0.73
1:A:317:HIS:O	1:A:320:SER:HB3	1.89	0.73
1:A:371:ARG:CG	1:A:371:ARG:HH21	2.06	0.67
1:A:380:ARG:HD3	1:A:400:GLU:OE1	1.95	0.67
2:A:750:HEM:CMC	2:A:750:HEM:HBC2	2.25	0.66
1:A:337:LEU:HD21	4:A:800:JK4:H03	1.77	0.66
1:A:371:ARG:HH21	1:A:371:ARG:HG2	1.64	0.62
1:B:481:ARG:HD3	1:B:498:ASN:ND2	2.14	0.62
1:B:494:GLY:O	1:B:496:PRO:HD3	2.00	0.60
2:A:750:HEM:HBA1	4:A:800:JK4:F15	1.90	0.60
1:A:371:ARG:HG2	1:A:371:ARG:NH2	2.15	0.59
1:A:391:THR:O	1:A:392:SER:OG	2.19	0.59
1:A:592:GLU:CD	4:A:800:JK4:H26	2.25	0.57
1:B:592:GLU:CD	4:B:800:JK4:H26	2.26	0.56
1:B:595:VAL:O	1:B:599:CYS:HB2	2.05	0.56
1:A:481:ARG:HD3	1:A:498:ASN:HD21	1.71	0.56
1:B:481:ARG:HD3	1:B:498:ASN:HD21	1.71	0.55
1:A:307[B]:GLU:OE2	1:B:602[B]:SER:OG	2.20	0.55
1:A:569:ASN:C	1:A:569:ASN:HD22	2.10	0.55
1:A:632:GLU:OE2	1:B:628:GLN:NE2	2.40	0.54
1:B:566:ALA:HB2	1:B:585:SER:HB3	1.89	0.53
1:A:473:ARG:NH2	1:A:710:PRO:HD3	2.23	0.53
1:B:566:ALA:CB	1:B:585:SER:HB3	2.38	0.53
1:A:439:PHE:HZ	1:A:537:PRO:HD3	1.72	0.53
1:B:349:ARG:HG2	1:B:353:GLN:HG3	1.91	0.52
2:B:750:HEM:O1A	3:B:760:H4B:N3	2.35	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:504:ILE:HG23	1:A:507:GLN:NE2	2.25	0.51
1:A:391:THR:O	1:A:392:SER:CB	2.59	0.51
1:A:480:ILE:HD13	1:A:541:VAL:HG13	1.93	0.50
1:A:336:MET:CE	1:A:678:TRP:HZ2	2.24	0.50
1:B:350:THR:O	1:B:353:GLN:HG2	2.12	0.50
1:B:478:GLN:HA	1:B:566:ALA:O	2.11	0.49
1:A:592:GLU:OE2	4:A:800:JK4:H26	2.13	0.49
1:B:608:GLU:HG3	7:B:24:HOH:O	2.13	0.48
1:A:710:PRO:O	1:A:714:HIS:N	2.45	0.48
1:B:480:ILE:HA	7:B:28:HOH:O	2.13	0.48
1:A:617:ASP:OD2	1:A:619:ARG:HB2	2.13	0.48
1:B:631:VAL:HG22	1:B:683:MET:HE1	1.95	0.48
1:A:455:LEU:HD12	1:A:587:TRP:CB	2.45	0.46
1:A:525:GLN:HG3	1:A:529:ASN:O	2.15	0.46
1:A:504:ILE:HG23	1:A:507:GLN:HE22	1.81	0.46
1:A:414:ARG:HD3	1:A:678:TRP:CD2	2.50	0.46
1:B:336:MET:HG2	4:B:800:JK4:H07B	1.96	0.46
1:A:664:ASN:O	1:A:668:CYS:HB2	2.16	0.46
1:B:487:GLN:HE22	1:B:514:ARG:NH2	2.12	0.46
1:A:607:LEU:HD11	1:B:687:ILE:HG21	1.97	0.46
1:B:587:TRP:O	2:B:750:HEM:HMB3	2.16	0.46
1:B:587:TRP:N	2:B:750:HEM:HAB	2.18	0.46
1:B:438:MET:O	1:B:442:ILE:HG13	2.16	0.45
1:A:330:ILE:HD11	1:B:696:LEU:HB3	1.98	0.45
1:A:588:TYR:CD1	1:A:593:ILE:HD11	2.50	0.45
2:B:750:HEM:HBC2	2:B:750:HEM:CMC	2.46	0.45
1:A:676:TRP:CE2	1:A:680:VAL:HG21	2.52	0.45
1:A:481:ARG:HD3	1:A:498:ASN:ND2	2.31	0.44
1:A:475:TRP:HB2	1:A:523:LEU:HB3	2.00	0.44
1:B:409:TRP:CZ3	2:B:750:HEM:HMC3	2.52	0.44
1:B:459:ILE:HD11	1:B:582:CYS:HB2	1.99	0.44
1:A:685:GLY:O	1:A:691:PHE:HB2	2.18	0.44
1:A:336:MET:HB3	1:A:336:MET:HE2	1.90	0.44
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.53	0.44
1:A:516:ARG:NH1	1:A:604:TYR:HD2	2.16	0.43
1:B:413:SER:OG	1:B:697:ASN:ND2	2.51	0.43
1:A:686:SER:HA	1:A:691:PHE:CG	2.52	0.43
1:B:317:HIS:O	1:B:320:SER:HB2	2.18	0.43
1:A:330:ILE:HG23	1:A:330:ILE:O	2.17	0.43
1:A:477:SER:HA	1:A:569:ASN:HB3	2.01	0.43
1:B:481:ARG:CD	1:B:498:ASN:HD21	2.32	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:431:ASP:OD2	1:A:431:ASP:C	2.57	0.42
1:A:550:LYS:HB2	1:A:550:LYS:HE3	1.75	0.42
1:B:537:PRO:HA	1:B:538:PRO:HD3	1.76	0.42
1:A:494:GLY:HA2	1:A:516:ARG:HG3	2.01	0.42
1:A:536:ILE:O	1:A:537:PRO:C	2.57	0.42
1:A:628:GLN:HG2	1:B:631:VAL:HG11	2.02	0.42
1:A:516:ARG:N	7:A:79:HOH:O	2.52	0.42
1:B:597:ASP:OD1	1:B:603:ARG:NH1	2.52	0.42
1:B:610:VAL:O	1:B:614:MET:HG3	2.19	0.42
1:B:512:ALA:HA	1:B:513:PRO:HD3	1.96	0.42
1:A:299:ARG:O	1:A:299:ARG:HG3	2.19	0.41
1:B:337:LEU:HA	1:B:338:PRO:HD2	1.94	0.41
1:B:403:TYR:C	1:B:403:TYR:CD2	2.92	0.41
1:B:478:GLN:NE2	4:B:800:JK4:H11A	2.36	0.41
1:A:696:LEU:HB3	1:B:330:ILE:HD11	2.02	0.41
1:B:584:PHE:CD1	2:B:750:HEM:CAC	3.04	0.41
1:A:475:TRP:CZ3	1:A:711:TRP:HB3	2.55	0.41
1:A:676:TRP:CZ2	1:A:680:VAL:HG21	2.56	0.41
1:A:609:GLU:O	1:A:613:LYS:HG2	2.21	0.41
1:A:530:ASP:HA	1:A:531:PRO:HD3	1.90	0.40
1:A:674:ALA:HB3	1:A:695:MET:HB3	2.03	0.40
1:B:572:LEU:HB3	1:B:579:PHE:HB2	2.03	0.40
1:A:403:TYR:CE1	1:A:407:HIS:CE1	3.10	0.40
1:B:709:ASP:HA	1:B:710:PRO:HD3	1.87	0.40
1:A:455:LEU:HD12	1:A:587:TRP:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	405/422 (96%)	388 (96%)	16 (4%)	1 (0%)	51 56

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	409/422 (97%)	395 (97%)	13 (3%)	1 (0%)	51 56
All	All	814/844 (96%)	783 (96%)	29 (4%)	2 (0%)	51 56

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	685	GLY
1	A	392	SER

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	365/377 (97%)	338 (93%)	27 (7%)	16 15
1	B	368/377 (98%)	354 (96%)	14 (4%)	38 45
All	All	733/754 (97%)	692 (94%)	41 (6%)	26 26

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	300	PHE
1	A	320	SER
1	A	321	THR
1	A	328	GLU
1	A	336	MET
1	A	371	ARG
1	A	380	ARG
1	A	381	LEU
1	A	382	GLU
1	A	390	SER
1	A	481	ARG
1	A	485	TYR
1	A	486	LYS
1	A	516	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	523	LEU
1	A	535	GLN
1	A	547	ARG
1	A	550	LYS
1	A	555	LYS
1	A	569	ASN
1	A	601[A]	ASN
1	A	601[B]	ASN
1	A	613	LYS
1	A	620	LYS
1	A	645	LYS
1	A	667	ARG
1	A	715	VAL
1	B	321	THR
1	B	350	THR
1	B	353	GLN
1	B	382	GLU
1	B	390	SER
1	B	454	ASN
1	B	481	ARG
1	B	486	LYS
1	B	507	GLN
1	B	527	ASN
1	B	547	ARG
1	B	627	ASP
1	B	668[A]	CYS
1	B	668[B]	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	454	ASN
1	A	507	GLN
1	A	527	ASN
1	A	569	ASN
1	A	605	ASN
1	A	642	GLN
1	A	697	ASN
1	B	454	ASN
1	B	601	ASN
1	B	605	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	628	GLN
1	B	642	GLN
1	B	697	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	750	1	28,50,50	2.33	15 (53%)	17,82,82	2.20	3 (17%)
3	H4B	A	760	-	14,18,18	0.90	0	12,26,26	2.54	6 (50%)
4	JK4	A	800	-	28,32,32	1.03	2 (7%)	34,45,45	2.13	9 (26%)
5	ACT	A	860	-	1,3,3	0.85	0	0,3,3	0.00	-
2	HEM	B	750	1	28,50,50	2.40	14 (50%)	17,82,82	2.02	5 (29%)
3	H4B	B	760	-	14,18,18	0.91	0	12,26,26	2.75	6 (50%)
4	JK4	B	800	-	28,32,32	0.91	1 (3%)	34,45,45	2.49	11 (32%)
5	ACT	B	860	-	1,3,3	1.00	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/6/54/54	0/0/8/8
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
4	JK4	A	800	-	-	0/17/29/29	0/3/3/3
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/6/54/54	0/0/8/8
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
4	JK4	B	800	-	-	0/17/29/29	0/3/3/3
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	C3B-C2B	-5.01	1.33	1.40
2	A	750	HEM	C3B-C2B	-4.85	1.34	1.40
2	B	750	HEM	C3C-C2C	-4.57	1.34	1.40
2	A	750	HEM	C3C-C2C	-3.76	1.35	1.40
4	A	800	JK4	F15-C14	-2.85	1.31	1.37
4	A	800	JK4	F16-C14	-2.43	1.32	1.37
4	B	800	JK4	F15-C14	-2.15	1.32	1.37
2	A	750	HEM	CMA-C3A	2.00	1.55	1.51
2	B	750	HEM	C4A-NA	2.02	1.40	1.36
2	B	750	HEM	C1A-NA	2.03	1.40	1.36
2	A	750	HEM	C4B-NB	2.03	1.40	1.36
2	A	750	HEM	CAD-C3D	2.07	1.56	1.52
2	A	750	HEM	C1D-ND	2.10	1.40	1.36
2	B	750	HEM	CMB-C2B	2.14	1.56	1.51
2	B	750	HEM	C4D-ND	2.14	1.39	1.36
2	B	750	HEM	CAA-C2A	2.17	1.55	1.52
2	B	750	HEM	CAD-C3D	2.18	1.56	1.52
2	B	750	HEM	CMD-C2D	2.21	1.56	1.51
2	B	750	HEM	C3B-CAB	2.28	1.52	1.47
2	A	750	HEM	CMD-C2D	2.29	1.56	1.51
2	B	750	HEM	CMA-C3A	2.35	1.56	1.51
2	A	750	HEM	C3C-CAC	2.41	1.52	1.47
2	A	750	HEM	C1C-NC	2.43	1.39	1.36
2	A	750	HEM	C1B-NB	2.67	1.39	1.36
2	A	750	HEM	C4D-ND	2.69	1.39	1.36
2	A	750	HEM	CMB-C2B	2.77	1.57	1.51
2	A	750	HEM	CMC-C2C	3.02	1.58	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	C3C-CAC	3.20	1.54	1.47
2	A	750	HEM	C3B-CAB	3.32	1.54	1.47
2	B	750	HEM	C1B-NB	4.54	1.42	1.36
2	B	750	HEM	C3D-C2D	4.86	1.52	1.37
2	A	750	HEM	C3D-C2D	5.01	1.52	1.37

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	HEM	CBA-CAA-C2A	-6.45	100.14	112.48
3	A	760	H4B	N3-C2-N1	-4.92	117.47	125.45
4	B	800	JK4	F23-C23-C22	-4.64	112.24	118.30
2	B	750	HEM	CBA-CAA-C2A	-4.49	103.90	112.48
2	B	750	HEM	CBD-CAD-C3D	-4.24	104.38	112.47
4	A	800	JK4	F23-C23-C22	-3.60	113.60	118.30
4	B	800	JK4	C04-C05-C06	-3.42	118.11	120.26
4	B	800	JK4	F22-C22-C23	-3.34	112.15	119.01
4	A	800	JK4	C05-C06-N01	-3.15	119.49	122.91
3	B	760	H4B	N3-C2-N1	-3.09	120.44	125.45
4	B	800	JK4	C05-C06-N01	-2.61	120.08	122.91
2	B	750	HEM	C1D-C2D-C3D	-2.45	105.29	107.00
4	B	800	JK4	C26-C25-C24	-2.20	117.14	120.24
4	A	800	JK4	F22-C22-C23	-2.16	114.57	119.01
3	B	760	H4B	C4A-C4-N3	-2.12	117.67	123.91
3	B	760	H4B	C6-C7-N8	-2.10	107.67	111.01
2	B	750	HEM	C3C-C4C-NC	-2.04	107.08	110.94
4	A	800	JK4	F16-C14-C21	2.09	113.34	110.39
3	A	760	H4B	N2-C2-N1	2.14	120.66	117.24
3	A	760	H4B	C4-C4A-C8A	2.34	116.69	114.56
2	A	750	HEM	CMC-C2C-C3C	2.46	129.45	124.89
3	B	760	H4B	C2-N1-C8A	2.69	120.56	114.51
4	B	800	JK4	C5'-N1'-C2'	2.70	111.67	105.40
4	B	800	JK4	C10-C11-N12	2.86	118.53	111.45
3	A	760	H4B	N2-C2-N3	2.90	121.87	117.24
2	B	750	HEM	C4C-C3C-C2C	2.92	108.94	106.90
4	B	800	JK4	C11-N12-C13	2.92	119.26	113.35
4	A	800	JK4	C11-N12-C13	3.35	120.12	113.35
3	A	760	H4B	C4-N3-C2	3.44	121.01	116.06
4	A	800	JK4	F22-C22-C21	3.51	123.19	118.52
4	B	800	JK4	F23-C23-C24	3.56	126.44	118.49
4	A	800	JK4	F23-C23-C24	3.64	126.62	118.49
4	A	800	JK4	C5'-N1'-C2'	3.65	113.89	105.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	760	H4B	C4-N3-C2	4.09	121.94	116.06
2	A	750	HEM	C4C-C3C-C2C	4.25	109.87	106.90
3	A	760	H4B	C2-N1-C8A	4.36	124.33	114.51
4	B	800	JK4	F22-C22-C21	5.58	125.95	118.52
3	B	760	H4B	C4-C4A-C8A	6.29	120.26	114.56
4	A	800	JK4	C02-N01-C06	6.79	122.97	118.17
4	B	800	JK4	C02-N01-C06	7.77	123.66	118.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	3	0
4	A	800	JK4	5	0
2	B	750	HEM	8	0
3	B	760	H4B	1	0
4	B	800	JK4	5	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/422 (96%)	0.82	55 (13%) 3 3	36, 65, 105, 134	0
1	B	411/422 (97%)	0.60	37 (9%) 10 10	33, 55, 83, 111	0
All	All	818/844 (96%)	0.71	92 (11%) 6 6	33, 59, 100, 134	0

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	488	PRO	9.3
1	B	300	PHE	6.3
1	B	351	LYS	5.8
1	A	715	VAL	5.8
1	A	351	LYS	5.7
1	B	350	THR	5.6
1	B	348	VAL	5.5
1	A	716	TRP	5.4
1	A	373	GLY	5.0
1	B	352	ASP	5.0
1	A	350	THR	4.4
1	B	619	ARG	4.2
1	B	718	GLY	4.1
1	A	300	PHE	4.1
1	A	713	THR	3.9
1	A	318	LEU	3.8
1	A	486	LYS	3.6
1	B	353	GLN	3.6
1	A	554	PHE	3.5
1	A	370	LYS	3.4
1	A	493	LEU	3.3
1	A	299	ARG	3.3
1	A	485	TYR	3.3
1	A	487	GLN	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	385	ASN	3.2
1	A	355	PHE	3.1
1	B	677	VAL	3.1
1	B	680	VAL	3.1
1	A	494	GLY	3.1
1	A	352	ASP	3.0
1	A	469	LYS	2.9
1	A	389	GLU	2.9
1	B	479	LEU	2.9
1	A	714	HIS	2.9
1	B	620	LYS	2.9
1	A	381	LEU	2.9
1	B	591	THR	2.8
1	A	676	TRP	2.8
1	B	717	LYS	2.8
1	A	507	GLN	2.8
1	B	593	ILE	2.7
1	A	706	TYR	2.7
1	B	392	SER	2.7
1	A	503	GLU	2.7
1	A	553	TRP	2.6
1	A	415	CYS	2.6
1	B	349	ARG	2.6
1	A	678	TRP	2.6
1	A	536	ILE	2.6
1	B	442	ILE	2.5
1	B	480	ILE	2.5
1	B	443	CYS	2.5
1	B	567	VAL	2.5
1	A	388	ILE	2.5
1	A	479	LEU	2.5
1	A	712	ASN	2.5
1	B	564	LEU	2.4
1	B	446	VAL	2.4
1	A	504	ILE	2.4
1	A	677	VAL	2.3
1	B	541	VAL	2.3
1	A	480	ILE	2.3
1	B	552	ASP	2.3
1	A	680	VAL	2.3
1	B	375	LYS	2.3
1	A	492	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	394	TYR	2.3
1	A	533	LEU	2.3
1	A	514	ARG	2.3
1	B	584	PHE	2.2
1	A	681	PRO	2.2
1	B	681	PRO	2.2
1	B	676	TRP	2.2
1	B	678	TRP	2.2
1	B	667	ARG	2.2
1	A	361	PHE	2.1
1	B	369	ILE	2.1
1	A	567	VAL	2.1
1	A	506	ILE	2.1
1	A	490	GLY	2.1
1	A	378	MET	2.1
1	B	355	PHE	2.1
1	B	583	PRO	2.1
1	A	390	SER	2.1
1	B	696	LEU	2.1
1	A	393	THR	2.1
1	A	691	PHE	2.1
1	A	392	SER	2.0
1	B	492	THR	2.0
1	A	369	ILE	2.0
1	B	616	LEU	2.0
1	A	371	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	JK4	A	800	30/30	0.75	0.31	3.62	73,90,96,96	0
4	JK4	B	800	30/30	0.72	0.28	2.98	61,68,73,78	0
2	HEM	A	750	43/43	0.97	0.21	1.68	37,45,59,67	0
2	HEM	B	750	43/43	0.98	0.20	0.92	38,45,55,59	0
3	H4B	B	760	17/17	0.93	0.21	0.44	39,44,51,54	0
3	H4B	A	760	17/17	0.91	0.18	0.18	45,52,58,58	0
5	ACT	B	860	4/4	0.95	0.12	-0.29	67,68,68,69	0
6	ZN	A	900	1/1	0.99	0.12	-0.74	56,56,56,56	0
5	ACT	A	860	4/4	0.96	0.12	-1.04	67,67,68,68	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.