



Full wwPDB X-ray Structure Validation Report i

Feb 28, 2018 – 12:24 PM EST

PDB ID : 3SXB
Title : Crystal structure of ABBB+UDP+Gal with MPD as the cryoprotectant
Authors : Johal, A.R.; Evans, S.V.
Deposited on : 2011-07-14
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2279 (1.50-1.50)
Clashscore	112137	2503 (1.50-1.50)
Ramachandran outliers	110173	2445 (1.50-1.50)
Sidechain outliers	110143	2443 (1.50-1.50)
RSRZ outliers	101464	2305 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GAL	A	2	-	-	-	X
4	PEG	A	3[A]	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	3[B]	-	-	-	X

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 2575 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histo-blood group ABO system transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	2256	1463	388	391	14	0	1	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	62	PHE	-	EXPRESSION TAG	UNP P16442
A	63	MET	-	EXPRESSION TAG	UNP P16442
A	235	SER	GLY	SEE REMARK 999	UNP P16442
A	266	MET	LEU	SEE REMARK 999	UNP P16442
A	268	ALA	GLY	SEE REMARK 999	UNP P16442

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C₉H₁₄N₂O₁₂P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	25	9	2	12	2	0	0

- Molecule 3 is BETA-D-GALACTOSE (three-letter code: GAL) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 12 6 6	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 14 8 6	0	1

- Molecule 5 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mn 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	267	Total O 267 267	0	0

3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histo-blood group ABO system transferase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	52.59 Å 149.84 Å 79.44 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.49 19.83 – 1.49	Depositor EDS
% Data completeness (in resolution range)	95.5 (20.00-1.49) 95.5 (19.83-1.49)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.60 (at 1.49 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R , R_{free}	0.194 , 0.221 0.186 , 0.214	Depositor DCC
R_{free} test set	2496 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	20.6	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 52.2	EDS
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2575	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, PEG, GAL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.53	17/2321 (0.7%)	1.37	12/3147 (0.4%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	210	VAL	CB-CG1	6.79	1.67	1.52
1	A	244	PHE	CE2-CZ	6.64	1.50	1.37
1	A	169	GLN	CB-CG	-6.50	1.35	1.52
1	A	113	ASN	CB-CG	6.49	1.66	1.51
1	A	338	LYS	CE-NZ	6.48	1.65	1.49
1	A	229	PHE	CE2-CZ	6.03	1.48	1.37
1	A	206	TYR	CE2-CZ	5.98	1.46	1.38
1	A	186	MET	C-O	5.95	1.34	1.23
1	A	244	PHE	CG-CD1	5.63	1.47	1.38
1	A	64	VAL	CB-CG1	5.63	1.64	1.52
1	A	215	GLU	CD-OE2	-5.61	1.19	1.25
1	A	241	ARG	CZ-NH2	5.46	1.40	1.33
1	A	323	TYR	CE1-CZ	5.41	1.45	1.38
1	A	110	ARG	CZ-NH2	5.19	1.39	1.33
1	A	271	GLY	N-CA	5.18	1.53	1.46
1	A	282	ARG	CG-CD	5.12	1.64	1.51
1	A	209	CYS	CB-SG	5.08	1.90	1.82

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	213	ASP	CB-CG-OD1	13.45	130.40	118.30
1	A	259	ASP	CB-CG-OD1	9.20	126.58	118.30
1	A	213	ASP	CB-CG-OD2	-7.05	111.96	118.30
1	A	340	ARG	NE-CZ-NH2	-6.92	116.84	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	312	ARG	NE-CZ-NH2	-6.83	116.89	120.30
1	A	282	ARG	NE-CZ-NH1	6.79	123.70	120.30
1	A	269	PHE	CB-CG-CD2	6.06	125.04	120.80
1	A	329	LEU	CB-CG-CD1	5.83	120.91	111.00
1	A	329	LEU	CB-CG-CD2	-5.65	101.39	111.00
1	A	341	PHE	CB-CG-CD1	-5.29	117.10	120.80
1	A	340	ARG	NE-CZ-NH1	5.08	122.84	120.30
1	A	103	ASP	CB-CG-OD2	5.00	122.80	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2256	0	2251	52	2
2	A	25	0	11	0	0
3	A	12	0	12	1	0
4	A	14	0	17	4	0
5	A	1	0	0	0	0
6	A	267	0	0	27	3
All	All	2575	0	2291	55	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:SER:HB3	1:A:242:GLU:OE1	1.49	1.12
1:A:186:MET:SD	1:A:297:GLU:OE2	2.08	1.11
1:A:187:ARG:HB3	1:A:191:MET:HE2	1.34	1.10
4:A:3[A]:PEG:H12	6:A:28:HOH:O	1.60	1.01
1:A:196:CYS:N	6:A:534:HOH:O	2.06	0.89
1:A:186:MET:HG2	6:A:446:HOH:O	1.75	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:SER:CB	1:A:242:GLU:OE1	2.25	0.84
1:A:233:HIS:HD2	1:A:235:SER:H	1.26	0.83
1:A:187:ARG:CB	1:A:191:MET:HE2	2.11	0.80
1:A:312:ARG:CZ	6:A:414:HOH:O	2.30	0.79
1:A:186:MET:CG	6:A:446:HOH:O	2.27	0.79
1:A:197:GLU:HA	1:A:278:GLN:NE2	2.01	0.75
1:A:63:MET:SD	1:A:112:GLN:NE2	2.58	0.73
1:A:196:CYS:HA	6:A:534:HOH:O	1.88	0.73
1:A:187:ARG:HB3	1:A:191:MET:CE	2.17	0.71
1:A:101:ASN:HD22	1:A:101:ASN:C	1.94	0.71
1:A:196:CYS:CA	6:A:534:HOH:O	2.36	0.70
1:A:123:ILE:HG22	1:A:124:LYS:HG3	1.75	0.69
1:A:187:ARG:CB	1:A:191:MET:CE	2.73	0.66
1:A:75:LYS:HB2	1:A:78:THR:HG23	1.79	0.65
1:A:312:ARG:NH2	6:A:414:HOH:O	2.28	0.64
1:A:101:ASN:HD21	1:A:103:ASP:HB2	1.63	0.63
4:A:3[A]:PEG:H31	6:A:25:HOH:O	2.00	0.62
1:A:186:MET:HG3	6:A:446:HOH:O	1.98	0.59
1:A:78:THR:HG22	6:A:530:HOH:O	2.03	0.58
1:A:62:PHE:N	6:A:527:HOH:O	2.39	0.56
1:A:312:ARG:NH1	6:A:414:HOH:O	2.34	0.56
1:A:257:PRO:HG2	1:A:260:GLU:OE2	2.05	0.56
1:A:176:ARG:HD2	6:A:540:HOH:O	2.06	0.55
1:A:63:MET:CE	6:A:541:HOH:O	2.56	0.54
1:A:197:GLU:HA	1:A:278:GLN:HE22	1.74	0.53
1:A:305:HIS:HE1	6:A:36:HOH:O	1.91	0.53
1:A:110:ARG:HH22	1:A:113:ASN:HD22	1.55	0.53
1:A:285:HIS:HD2	6:A:419:HOH:O	1.90	0.52
1:A:106:ASN:OD1	1:A:145:HIS:HE1	1.93	0.52
1:A:242:GLU:H	1:A:242:GLU:CD	2.13	0.51
1:A:305:HIS:HD2	6:A:46:HOH:O	1.93	0.51
1:A:233:HIS:HE1	3:A:2:GAL:O4	1.94	0.51
1:A:217:ARG:NH2	6:A:511:HOH:O	2.08	0.50
4:A:3[B]:PEG:O1	6:A:34:HOH:O	2.20	0.49
1:A:187:ARG:HB2	1:A:191:MET:CE	2.43	0.49
1:A:193:SER:HB3	1:A:281:THR:OG1	2.14	0.48
1:A:80:CYS:HB3	1:A:98:GLY:O	2.14	0.47
1:A:102:ILE:HG21	4:A:3[B]:PEG:H11	1.98	0.46
1:A:233:HIS:CD2	1:A:235:SER:H	2.17	0.46
1:A:161:ARG:NH1	6:A:463:HOH:O	2.48	0.46
1:A:145:HIS:HD2	6:A:9:HOH:O	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:ARG:HA	1:A:187:ARG:HD2	1.76	0.45
1:A:186:MET:N	6:A:376:HOH:O	2.49	0.44
1:A:176:ARG:HB3	6:A:540:HOH:O	2.17	0.43
1:A:62:PHE:N	6:A:502:HOH:O	2.51	0.43
1:A:110:ARG:HH22	1:A:113:ASN:ND2	2.16	0.43
1:A:63:MET:HE2	6:A:541:HOH:O	2.15	0.42
1:A:266:MET:HB3	1:A:324:LEU:CD2	2.49	0.42
1:A:78:THR:CG2	6:A:530:HOH:O	2.65	0.42

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:282:ARG:NH1	6:A:544:HOH:O[8_555]	1.46	0.74
1:A:312:ARG:NE	6:A:546:HOH:O[3_655]	1.92	0.28
6:A:374:HOH:O	6:A:538:HOH:O[3_655]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	270/293 (92%)	263 (97%)	6 (2%)	1 (0%)	38 14

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	187	ARG

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	246/262 (94%)	239 (97%)	7 (3%)	49 16

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	VAL
1	A	78	THR
1	A	101	ASN
1	A	113	ASN
1	A	186	MET
1	A	269	PHE
1	A	329	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	GLN
1	A	101	ASN
1	A	113	ASN
1	A	145	HIS
1	A	233	HIS
1	A	275	GLN
1	A	278	GLN
1	A	285	HIS
1	A	286	GLN
1	A	294	ASN
1	A	305	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UDP	A	1	5	21,26,26	2.17	6 (28%)	22,40,40	2.17	2 (9%)
3	GAL	A	2	-	12,12,12	0.92	1 (8%)	17,17,17	1.92	8 (47%)
4	PEG	A	3[A]	-	6,6,6	1.13	1 (16%)	5,5,5	3.56	4 (80%)
4	PEG	A	3[B]	-	6,6,6	1.17	0	5,5,5	1.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	A	1	5	-	0/12/32/32	0/2/2/2
3	GAL	A	2	-	-	0/2/22/22	0/1/1/1
4	PEG	A	3[A]	-	-	0/4/4/4	0/0/0/0
4	PEG	A	3[B]	-	-	0/4/4/4	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	UDP	PB-O3A	-2.02	1.56	1.60
4	A	3[A]	PEG	C3-C4	2.13	1.60	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2	GAL	C6-C5	2.14	1.59	1.51
2	A	1	UDP	O2'-C2'	2.22	1.48	1.43
2	A	1	UDP	C4-N3	3.08	1.38	1.33
2	A	1	UDP	PB-O1B	3.20	1.61	1.50
2	A	1	UDP	C6-N1	5.20	1.42	1.35
2	A	1	UDP	O4'-C1'	5.37	1.48	1.41

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	UDP	C4'-O4'-C1'	-5.32	104.11	109.77
3	A	2	GAL	C1-O5-C5	-3.06	107.88	113.39
3	A	2	GAL	O1-C1-O5	-2.60	102.50	110.20
3	A	2	GAL	C4-C3-C2	-2.22	106.92	110.84
3	A	2	GAL	O2-C2-C1	-2.11	105.36	109.75
3	A	2	GAL	O1-C1-C2	2.02	115.11	109.42
4	A	3[A]	PEG	C3-O2-C2	2.43	123.84	113.30
3	A	2	GAL	O3-C3-C2	2.55	115.91	110.36
4	A	3[A]	PEG	O1-C1-C2	2.69	127.31	111.89
3	A	2	GAL	O5-C5-C4	2.81	114.84	109.66
3	A	2	GAL	O5-C1-C2	2.98	114.98	110.04
4	A	3[A]	PEG	O4-C4-C3	4.68	138.79	111.89
4	A	3[A]	PEG	O2-C3-C4	5.06	133.49	110.15
2	A	1	UDP	C4-N3-C2	7.30	120.40	114.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2	GAL	1	0
4	A	3[A]	PEG	2	0
4	A	3[B]	PEG	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/293 (93%)	0.62	28 (10%) 7 8	9, 17, 35, 52	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	80	CYS	8.2
1	A	177	ALA	7.1
1	A	187	ARG	4.4
1	A	62	PHE	4.1
1	A	186	MET	4.1
1	A	176	ARG	4.0
1	A	345	PRO	3.9
1	A	346	LYS	3.6
1	A	64	VAL	3.4
1	A	191	MET	3.2
1	A	198	ARG	3.2
1	A	201	LEU	3.1
1	A	202	SER	3.1
1	A	344	VAL	3.1
1	A	294	ASN	3.0
1	A	197	GLU	2.9
1	A	161	ARG	2.9
1	A	293	ALA	2.8
1	A	199	ARG	2.8
1	A	78	THR	2.7
1	A	192	ILE	2.6
1	A	63	MET	2.5
1	A	256	ILE	2.4
1	A	292	GLN	2.3
1	A	328	GLN	2.1
1	A	251	GLN	2.1
1	A	196	CYS	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	151	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GAL	A	2	12/12	0.90	0.18	5.27	21,26,34,38	0
4	PEG	A	3[A]	7/7	0.78	0.21	4.41	13,22,26,34	7
4	PEG	A	3[B]	7/7	0.78	0.21	4.25	15,24,28,33	7
2	UDP	A	1	25/25	0.85	0.14	0.11	13,21,26,30	18
5	MN	A	355	1/1	0.98	0.09	-	18,18,18,18	1

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.