



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 16, 2017 – 01:21 PM EDT

PDB ID : 5T00
Title : Human CTCF ZnF3-7 and methylated DNA complex
Authors : Hashimoto, H.; Wang, D.; Cheng, X.
Deposited on : unknown
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	150	<div> <div>6%</div> <div>89%</div> <div>6%</div> <div>5%</div> </div>
1	D	150	<div> <div>27%</div> <div>91%</div> <div>5%</div> <div>5%</div> </div>
2	B	17	<div> <div>100%</div> </div>
2	E	17	<div> <div>94%</div> <div>6%</div> </div>
3	C	17	<div> <div>94%</div> <div>6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	17	 76% 24%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6860 atoms, of which 3028 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional repressor CTCF.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	143	Total	C	H	N	O	S	0	0	0
			2306	726	1135	228	202	15			
1	D	143	Total	C	H	N	O	S	0	0	0
			2289	723	1121	226	204	15			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	316	GLY	-	expression tag	UNP P49711
A	317	PRO	-	expression tag	UNP P49711
A	318	LEU	-	expression tag	UNP P49711
A	319	GLY	-	expression tag	UNP P49711
A	320	SER	-	expression tag	UNP P49711
D	316	GLY	-	expression tag	UNP P49711
D	317	PRO	-	expression tag	UNP P49711
D	318	LEU	-	expression tag	UNP P49711
D	319	GLY	-	expression tag	UNP P49711
D	320	SER	-	expression tag	UNP P49711

- Molecule 2 is a DNA chain called DNA (5'-TAG(5CM)GCCCCCTGCTGGC-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	17	Total	C	H	N	O	P	0	0	0
			535	163	194	60	102	16			
2	E	17	Total	C	H	N	O	P	0	0	0
			535	163	194	60	102	16			

- Molecule 3 is a DNA chain called DNA (5'-GCCAGCAGGGGG(5CM)GCTA-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	17	Total	C	H	N	O	P	0	0	0
			544	166	192	72	98	16			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	F	17	Total	C	H	N	O	P	0	0	0
			544	166	192	72	98	16			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	Zn	0	0
			5	5		
4	D	5	Total	Zn	0	0
			5	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	28	Total	O	0	0
			28	28		
5	B	22	Total	O	0	0
			22	22		
5	C	13	Total	O	0	0
			13	13		
5	D	15	Total	O	0	0
			15	15		
5	E	10	Total	O	0	0
			10	10		
5	F	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcriptional repressor CTCF



- Molecule 1: Transcriptional repressor CTCF



- Molecule 2: DNA (5'-TAG(5CM)GCCCCCTGCTGGC-3')



There are no outlier residues recorded for this chain.


- Molecule 2: DNA (5'-TAG(5CM)GCCCCCTGCTGGC-3')



- Molecule 3: DNA (5'-GCCAGCAGGGGG(5CM)GCTA-3')



- Molecule 3: DNA (5'-GCCAGCAGGGGG(5CM)GCTA-3')

Chain F:  76% 24%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	40.99 Å 44.91 Å 86.80 Å 98.33° 92.40° 94.80°	Depositor
Resolution (Å)	41.96 – 2.19 85.76 – 2.18	Depositor EDS
% Data completeness (in resolution range)	98.7 (41.96-2.19) 92.6 (85.76-2.18)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.18 Å)	Xtriage
Refinement program	PHENIX (dev_2257: ???)	Depositor
R, R_{free}	0.218 , 0.243 0.218 , 0.241	Depositor DCC
R_{free} test set	1520 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	41.6	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6860	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 5CM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/1203	0.43	0/1612
1	D	0.24	0/1199	0.41	0/1608
2	B	0.53	0/357	0.92	0/546
2	E	0.50	0/357	0.90	0/546
3	C	0.54	0/373	0.82	0/573
3	F	0.53	0/373	0.81	0/573
All	All	0.38	0/3862	0.64	0/5458

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1171	1135	1135	5	0
1	D	1168	1121	1121	5	0
2	B	341	194	194	0	0
2	E	341	194	194	1	0
3	C	352	192	192	1	0
3	F	352	192	192	6	0
4	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	5	0	0	0	0
5	A	28	0	0	0	0
5	B	22	0	0	0	0
5	C	13	0	0	0	0
5	D	15	0	0	0	0
5	E	10	0	0	1	0
5	F	9	0	0	2	0
All	All	3832	3028	3028	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:11:DT:OP2	5:E:101:HOH:O	2.06	0.74
1:D:393:LYS:NZ	3:F:10:DG:O6	2.23	0.71
1:A:339:ARG:NH1	3:C:14:DG:N7	2.45	0.61
3:F:2:DC:OP1	5:F:101:HOH:O	2.16	0.58
3:F:10:DG:OP2	5:F:102:HOH:O	2.17	0.57
1:D:396:ARG:NH1	3:F:7:DA:N7	2.52	0.55
1:D:340:HIS:ND1	3:F:13:5CM:OP1	2.41	0.51
1:A:457:ARG:HA	1:A:462:TYR:HB2	1.98	0.46
1:D:416:PHE:CZ	1:D:425:HIS:CG	3.04	0.45
1:A:342:ARG:O	1:A:347:HIS:HA	2.18	0.43
1:D:397:HIS:ND1	3:F:7:DA:OP1	2.47	0.43
1:A:380:GLN:NE2	1:A:381:CYS:O	2.45	0.42
1:A:416:PHE:CZ	1:A:425:HIS:CG	3.08	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/150 (94%)	137 (97%)	4 (3%)	0	100	100
1	D	141/150 (94%)	138 (98%)	3 (2%)	0	100	100
All	All	282/300 (94%)	275 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/137 (96%)	131 (100%)	0	100	100
1	D	130/137 (95%)	129 (99%)	1 (1%)	85	92
All	All	261/274 (95%)	260 (100%)	1 (0%)	93	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	343	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	5CM	B	4	3,2	14,21,22	1.02	2 (14%)	18,30,33	0.75	1 (5%)
3	5CM	C	13	3,2	14,21,22	1.03	2 (14%)	18,30,33	0.86	1 (5%)
2	5CM	E	4	3,2	14,21,22	0.98	2 (14%)	18,30,33	0.71	1 (5%)
3	5CM	F	13	3,2	14,21,22	1.00	2 (14%)	18,30,33	0.82	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5CM	B	4	3,2	-	0/3/21/22	0/2/2/2
3	5CM	C	13	3,2	-	0/3/21/22	0/2/2/2
2	5CM	E	4	3,2	-	0/3/21/22	0/2/2/2
3	5CM	F	13	3,2	-	0/3/21/22	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	13	5CM	C6-C5	-2.35	1.33	1.40
2	E	4	5CM	C6-C5	-2.34	1.33	1.40
2	B	4	5CM	C6-C5	-2.30	1.34	1.40
3	F	13	5CM	C6-C5	-2.28	1.34	1.40
2	E	4	5CM	C5-C4	2.12	1.44	1.41
2	B	4	5CM	C5-C4	2.25	1.44	1.41
3	F	13	5CM	C5-C4	2.25	1.44	1.41
3	C	13	5CM	C5-C4	2.31	1.44	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	13	5CM	C5A-C5-C4	2.09	123.80	121.65
2	E	4	5CM	C5A-C5-C4	2.31	124.03	121.65
2	B	4	5CM	C5A-C5-C4	2.31	124.03	121.65
3	C	13	5CM	C5A-C5-C4	2.39	124.11	121.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	13	5CM	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	143/150 (95%)	0.77	9 (6%) 21 19	29, 52, 103, 126	0
1	D	143/150 (95%)	1.53	40 (27%) 1 0	31, 85, 130, 145	0
2	B	16/17 (94%)	0.20	0 100 100	39, 50, 83, 88	0
2	E	16/17 (94%)	0.18	0 100 100	42, 63, 94, 96	0
3	C	16/17 (94%)	0.28	0 100 100	32, 39, 86, 90	0
3	F	16/17 (94%)	0.27	0 100 100	33, 67, 94, 104	0
All	All	350/368 (95%)	0.98	49 (14%) 3 3	29, 63, 116, 145	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	462	TYR	6.2
1	A	332	VAL	6.0
1	D	357	ASP	5.9
1	D	370	ILE	5.5
1	D	330	ALA	5.4
1	D	332	VAL	5.2
1	A	320	SER	4.8
1	D	366	LEU	4.7
1	D	367	LYS	4.2
1	D	382	SER	4.1
1	D	363	VAL	3.9
1	D	335	GLY	3.9
1	D	426	ILE	3.5
1	D	463	ILE	3.4
1	D	331	PHE	3.3
1	D	394	LEU	3.2
1	D	341	ARG	3.2
1	D	372	SER	3.2
1	D	411	ILE	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	386	TYR	3.0
1	D	346	THR	2.9
1	D	387	ALA	2.9
1	D	378	PRO	2.8
1	A	445	VAL	2.8
1	D	379	PHE	2.7
1	D	437	PHE	2.6
1	D	388	SER	2.6
1	D	354	SER	2.6
1	A	323	LYS	2.6
1	D	352	LYS	2.6
1	D	398	MET	2.5
1	D	400	THR	2.5
1	D	393	LYS	2.5
1	D	389	ARG	2.5
1	D	359	ALA	2.4
1	D	337	LEU	2.4
1	D	383	LEU	2.3
1	A	459	GLN	2.3
1	D	373	HIS	2.3
1	D	380	GLN	2.3
1	A	329	MET	2.2
1	A	334	SER	2.2
1	D	449	LYS	2.1
1	D	391	THR	2.1
1	A	338	VAL	2.1
1	D	405	LYS	2.1
1	D	454	VAL	2.1
1	D	390	ASP	2.0
1	A	335	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	5CM	F	13	20/21	0.93	0.15	-	62,74,92,93	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	5CM	E	4	20/21	0.86	0.15	-	88,104,122,125	0
2	5CM	B	4	20/21	0.87	0.15	-	67,72,87,87	0
3	5CM	C	13	20/21	0.95	0.16	-	60,73,96,97	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ZN	A	502	1/1	0.99	0.15	1.42	46,46,46,46	0
4	ZN	A	503	1/1	0.98	0.17	1.15	45,45,45,45	0
4	ZN	D	505	1/1	0.97	0.17	0.69	39,39,39,39	0
4	ZN	A	504	1/1	0.99	0.17	0.47	38,38,38,38	0
4	ZN	A	505	1/1	0.99	0.17	-0.23	47,47,47,47	0
4	ZN	D	504	1/1	0.99	0.13	-0.93	54,54,54,54	0
4	ZN	D	501	1/1	0.97	0.08	-1.68	89,89,89,89	0
4	ZN	A	501	1/1	0.98	0.10	-1.88	72,72,72,72	0
4	ZN	D	502	1/1	0.93	0.07	-2.34	93,93,93,93	0
4	ZN	D	503	1/1	0.95	0.05	-2.51	74,74,74,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.