



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 07:07 am GMT

PDB ID : 1T2W
Title : Crystal Structure of Sortase A in Complex with a LPETG peptide
Authors : Zong, Y.; Bice, T.W.; Ton-That, H.; Schneewind, O.; Narayana, S.V.
Deposited on : 2004-04-23
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	145	<div> <div>12%</div> <div> <div></div> <div>71%</div> <div>26%</div> <div>••</div> </div> </div>
1	B	145	<div> <div>3%</div> <div> <div></div> <div>69%</div> <div>21%</div> <div>• 9%</div> </div> </div>
1	C	145	<div> <div>5%</div> <div> <div></div> <div>78%</div> <div>22%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PRO	A	332	-	-	X	X
2	GLU	A	333	-	-	X	X
2	GLY	A	335	-	-	X	X

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3622 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called sortase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	145	Total	C	N	O	S	0	0	0
			1154	730	195	227	2			
1	B	132	Total	C	N	O	S	0	0	0
			1054	669	177	206	2			
1	C	145	Total	C	N	O	S	0	0	0
			1154	730	195	227	2			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	184	ALA	CYS	ENGINEERED	GB 5726436
B	184	ALA	CYS	ENGINEERED	GB 5726436
C	184	ALA	CYS	ENGINEERED	GB 5726436

- Molecule 2 is LEUCINE (three-letter code: GLY, PRO, LEU, THR, GLU) (formula: $C_2H_5NO_2$, $C_5H_9NO_2$, $C_6H_{13}NO_2$, $C_4H_9NO_3$, $C_5H_9NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	5	Total	C	N	O	0	0
			36	22	5	9		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	73	Total	O	0	0
			73	73		
3	B	67	Total	O	0	0
			67	67		
3	C	84	Total	O	0	0
			84	84		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: sortase



- Molecule 1: sortase



- Molecule 1: sortase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.64Å 82.67Å 56.29Å 90.00° 103.96° 90.00°	Depositor
Resolution (Å)	27.31 – 1.80 27.31 – 1.60	Depositor EDS
% Data completeness (in resolution range)	74.8 (27.31-1.80) 84.1 (27.31-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.83 (at 1.61Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.186 , 0.230 0.188 , 0.230	Depositor DCC
R_{free} test set	2806 reflections (9.97%)	DCC
Wilson B-factor (Å ²)	20.1	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 61.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3622	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/1176	0.62	0/1585
1	B	0.31	0/1074	0.57	0/1448
1	C	0.30	0/1176	0.59	0/1585
All	All	0.31	0/3426	0.60	0/4618

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1154	0	1159	44	0
1	B	1054	0	1056	26	0
1	C	1154	0	1159	22	0
2	A	36	0	33	16	0
3	A	73	0	0	2	0
3	B	67	0	0	4	0
3	C	84	0	0	6	0
All	All	3622	0	3407	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:PRO:HB3	1:A:146:VAL:HG11	1.50	0.93
1:A:104:ALA:CA	2:A:335:GLY:OXT	2.16	0.93
1:A:104:ALA:HA	2:A:335:GLY:OXT	1.70	0.91
1:A:104:ALA:CB	2:A:335:GLY:OXT	2.20	0.89
1:C:91:PRO:HD3	1:C:106:GLU:HB2	1.61	0.82
1:A:194:TRP:O	1:A:197:ARG:HD3	1.83	0.79
2:A:332:PRO:O	2:A:333:GLU:CG	2.31	0.79
2:A:332:PRO:O	2:A:333:GLU:CB	2.31	0.77
1:A:104:ALA:HB1	2:A:335:GLY:OXT	1.81	0.77
1:B:96:GLN:HE22	1:B:99:ARG:NH1	1.85	0.75
1:A:91:PRO:HD3	1:A:106:GLU:HB2	1.69	0.73
1:A:175:LYS:O	1:A:176:ASP:HB2	1.88	0.73
1:B:96:GLN:HE22	1:B:99:ARG:HH11	1.33	0.73
1:B:98:ASN:HB3	1:C:206:LYS:NZ	2.08	0.69
1:A:171:GLU:HB2	1:A:178:GLN:OE1	1.95	0.67
2:A:332:PRO:O	2:A:333:GLU:HB2	1.94	0.67
1:A:92:ALA:HB3	2:A:335:GLY:C	2.16	0.65
1:B:103:PHE:HZ	1:B:144:PHE:HZ	1.45	0.62
1:C:114:ASN:HD22	1:C:178:GLN:HG3	1.65	0.62
1:A:171:GLU:OE2	1:A:203:THR:HG21	2.00	0.61
1:A:146:VAL:HG12	1:A:147:GLY:N	2.16	0.61
1:B:98:ASN:HB3	1:C:206:LYS:HZ2	1.63	0.61
1:B:103:PHE:CZ	1:B:144:PHE:HZ	2.19	0.61
1:A:109:SER:HB3	1:A:111:ASP:OD1	2.01	0.60
1:A:162:LYS:HB2	1:A:166:VAL:CG2	2.32	0.59
1:B:151:ARG:HH22	1:B:177:LYS:HD3	1.67	0.59
1:A:146:VAL:HG12	1:A:147:GLY:H	1.68	0.59
1:B:121:THR:O	1:B:121:THR:HG23	2.03	0.59
1:C:169:LEU:O	1:C:173:LYS:HD3	2.04	0.58
1:A:158:ILE:HG13	1:A:170:ASP:HB2	1.84	0.58
1:C:154:LYS:HE3	1:C:205:VAL:HG21	1.83	0.58
1:A:120:HIS:HE1	3:A:353:HOH:O	1.87	0.57
1:A:148:ASN:HB2	3:B:269:HOH:O	2.06	0.56
1:B:188:ASN:C	1:B:188:ASN:HD22	2.09	0.55
1:C:172:GLN:H	1:C:178:GLN:HE22	1.53	0.55
1:B:205:VAL:O	1:B:206:LYS:OXT	2.25	0.55
1:A:145:LYS:HE3	3:C:225:HOH:O	2.05	0.55
1:C:159:ARG:HD3	3:C:246:HOH:O	2.07	0.54
1:A:114:ASN:HD22	1:A:178:GLN:HG3	1.73	0.53
1:C:153:TYR:CZ	1:C:179:LEU:HD13	2.44	0.53
1:C:121:THR:HG23	1:C:121:THR:O	2.09	0.53
1:B:95:GLU:HG3	3:B:214:HOH:O	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176:ASP:HB3	3:A:363:HOH:O	2.08	0.52
1:A:158:ILE:HG12	1:A:170:ASP:OD2	2.10	0.52
1:B:191:THR:OG1	1:B:193:VAL:HG22	2.11	0.51
1:A:197:ARG:HD2	2:A:334:THR:O	2.11	0.51
1:B:109:SER:HB3	1:B:112:ASP:OD1	2.11	0.51
1:C:95:GLU:HB2	3:C:233:HOH:O	2.11	0.50
1:B:151:ARG:HH22	1:B:177:LYS:CD	2.24	0.49
1:A:171:GLU:HB3	1:A:174:GLY:O	2.13	0.48
1:B:65:ILE:HD13	1:B:108:GLU:O	2.13	0.48
1:B:177:LYS:N	3:B:258:HOH:O	2.47	0.47
2:A:332:PRO:C	2:A:333:GLU:HG2	2.35	0.47
1:A:161:VAL:HB	1:A:166:VAL:HG12	1.97	0.47
1:A:163:PRO:HD3	1:A:197:ARG:O	2.14	0.46
1:B:182:ILE:HD12	1:B:182:ILE:N	2.29	0.46
1:B:166:VAL:HG21	1:B:199:ILE:HD11	1.97	0.46
1:A:169:LEU:HD23	1:A:169:LEU:H	1.80	0.46
1:C:113:GLN:HA	1:C:177:LYS:CD	2.45	0.46
1:C:114:ASN:HD22	1:C:178:GLN:CG	2.26	0.46
1:C:158:ILE:HG13	1:C:170:ASP:HB3	1.98	0.46
1:A:175:LYS:O	1:A:176:ASP:CB	2.62	0.45
2:A:332:PRO:C	2:A:333:GLU:CG	2.85	0.45
1:B:76:ILE:HD11	1:B:142:VAL:CG1	2.46	0.45
1:C:183:THR:OG1	1:C:198:LYS:HB2	2.17	0.45
1:A:156:THR:HG21	1:A:203:THR:HG23	1.98	0.45
1:A:114:ASN:ND2	1:A:178:GLN:HG3	2.32	0.45
1:A:162:LYS:HB2	1:A:166:VAL:HG21	1.99	0.44
1:A:118:ALA:HB2	2:A:335:GLY:HA2	1.99	0.44
2:A:332:PRO:O	2:A:333:GLU:HG2	2.15	0.44
1:C:172:GLN:H	1:C:178:GLN:NE2	2.15	0.43
1:B:113:GLN:NE2	1:B:177:LYS:HG2	2.32	0.43
1:A:205:VAL:O	1:A:206:LYS:HB2	2.18	0.43
1:C:163:PRO:HD2	3:C:262:HOH:O	2.17	0.43
1:A:66:PRO:HD2	1:A:73:ALA:HB2	1.99	0.43
1:B:138:LYS:HD2	1:B:157:SER:OG	2.19	0.43
2:A:332:PRO:O	2:A:333:GLU:HG3	2.15	0.43
1:A:113:GLN:HA	1:A:177:LYS:HD3	2.00	0.43
1:A:156:THR:CG2	1:A:203:THR:HG23	2.49	0.42
1:C:120:HIS:HE1	3:C:215:HOH:O	2.02	0.42
1:B:110:LEU:HD11	1:B:151:ARG:HD3	2.02	0.42
1:C:75:TYR:CZ	1:C:86:PRO:HB3	2.55	0.42
1:B:113:GLN:HA	1:B:113:GLN:NE2	2.35	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:162:LYS:NZ	1:C:162:LYS:HB2	2.34	0.42
1:A:69:LYS:NZ	1:A:106:GLU:HG3	2.35	0.42
1:C:193:VAL:HG13	3:C:260:HOH:O	2.19	0.42
2:A:333:GLU:C	2:A:334:THR:HG23	2.40	0.41
1:A:205:VAL:O	1:A:205:VAL:HG12	2.20	0.41
1:B:112:ASP:HB3	3:B:255:HOH:O	2.20	0.41
1:A:97:LEU:HA	1:A:100:GLY:O	2.20	0.41
1:B:78:ILE:HD11	1:B:181:LEU:HD21	2.02	0.41
1:B:85:GLU:HA	1:B:86:PRO:HD3	1.93	0.41
1:A:162:LYS:HB2	1:A:166:VAL:HG23	2.01	0.40
1:A:143:TYR:N	1:A:143:TYR:CD1	2.89	0.40
1:A:194:TRP:O	1:A:197:ARG:CD	2.61	0.40
1:A:197:ARG:NE	2:A:334:THR:HA	2.36	0.40
1:A:109:SER:HB2	1:A:112:ASP:OD2	2.20	0.40
1:C:97:LEU:HD12	1:C:194:TRP:CZ2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/145 (99%)	135 (94%)	6 (4%)	2 (1%)	13	3
1	B	128/145 (88%)	123 (96%)	5 (4%)	0	100	100
1	C	143/145 (99%)	138 (96%)	5 (4%)	0	100	100
All	All	414/435 (95%)	396 (96%)	16 (4%)	2 (0%)	25	17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	169	LEU
1	A	176	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/127 (100%)	123 (97%)	4 (3%)	45	29
1	B	116/127 (91%)	113 (97%)	3 (3%)	51	36
1	C	127/127 (100%)	124 (98%)	3 (2%)	54	40
All	All	370/381 (97%)	360 (97%)	10 (3%)	50	35

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	120	HIS
1	A	165	ASP
1	A	169	LEU
1	A	171	GLU
1	B	165	ASP
1	B	181	LEU
1	B	188	ASN
1	C	107	ASN
1	C	171	GLU
1	C	188	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	ASN
1	A	120	HIS
1	B	96	GLN
1	B	113	GLN
1	B	120	HIS
1	B	188	ASN
1	C	98	ASN
1	C	107	ASN
1	C	113	GLN
1	C	114	ASN
1	C	120	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	178	GLN
1	C	188	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LEU	A	331	2	7,7,8	0.57	0	6,8,10	0.56	0
2	PRO	A	332	2	6,7,8	1.03	1 (16%)	7,8,10	2.06	1 (14%)
2	GLU	A	333	2	5,8,9	1.21	1 (20%)	2,9,11	0.57	0
2	THR	A	334	2	6,6,7	2.55	1 (16%)	6,7,9	3.09	2 (33%)
2	GLY	A	335	2	1,4,4	0.43	0	0,4,4	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LEU	A	331	2	-	0/4/6/8	0/0/0/0
2	PRO	A	332	2	-	0/0/9/11	0/1/1/1
2	GLU	A	333	2	-	0/3/7/9	0/0/0/0
2	THR	A	334	2	-	0/4/6/8	0/0/0/0
2	GLY	A	335	2	-	0/0/2/2	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	332	PRO	CA-C	2.27	1.53	1.50
2	A	333	GLU	CA-C	2.39	1.53	1.50
2	A	334	THR	CA-C	6.14	1.58	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	334	THR	O-C-CA	-2.80	118.63	125.15
2	A	332	PRO	CB-CA-C	4.73	119.22	112.70
2	A	334	THR	C-CA-N	6.75	123.48	109.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	332	PRO	7	0
2	A	333	GLU	8	0
2	A	334	THR	3	0
2	A	335	GLY	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	145/145 (100%)	0.85	17 (11%) 5 4	14, 25, 108, 137	0
1	B	132/145 (91%)	0.06	4 (3%) 51 45	12, 24, 65, 95	0
1	C	145/145 (100%)	0.09	7 (4%) 31 26	14, 24, 69, 112	0
All	All	422/435 (97%)	0.34	28 (6%) 19 16	12, 25, 78, 137	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	169	LEU	16.8
1	A	170	ASP	11.3
1	A	173	LYS	10.9
1	A	166	VAL	10.5
1	A	168	VAL	9.3
1	A	167	GLY	8.3
1	C	62	LYS	8.0
1	A	171	GLU	7.8
1	A	206	LYS	7.3
1	A	174	GLY	7.1
1	A	165	ASP	6.9
1	A	62	LYS	6.3
1	B	65	ILE	6.3
1	A	175	LYS	6.3
1	C	206	LYS	5.8
1	A	164	THR	5.4
1	A	63	PRO	5.1
1	C	190	LYS	5.1
1	B	206	LYS	4.6
1	C	191	THR	4.4
1	A	172	GLN	4.4
1	B	67	LYS	3.1
1	A	176	ASP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	189	GLU	2.8
1	B	110	LEU	2.7
1	C	63	PRO	2.6
1	C	192	GLY	2.4
1	A	146	VAL	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q < 0.9
2	GLY	A	335	5/5	0.82	0.49	17.34	90,91,91,105	0
2	GLU	A	333	9/10	0.46	0.38	12.58	55,95,97,146	0
2	PRO	A	332	7/8	0.44	0.70	1.54	65,70,119,119	0
2	THR	A	334	7/8	0.59	0.43	-	76,77,89,90	0
2	LEU	A	331	8/9	0.30	0.51	-	70,87,141,143	0

6.5 Other polymers [i](#)

There are no such residues in this entry.