



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 10, 2017 – 02:59 PM EDT

PDB ID : 5T2G
Title : mPI3Kd IN COMPLEX WITH γ i
Authors : Petersen, J.; Terstige, I.; Perry, M.; Svensson, T.; Tyrchan, C.; Lindmark, H.; Oster, L.
Deposited on : 2016-08-23
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029077
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029077


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4993 (2.58-2.50)
Clashscore	112137	5755 (2.58-2.50)
Ramachandran outliers	110173	5652 (2.58-2.50)
Sidechain outliers	110143	5654 (2.58-2.50)
RSRZ outliers	101464	5026 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	939	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit delta isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	820	Total	C	N	O	S	0	0	0
			6618	4246	1124	1194	54			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	508	GLN	-	insertion	UNP O35904

- Molecule 2 is 4-[3-azanyl-6-[1-methyl-5-(1-phenylcyclopropyl)-1,2,4-triazol-3-yl]pyrazin-2-yl]pyrazole-1-carboxamide (three-letter code: 74K) (formula: C₂₀H₁₉N₉O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			30	20	9	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	49	Total	O	0	0
			49	49		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	64.59Å 142.16Å 221.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.84 – 2.55 56.84 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.9 (56.84-2.55) 99.9 (56.84-2.55)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.55Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.234 , 0.278 0.247 , 0.293	Depositor DCC
R_{free} test set	1688 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	46.0	Xtriage
Anisotropy	0.268	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 57.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6697	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 74K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/6760	0.66	0/9118

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6618	0	6604	29	0
2	A	30	0	0	0	0
3	A	49	0	0	0	0
All	All	6697	0	6604	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:549:LEU:HG	1:A:564:MET:CE	2.27	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:549:LEU:HG	1:A:564:MET:HE1	1.82	0.62
1:A:784:LEU:HD12	1:A:823:GLY:HA3	1.85	0.58
1:A:756:MET:HG2	1:A:780:ASN:O	2.05	0.56
1:A:162:TRP:CE3	1:A:286:ARG:HG3	2.41	0.55
1:A:317:TRP:HA	1:A:382:CYS:HB2	1.89	0.55
1:A:553:THR:HG21	1:A:564:MET:HG2	1.89	0.54
1:A:1002:LEU:HB3	1:A:1004:LEU:HD23	1.89	0.54
1:A:756:MET:HG2	1:A:781:GLY:HA3	1.91	0.53
1:A:834:ILE:HD11	1:A:901:ILE:HD11	1.90	0.52
1:A:192:VAL:HG13	1:A:272:PRO:HB2	1.91	0.52
1:A:883:CYS:HB3	1:A:932:PHE:CZ	2.45	0.51
1:A:326:GLU:HB3	1:A:474:TYR:HB3	1.94	0.49
1:A:512:ARG:HG2	1:A:530:LEU:HD12	1.94	0.49
1:A:568:LEU:HA	1:A:571:TRP:HB2	1.94	0.49
1:A:154:ARG:HH11	1:A:154:ARG:HG2	1.78	0.48
1:A:135:GLU:HG3	1:A:428:TYR:CG	2.48	0.47
1:A:113:ILE:HA	1:A:116:GLN:HE21	1.79	0.47
1:A:549:LEU:HG	1:A:564:MET:HE2	1.96	0.47
1:A:647:LEU:O	1:A:651:LEU:HG	2.18	0.44
1:A:620:LYS:HG2	1:A:660:VAL:HG11	2.00	0.43
1:A:169:LEU:HD22	1:A:259:PHE:HE1	1.83	0.43
1:A:808:LEU:HD11	1:A:963:ALA:HB2	2.01	0.43
1:A:340:LYS:HG2	1:A:362:GLU:HB3	2.00	0.43
1:A:209:THR:HB	1:A:257:CYS:HB3	2.00	0.42
1:A:886:THR:HA	1:A:891:ILE:HD12	2.01	0.42
1:A:971:GLY:HA3	1:A:1004:LEU:HD21	2.01	0.42
1:A:278:HIS:CD2	1:A:280:SER:H	2.39	0.41
1:A:870:ARG:HD2	1:A:870:ARG:HA	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	796/939 (85%)	764 (96%)	29 (4%)	3 (0%)	38 57

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	328	ILE
1	A	742	GLU
1	A	755	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	726/827 (88%)	677 (93%)	49 (7%)	19 33

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	LEU
1	A	154	ARG
1	A	203	PHE
1	A	216	LEU
1	A	247	HIS
1	A	270	LEU
1	A	316	LEU
1	A	317	TRP
1	A	329	GLU
1	A	331	ARG
1	A	332	LYS
1	A	340	LYS
1	A	394	LEU
1	A	398	VAL
1	A	423	LEU
1	A	439	LEU
1	A	471	LEU
1	A	472	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	475	LEU
1	A	483	VAL
1	A	511	LEU
1	A	512	ARG
1	A	514	ILE
1	A	516	GLU
1	A	517	ARG
1	A	523	LEU
1	A	530	LEU
1	A	548	ARG
1	A	553	THR
1	A	586	SER
1	A	617	GLN
1	A	634	LEU
1	A	722	GLU
1	A	731	LEU
1	A	748	GLN
1	A	756	MET
1	A	757	LYS
1	A	841	LYS
1	A	855	LEU
1	A	898	ASN
1	A	915	PHE
1	A	919	PHE
1	A	930	VAL
1	A	951	GLU
1	A	962	ARG
1	A	986	LEU
1	A	998	LEU
1	A	1004	LEU
1	A	1027	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	GLN
1	A	170	GLN
1	A	273	HIS
1	A	278	HIS
1	A	344	GLN
1	A	773	ASN
1	A	780	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	898	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	74K	A	1101	-	32,34,34	1.15	2 (6%)	33,51,51	3.07	10 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	74K	A	1101	-	-	0/10/28/28	0/4/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	74K	C14-N13	-2.90	1.30	1.34
2	A	1101	74K	N16-N15	3.11	1.40	1.37

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	74K	N16-C12-N13	-10.17	108.30	114.52
2	A	1101	74K	C7-C8-N9	-3.49	117.09	120.45
2	A	1101	74K	O30-C28-N29	-3.38	116.46	122.65
2	A	1101	74K	C4-C5-N6	-2.58	118.93	122.38
2	A	1101	74K	N13-C14-N15	-2.45	108.16	113.51
2	A	1101	74K	C12-N16-N15	2.30	107.20	103.11
2	A	1101	74K	C5-N6-C7	2.60	121.74	118.84
2	A	1101	74K	C8-C7-N10	2.98	123.48	121.12
2	A	1101	74K	C27-N26-N25	8.09	109.47	103.70
2	A	1101	74K	C12-N13-C14	8.15	110.35	101.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	820/939 (87%)	0.40	46 (5%) 25 27	17, 43, 81, 149	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	357	THR	4.9
1	A	334	ASN	4.1
1	A	373	GLN	4.0
1	A	330	GLY	3.9
1	A	347	LEU	3.9
1	A	470	ALA	3.9
1	A	1027	TRP	3.9
1	A	530	LEU	3.8
1	A	843	ASN	3.7
1	A	517	ARG	3.6
1	A	846	ALA	3.6
1	A	418	ILE	3.5
1	A	396	ALA	3.3
1	A	844	MET	3.2
1	A	341	LEU	3.2
1	A	366	CYS	3.1
1	A	511	LEU	3.0
1	A	945	GLY	3.0
1	A	317	TRP	3.0
1	A	190	LEU	2.9
1	A	226	THR	2.9
1	A	343	VAL	2.9
1	A	332	LYS	2.9
1	A	372	LYS	2.8
1	A	381	VAL	2.7
1	A	492	LEU	2.6
1	A	514	ILE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	936	TYR	2.6
1	A	206	GLN	2.6
1	A	515	LEU	2.5
1	A	325	ILE	2.5
1	A	1025	GLU	2.5
1	A	841	LYS	2.5
1	A	395	TYR	2.3
1	A	316	LEU	2.3
1	A	173	PRO	2.3
1	A	441	MET	2.2
1	A	1004	LEU	2.2
1	A	840	ASN	2.1
1	A	394	LEU	2.1
1	A	532	TRP	2.1
1	A	153	HIS	2.1
1	A	270	LEU	2.1
1	A	377	PHE	2.0
1	A	919	PHE	2.0
1	A	479	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	74K	A	1101	30/30	0.93	0.21	1.61	19,27,50,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.