



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2017 – 11:19 AM EDT

PDB ID : 1T40
Title : Crystal structure of human aldose reductase complexed with NADP and
IDD552 at ph 5
Authors : Ruiz, F.; Hazemann, I.; Mitschler, A.; Chevrier, B.; Schneider, T.; Joachimiak,
A.; Karplus, M.; Podjarny, A.
Deposited on : unknown
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div>2%</div> <div>83%</div> <div>15%</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldose reductase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	316	2525	1622	426	464	13	0	0	0

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0

- Molecule 3 is [5-FLUORO-2-({[(4,5,7-TRIFLUORO-1,3-BENZOTHAZOL-2-YL)METHYL]AMINO}CARBONYL)PHENOXY]ACETIC ACID (three-letter code: ID5) (formula: C₁₇H₁₀F₄N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	S	
			28	17	4	2	4	1	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	263	Total	O		
			263	263	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldose reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.00Å 67.24Å 47.78Å 90.00° 92.50° 90.00°	Depositor
Resolution (Å)	7.99 – 1.80 15.67 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.3 (7.99-1.80) 98.3 (15.67-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 1.80Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.159 , 0.193 0.158 , 0.191	Depositor DCC
R_{free} test set	2845 reflections (11.05%)	DCC
Wilson B-factor (Å ²)	14.2	Xtriage
Anisotropy	0.132	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for -l,k,h 0.043 for -h,-k,l 0.031 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2864	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, ID5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2587	0.69	0/3512

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	209	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2525	0	2542	34	0
2	A	48	0	25	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	28	0	9	1	0
4	A	263	0	0	3	0
All	All	2864	0	2576	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:GLU:OE1	1:A:223:GLU:HA	1.52	1.08
1:A:229:GLU:HA	1:A:234:LYS:HE3	1.63	0.81
1:A:60:GLU:HG2	4:A:573:HOH:O	1.81	0.79
1:A:271:GLU:HA	1:A:274:LYS:HE3	1.63	0.79
1:A:4:LEU:HD12	1:A:5:LEU:H	1.61	0.65
1:A:217:ARG:HG2	1:A:220:ALA:HB2	1.81	0.63
1:A:187:HIS:HD2	1:A:189:TYR:H	1.46	0.63
1:A:305:SER:HB2	4:A:763:HOH:O	2.00	0.61
1:A:20:TRP:O	1:A:21:LYS:HB2	2.06	0.56
1:A:59:GLN:HG2	4:A:550:HOH:O	2.04	0.56
1:A:4:LEU:HD12	1:A:5:LEU:N	2.23	0.54
1:A:136:ASN:ND2	1:A:138:LEU:HB2	2.23	0.54
1:A:185:GLU:OE2	1:A:187:HIS:HE1	1.94	0.51
1:A:242:LYS:HE3	1:A:278:PHE:CD1	2.47	0.50
1:A:185:GLU:HB2	1:A:209:TYR:CE1	2.46	0.49
1:A:210:SER:N	1:A:211:PRO:CD	2.76	0.48
1:A:136:ASN:HD22	1:A:138:LEU:H	1.62	0.48
1:A:3:ARG:HG2	1:A:13:PRO:HA	1.95	0.48
1:A:167:GLU:HG2	1:A:171:ASN:ND2	2.29	0.47
1:A:234:LYS:HA	1:A:244:THR:HG22	1.98	0.46
1:A:206:VAL:HB	1:A:257:LEU:HD22	1.98	0.45
1:A:50:ASN:C	1:A:50:ASN:HD22	2.20	0.45
1:A:223:GLU:OE1	1:A:223:GLU:CA	2.43	0.44
1:A:251:PHE:HB3	1:A:252:PRO:CD	2.48	0.44
1:A:217:ARG:CG	1:A:220:ALA:HB2	2.46	0.44
1:A:234:LYS:HG2	1:A:244:THR:HG21	2.00	0.43
1:A:7:ASN:C	1:A:7:ASN:HD22	2.22	0.43
1:A:222:PRO:O	1:A:223:GLU:OE1	2.38	0.42
1:A:20:TRP:CE3	1:A:21:LYS:HE3	2.54	0.41
1:A:113:THR:OG1	3:A:320:ID5:H28	2.21	0.41
1:A:146:GLU:O	1:A:150:GLU:HG3	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:251:PHE:HB3	1:A:252:PRO:HD3	2.01	0.41
1:A:202:LYS:HA	1:A:202:LYS:HD3	1.84	0.40
1:A:20:TRP:O	1:A:21:LYS:CB	2.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	314/316 (99%)	308 (98%)	5 (2%)	1 (0%)	44 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	222	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	281/281 (100%)	262 (93%)	19 (7%)	18 6

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	6	LEU
1	A	7	ASN
1	A	50	ASN
1	A	65	GLN
1	A	85	LYS
1	A	104	LEU
1	A	119	LYS
1	A	138	LEU
1	A	154	LYS
1	A	175	LEU
1	A	182	ASN
1	A	193	GLU
1	A	221	LYS
1	A	223	GLU
1	A	228	LEU
1	A	288	LEU
1	A	293	ARG
1	A	307	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	93	GLN
1	A	136	ASN
1	A	171	ASN
1	A	182	ASN
1	A	187	HIS
1	A	283	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	A	318	-	44,52,52	1.17	5 (11%)	51,80,80	2.20	8 (15%)
3	ID5	A	320	-	25,30,30	1.96	10 (40%)	28,43,43	1.37	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	318	-	-	0/27/67/67	0/5/5/5
3	ID5	A	320	-	-	0/10/14/14	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	318	NAP	C5A-N7A	-2.17	1.32	1.39
3	A	320	ID5	C6-C3	2.12	1.42	1.38
3	A	320	ID5	C7-C4	2.21	1.44	1.40
2	A	318	NAP	C3N-C7N	2.21	1.54	1.50
2	A	318	NAP	O4B-C1B	2.22	1.44	1.41
3	A	320	ID5	C3-C7	2.29	1.43	1.39
2	A	318	NAP	O4D-C1D	2.31	1.44	1.41
3	A	320	ID5	C2-C5	2.37	1.41	1.37
3	A	320	ID5	C2-C4	2.60	1.43	1.38
3	A	320	ID5	C11-N36	2.84	1.40	1.33
3	A	320	ID5	C25-C29	2.89	1.43	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	320	ID5	C6-C5	3.58	1.44	1.37
3	A	320	ID5	C28-C27	3.71	1.42	1.36
3	A	320	ID5	C28-C25	3.79	1.44	1.37
2	A	318	NAP	C6N-N1N	3.85	1.45	1.35

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	318	NAP	N3A-C2A-N1A	-11.36	118.96	128.86
3	A	320	ID5	C25-C29-C26	-2.87	118.57	120.94
3	A	320	ID5	C28-C27-C24	-2.80	120.08	123.85
2	A	318	NAP	C6N-C5N-C4N	-2.70	115.37	119.44
3	A	320	ID5	C6-C5-C2	-2.03	120.62	123.29
2	A	318	NAP	O5D-C5D-C4D	2.07	116.35	109.00
2	A	318	NAP	C2A-N1A-C6A	2.84	123.74	118.77
2	A	318	NAP	O3D-C3D-C2D	3.05	121.61	111.83
2	A	318	NAP	O3D-C3D-C4D	3.17	120.35	111.09
3	A	320	ID5	C27-C24-S22	3.95	131.41	124.70
2	A	318	NAP	C4A-C5A-N7A	4.07	113.35	109.41
2	A	318	NAP	C5N-C4N-C3N	4.97	126.19	120.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	320	ID5	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/316 (100%)	-0.32	5 (1%) 72 69	6, 13, 28, 49	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	222	PRO	5.1
1	A	223	GLU	4.7
1	A	221	LYS	3.1
1	A	1(B)	ALA	2.7
1	A	1(A)	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ID5	A	320	28/28	0.97	0.09	-0.25	10,12,16,16	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAP	A	318	48/48	0.98	0.07	-0.87	5,9,13,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.