



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 12, 2017 – 02:02 PM EDT

PDB ID : 5T7X
Title : Crystal structure of HHV-4 EBNA1 DNA binding domain (patient-derived, nasopharyngeal carcinoma) bound to DNA
Authors : Malecka, K.A.; Messick, T.E.; Lieberman, P.M.
Deposited on : unknown
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20029824
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1522 (2.38-2.34)
Clashscore	112137	1626 (2.38-2.34)
Ramachandran outliers	110173	1605 (2.38-2.34)
Sidechain outliers	110143	1606 (2.38-2.34)
RSRZ outliers	101464	1528 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	18	<div> <div></div> <div>72%</div> <div>28%</div> </div>
2	D	18	<div> <div></div> <div>72%</div> <div>28%</div> </div>
3	A	149	<div> <div>5%</div> <div>86%</div> <div>11%</div> <div>••</div> </div>
3	B	149	<div> <div>5%</div> <div>85%</div> <div>13%</div> <div>•</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3126 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*GP*GP*AP*TP*AP*GP*CP*CP*TP*AP*TP*GP*CP*TP*AP*CP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	18	Total	C	N	O	P	0	0	0
			363	174	66	106	17			

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*GP*GP*TP*AP*GP*CP*AP*TP*AP*GP*GP*CP*TP*AP*TP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	18	Total	C	N	O	P	0	0	0
			369	176	70	106	17			

- Molecule 3 is a protein called Epstein-Barr nuclear antigen 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	147	Total	C	N	O	S	0	0	0
			1128	724	199	198	7			
3	B	149	Total	C	N	O	S	0	0	0
			1144	733	204	200	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	585	ILE	THR	engineered mutation	UNP Q3KSS4
B	585	ILE	THR	engineered mutation	UNP Q3KSS4

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	18	Total	O	0	0
			18	18		
4	D	16	Total	O	0	0
			16	16		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	45	Total	O	0	0
			45	45		
4	B	43	Total	O	0	0
			43	43		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(*GP*GP*AP*TP*AP*GP*CP*CP*TP*AP*TP*GP*CP*TP*AP*CP*CP*C)-3')

Chain C:  72% 28%





- Molecule 2: DNA (5'-D(*GP*GP*GP*TP*AP*GP*CP*AP*TP*AP*GP*GP*CP*TP*AP*TP*CP*C)-3')

Chain D:  72% 28%




- Molecule 3: Epstein-Barr nuclear antigen 1

Chain A:  5% 86% 11%



- Molecule 3: Epstein-Barr nuclear antigen 1

Chain B:  5% 85% 13%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.02Å 64.90Å 111.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.37 – 2.35 23.37 – 2.29	Depositor EDS
% Data completeness (in resolution range)	62.2 (23.37-2.35) 76.9 (23.37-2.29)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.14 (at 2.28Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.185 , 0.233 0.211 , 0.256	Depositor DCC
R_{free} test set	1541 reflections (10.00%)	DCC
Wilson B-factor (Å ²)	19.5	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 17.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3126	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.60	0/406	0.94	0/624
2	D	0.65	0/414	0.96	0/638
3	A	0.30	0/1157	0.47	0/1567
3	B	0.28	0/1173	0.49	0/1588
All	All	0.40	0/3150	0.65	0/4417

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	363	0	204	5	0
2	D	369	0	204	3	0
3	A	1128	0	1143	13	0
3	B	1144	0	1158	13	0
4	A	45	0	0	2	0
4	B	43	0	0	2	0
4	C	18	0	0	0	0
4	D	16	0	0	0	0
All	All	3126	0	2709	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:479:GLU:OE2	3:B:486:ARG:NH2	2.16	0.78
3:A:583:VAL:HG13	3:A:591:CYS:HA	1.77	0.66
3:A:583:VAL:HG22	3:A:590:THR:HG22	1.79	0.63
3:B:576:LYS:HD3	3:B:597:VAL:HG23	1.80	0.63
3:A:479:GLU:OE2	3:A:486:ARG:NH1	2.31	0.61
3:A:491:ARG:NH2	3:A:581:ASP:OD2	2.27	0.61
3:B:583:VAL:HG13	3:B:591:CYS:HA	1.86	0.57
3:A:488:LEU:O	3:A:491:ARG:HD3	2.06	0.56
2:D:214:DT:H2'	2:D:215:DA:C8	2.42	0.54
3:B:459:ARG:N	4:B:707:HOH:O	2.43	0.52
3:B:573:GLU:OE2	3:B:576:LYS:NZ	2.42	0.52
3:A:576:LYS:HD3	3:A:597:VAL:HG23	1.92	0.52
3:B:462:GLY:HA3	3:B:465:PHE:CE1	2.48	0.49
3:A:507:VAL:HG22	3:A:564:VAL:HB	1.94	0.49
3:B:462:GLY:HA3	3:B:465:PHE:HE1	1.77	0.48
2:D:213:DC:H2'	2:D:214:DT:C6	2.49	0.48
1:C:110:DA:H5''	3:A:538:ARG:NH1	2.29	0.47
3:B:555:ARG:NH1	4:B:705:HOH:O	2.42	0.47
1:C:110:DA:H5''	3:A:538:ARG:CZ	2.47	0.44
1:C:113:DC:H2'	1:C:114:DT:C6	2.52	0.44
3:B:521:ARG:HG3	3:B:533:ILE:HG21	1.99	0.44
2:D:209:DT:H1'	2:D:210:DA:C8	2.52	0.44
3:B:586:LYS:O	3:B:591:CYS:HB3	2.18	0.44
3:B:489:LEU:HD12	3:B:574:VAL:HG12	2.00	0.43
3:A:555:ARG:NH2	4:A:703:HOH:O	2.42	0.43
3:B:583:VAL:HG22	3:B:590:THR:HG22	2.01	0.43
1:C:103:DA:C8	3:B:515:THR:HG21	2.54	0.43
3:A:496:ARG:HD3	4:A:726:HOH:O	2.18	0.42
3:A:517:LEU:HD23	3:A:517:LEU:HA	1.89	0.42
1:C:112:DG:H5''	3:A:535:PRO:HB3	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	145/149 (97%)	142 (98%)	3 (2%)	0	100	100
3	B	147/149 (99%)	141 (96%)	5 (3%)	1 (1%)	25	27
All	All	292/298 (98%)	283 (97%)	8 (3%)	1 (0%)	44	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	461	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	121/123 (98%)	116 (96%)	5 (4%)	35	44
3	B	122/123 (99%)	114 (93%)	8 (7%)	19	21
All	All	243/246 (99%)	230 (95%)	13 (5%)	26	32

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	491	ARG
3	A	496	ARG
3	A	539	LEU
3	A	575	LEU
3	A	583	VAL
3	B	459	ARG
3	B	489	LEU
3	B	496	ARG
3	B	539	LEU
3	B	563	MET
3	B	575	LEU
3	B	576	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	B	583	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	C	18/18 (100%)	-0.49	0	100	100	19, 21, 25, 25	0
2	D	18/18 (100%)	-0.62	0	100	100	19, 20, 24, 25	0
3	A	147/149 (98%)	0.12	8 (5%)	26	37	16, 20, 36, 62	0
3	B	149/149 (100%)	0.12	7 (4%)	32	44	16, 22, 35, 53	0
All	All	332/334 (99%)	0.05	15 (4%)	34	46	16, 21, 35, 62	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	546	GLY	4.9
3	B	462	GLY	3.4
3	A	499	GLU	3.3
3	B	500	GLU	3.0
3	A	500	GLU	2.8
3	B	547	PRO	2.6
3	A	598	CYS	2.6
3	A	558	ILE	2.5
3	A	570	ILE	2.4
3	A	545	PRO	2.4
3	B	546	GLY	2.3
3	B	549	PRO	2.3
3	B	545	PRO	2.1
3	A	547	PRO	2.1
3	B	499	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.