



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 12:54 am GMT

PDB ID : 3TDC  
Title : Crystal Structure of Human Acetyl-CoA carboxylase 2  
Authors : Dougan, D.R.; Mol, C.D.  
Deposited on : 2011-08-10  
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3709 (2.44-2.40)
Clashscore	112137	4241 (2.44-2.40)
Ramachandran outliers	110173	4178 (2.44-2.40)
Sidechain outliers	110143	4179 (2.44-2.40)
RSRZ outliers	101464	3740 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	762	<div> <div>6%</div> <div>89%</div> <div>5%</div> <div>6%</div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6284 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Acetyl-CoA carboxylase 2 variant.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	720	5792	3713	990	1065	24	0	9	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2471	HIS	-	EXPRESSION TAG	UNP Q59GJ9
A	2472	HIS	-	EXPRESSION TAG	UNP Q59GJ9
A	2473	HIS	-	EXPRESSION TAG	UNP Q59GJ9
A	2474	HIS	-	EXPRESSION TAG	UNP Q59GJ9
A	2475	HIS	-	EXPRESSION TAG	UNP Q59GJ9
A	2476	HIS	-	EXPRESSION TAG	UNP Q59GJ9

- Molecule 2 is 1-[3-({4-[(5S)-3,3-DIMETHYL-1-OXO-2-OXA-7-AZASPIRO[4.5]DEC-7-YL]P IPERIDIN-1-YL}CARBONYL)-1-BENZOTHIOPHEN-2-YL]-3-ETHYLUREA (three-letter code: 0EU) (formula: C<sub>27</sub>H<sub>36</sub>N<sub>4</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			36	27	4	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	456	Total	O	0	0
			456	456		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetyl-CoA carboxylase 2 variant



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.92Å 119.76Å 146.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.41 29.94 – 2.41	Depositor EDS
% Data completeness (in resolution range)	96.8 (30.00-2.41) 96.8 (29.94-2.41)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.90 (at 2.42Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.169 , 0.218 0.168 , 0.216	Depositor DCC
$R_{free}$ test set	1874 reflections (5.24%)	DCC
Wilson B-factor (Å <sup>2</sup> )	40.9	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 54.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.020 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6284	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0EU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/5967	0.59	0/8088

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5792	0	5759	19	0
2	A	36	0	36	1	0
3	A	456	0	0	2	0
All	All	6284	0	5795	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1991:ASN:HD21	1:A:1999:TYR:H	1.33	0.75

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2386:ARG:O	1:A:2390:GLU:HG2	1.88	0.74
1:A:2402:ASN:HD22	1:A:2405:VAL:H	1.48	0.61
1:A:1868:HIS:CD2	3:A:4040:HOH:O	2.53	0.61
1:A:1868:HIS:HD2	3:A:4040:HOH:O	1.84	0.61
2:A:3000:0EU:O10	2:A:3000:0EU:S7	2.62	0.57
1:A:1727:LEU:HD11	1:A:1788:PRO:HB2	1.89	0.55
1:A:1907:HIS:HE1	1:A:1914:SER:OG	1.90	0.53
1:A:2009:GLN:H	1:A:2009:GLN:NE2	2.07	0.53
1:A:1757:TRP:CD1	1:A:1767:ILE:HD11	2.44	0.52
1:A:1991:ASN:ND2	1:A:1998:VAL:H	2.13	0.47
1:A:1843:ARG:HD3	1:A:1946:ALA:HA	1.98	0.45
1:A:2402:ASN:ND2	1:A:2405:VAL:H	2.11	0.45
1:A:2289:GLU:HA	1:A:2290:PRO:HD3	1.83	0.44
1:A:2364:VAL:HG11	1:A:2384:LEU:HD21	1.98	0.43
1:A:1775:LEU:HG	1:A:1834:LEU:HD12	2.00	0.43
1:A:2316:HIS:O	1:A:2320:VAL:HG23	2.18	0.42
1:A:2042:ASN:C	1:A:2042:ASN:HD22	2.22	0.42
1:A:1947:TYR:CE2	1:A:1972:GLN:HG3	2.56	0.40
1:A:2009:GLN:H	1:A:2009:GLN:HE21	1.68	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	724/762 (95%)	698 (96%)	25 (4%)	1 (0%)	55 71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2449	GLU



### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	624/653 (96%)	606 (97%)	18 (3%)	48 67

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1719	THR
1	A	1728	GLN
1	A	1766	ASP
1	A	1786	ARG
1	A	1997	GLU
1	A	2009	GLN
1	A	2042	ASN
1	A	2130	GLU
1	A	2210	TYR
1	A	2264	PHE
1	A	2274	ARG
1	A	2340	ASP
1	A	2374	GLU
1	A	2375	LEU
1	A	2382	SER
1	A	2402	ASN
1	A	2408	GLN
1	A	2441	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1728	GLN
1	A	1805	GLN
1	A	1818	ASN
1	A	1868	HIS
1	A	1901	ASN
1	A	1907	HIS
1	A	1991	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	2009	GLN
1	A	2042	ASN
1	A	2139	ASN
1	A	2286	GLN
1	A	2316	HIS
1	A	2363	GLN
1	A	2381	GLN
1	A	2402	ASN
1	A	2413	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	0EU	A	3000	-	34,40,40	1.21	3 (8%)	42,60,60	1.71	8 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0EU	A	3000	-	-	0/14/58/58	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	3000	0EU	C14-C15	-2.65	1.50	1.51
2	A	3000	0EU	C14-C8	2.67	1.47	1.41
2	A	3000	0EU	O27-C26	3.85	1.44	1.36

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3000	0EU	C17-C16-N15	-3.58	105.55	110.82
2	A	3000	0EU	C1-C14-C15	-2.33	122.07	125.80
2	A	3000	0EU	O10-C10-N11	-2.25	118.52	122.62
2	A	3000	0EU	C2-C1-C14	-2.15	131.08	135.53
2	A	3000	0EU	C35-N11-C10	2.66	125.45	121.51
2	A	3000	0EU	C14-C15-N15	2.83	120.46	117.70
2	A	3000	0EU	N9-C10-N11	3.87	120.32	113.91
2	A	3000	0EU	C29-C28-C25	6.61	108.13	100.28

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	3000	0EU	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	720/762 (94%)	-0.04	43 (5%)	23 20	18, 28, 62, 88	1 (0%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2290	PRO	6.0
1	A	2450	VAL	5.8
1	A	1759	SER	4.9
1	A	1761	ASP	4.6
1	A	2444	VAL	4.6
1	A	1725	ASP	4.5
1	A	2291	ASP	4.4
1	A	2085	THR	4.2
1	A	2395	VAL	4.1
1	A	2082	LEU	3.9
1	A	2441	ARG	3.5
1	A	1762	LYS	3.5
1	A	1760	PRO	3.2
1	A	2282	LYS	3.2
1	A	1763	TYR	3.1
1	A	2084	GLY	3.1
1	A	1726	LEU	3.1
1	A	1851	TYR	2.9
1	A	2288	GLY	2.9
1	A	2448	PRO	2.9
1	A	2414	TRP	2.7
1	A	1815	VAL	2.7
1	A	2446	GLU	2.7
1	A	2141	ASP	2.7
1	A	2289	GLU	2.6
1	A	2292	LEU	2.6
1	A	2379	HIS	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	2294	ASP	2.6
1	A	1816	ILE	2.6
1	A	1852	VAL	2.6
1	A	1817	GLY	2.5
1	A	2451	ALA	2.4
1	A	2394	ALA	2.4
1	A	2445	GLU	2.4
1	A	2449	GLU	2.2
1	A	1850[A]	ILE	2.2
1	A	1732	PHE	2.2
1	A	2374	GLU	2.1
1	A	1785	ASN	2.1
1	A	2285	GLU	2.1
1	A	1724	LYS	2.1
1	A	2296	ASP	2.0
1	A	2428	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	0EU	A	3000	36/36	0.96	0.10	-0.53	11,18,26,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.