



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2017 – 05:05 PM EDT

PDB ID : 5TDX
Title : Resurrected Ancestral Hydroxynitrile Lyase from Flowering Plants
Authors : Jones, B.J.; Evans, R.; Wilmot, C.M.; Kazlauskas, R.J.
Deposited on : unknown
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	<div> <div>5%</div> <div> <div></div> <div>81%</div> <div>13%</div> <div>• •</div> </div> </div>
1	B	272	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>15%</div> <div>• 5%</div> </div> </div>
1	C	272	<div> <div>6%</div> <div> <div></div> <div>84%</div> <div>12%</div> <div>• •</div> </div> </div>
1	D	272	<div> <div>13%</div> <div> <div></div> <div>79%</div> <div>14%</div> <div>• •</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	300	-	-	-	X
2	GOL	B	301	-	-	-	X

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8741 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ancestral Hydroxynitrile Lyase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	5	0
			2131	1373	341	408	9			
1	B	259	Total	C	N	O	S	0	4	0
			2112	1361	340	402	9			
1	C	265	Total	C	N	O	S	0	2	0
			2138	1380	342	407	9			
1	D	260	Total	C	N	O	S	0	5	0
			2132	1374	344	405	9			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total 6	C 3	O 3	0	0
2	C	1	Total 6	C 3	O 3	0	0
2	C	1	Total 6	C 3	O 3	0	0
2	D	1	Total 6	C 3	O 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	58	Total 58	O 58	0	0
3	B	66	Total 66	O 66	0	0
3	C	54	Total 54	O 54	0	0
3	D	14	Total 14	O 14	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Ancestral Hydroxynitrile Lyase 1




• Molecule 1: Ancestral Hydroxynitrile Lyase 1




• Molecule 1: Ancestral Hydroxynitrile Lyase 1



• Molecule 1: Ancestral Hydroxynitrile Lyase 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	108.27Å 108.27Å 201.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.32 – 1.96 29.32 – 1.96	Depositor EDS
% Data completeness (in resolution range)	99.3 (29.32-1.96) 99.3 (29.32-1.96)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.177 , 0.227 0.186 , 0.230	Depositor DCC
R_{free} test set	4314 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	30.8	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8741	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.43	7/2188 (0.3%)	1.26	10/2969 (0.3%)
1	B	1.51	15/2172 (0.7%)	1.30	22/2947 (0.7%)
1	C	1.36	9/2195 (0.4%)	1.19	14/2980 (0.5%)
1	D	1.16	2/2192 (0.1%)	1.16	13/2974 (0.4%)
All	All	1.37	33/8747 (0.4%)	1.23	59/11870 (0.5%)

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	167	GLU	CD-OE2	8.75	1.35	1.25
1	B	213	GLU	CD-OE2	7.91	1.34	1.25
1	B	164	GLU	CD-OE1	7.71	1.34	1.25
1	A	244	GLU	CD-OE2	6.89	1.33	1.25
1	A	136	TYR	CB-CG	6.85	1.61	1.51
1	A	217	TRP	CE3-CZ3	6.85	1.50	1.38
1	A	167[A]	GLU	CD-OE1	6.78	1.33	1.25
1	A	167[B]	GLU	CD-OE1	6.78	1.33	1.25
1	B	244	GLU	CD-OE1	6.72	1.33	1.25
1	B	211	PRO	N-CA	-6.58	1.36	1.47
1	B	220	GLU	CD-OE2	6.56	1.32	1.25
1	B	192	GLU	CD-OE1	6.53	1.32	1.25
1	A	20	TYR	CE2-CZ	-6.26	1.30	1.38
1	B	213	GLU	CG-CD	6.23	1.61	1.51
1	C	166	TYR	CE1-CZ	-6.04	1.30	1.38
1	C	256	TYR	CG-CD1	5.99	1.47	1.39
1	C	42	GLY	C-O	-5.82	1.14	1.23
1	C	158	TYR	CE2-CZ	-5.71	1.31	1.38
1	B	213	GLU	CD-OE1	5.56	1.31	1.25
1	B	203	TRP	CE3-CZ3	-5.50	1.29	1.38
1	B	256	TYR	CE1-CZ	-5.43	1.31	1.38
1	A	116	TYR	CG-CD1	5.38	1.46	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	158	TYR	CE1-CZ	5.35	1.45	1.38
1	D	244	GLU	CD-OE2	5.34	1.31	1.25
1	C	251	GLU	CD-OE1	5.33	1.31	1.25
1	B	203	TRP	CB-CG	5.33	1.59	1.50
1	B	194	TYR	CE1-CZ	-5.31	1.31	1.38
1	C	155	GLU	CD-OE2	-5.25	1.19	1.25
1	D	240	SER	CB-OG	5.20	1.49	1.42
1	C	27	GLU	CD-OE1	-5.14	1.20	1.25
1	C	264	GLY	C-O	-5.08	1.15	1.23
1	B	67[A]	SER	CB-OG	5.00	1.48	1.42
1	B	67[B]	SER	CB-OG	5.00	1.48	1.42

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	174	ARG	NE-CZ-NH2	-10.34	115.13	120.30
1	A	174	ARG	NE-CZ-NH1	10.33	125.46	120.30
1	A	55	ASP	CB-CG-OD2	-9.97	109.33	118.30
1	A	229	ARG	NE-CZ-NH1	9.76	125.18	120.30
1	B	205	ASP	CB-CG-OD1	8.37	125.83	118.30
1	B	229[A]	ARG	NE-CZ-NH2	-8.25	116.17	120.30
1	B	229[B]	ARG	NE-CZ-NH2	-8.25	116.17	120.30
1	C	229	ARG	NE-CZ-NH2	-7.97	116.31	120.30
1	C	205	ASP	CB-CG-OD1	7.92	125.43	118.30
1	B	91	ASP	CB-CG-OD1	7.53	125.08	118.30
1	A	186	ARG	NE-CZ-NH2	-7.20	116.70	120.30
1	C	37	ASP	CB-CG-OD1	7.16	124.74	118.30
1	C	229	ARG	NE-CZ-NH1	7.10	123.85	120.30
1	B	258	ASP	CB-CG-OD1	7.08	124.67	118.30
1	D	163[A]	ILE	CB-CA-C	-7.05	97.50	111.60
1	D	163[B]	ILE	CB-CA-C	-7.05	97.50	111.60
1	B	46	ARG	NE-CZ-NH1	7.00	123.80	120.30
1	C	174	ARG	NE-CZ-NH2	-6.80	116.90	120.30
1	D	199	ARG	NE-CZ-NH1	6.49	123.55	120.30
1	D	153	MET	CG-SD-CE	6.49	110.58	100.20
1	A	199	ARG	NE-CZ-NH1	6.43	123.51	120.30
1	B	163	ILE	N-CA-CB	6.39	125.51	110.80
1	A	254	ASP	CB-CG-OD2	6.29	123.97	118.30
1	C	225	ASP	CB-CG-OD1	-6.29	112.64	118.30
1	C	199	ARG	NE-CZ-NH1	6.25	123.42	120.30
1	B	164	GLU	OE1-CD-OE2	6.15	130.68	123.30
1	C	186	ARG	NE-CZ-NH1	6.14	123.37	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	175	LYS	CD-CE-NZ	-5.94	98.04	111.70
1	B	163	ILE	CB-CA-C	-5.92	99.77	111.60
1	D	174	ARG	NE-CZ-NH2	-5.86	117.37	120.30
1	B	167	GLU	OE1-CD-OE2	5.84	130.31	123.30
1	A	207	ASP	CB-CG-OD2	-5.83	113.05	118.30
1	B	229[A]	ARG	NE-CZ-NH1	5.81	123.21	120.30
1	B	229[B]	ARG	NE-CZ-NH1	5.81	123.21	120.30
1	B	257	ALA	N-CA-C	-5.80	95.34	111.00
1	D	130	ASP	CB-CG-OD1	5.70	123.43	118.30
1	C	91	ASP	CB-CG-OD2	-5.68	113.18	118.30
1	A	146	MET	CG-SD-CE	-5.67	91.13	100.20
1	C	73	LYS	CD-CE-NZ	-5.66	98.68	111.70
1	D	186[A]	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	D	186[B]	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	91	ASP	CB-CG-OD1	5.66	123.39	118.30
1	B	227	VAL	CG1-CB-CG2	-5.60	101.94	110.90
1	D	119	ASP	CB-CG-OD1	5.52	123.27	118.30
1	B	86	ILE	CB-CA-C	-5.51	100.57	111.60
1	B	186	ARG	CB-CA-C	-5.51	99.38	110.40
1	C	146	MET	CA-CB-CG	5.48	122.62	113.30
1	B	95	GLU	OE1-CD-OE2	-5.46	116.75	123.30
1	B	202	VAL	CG1-CB-CG2	-5.44	102.19	110.90
1	B	148	LEU	CB-CG-CD1	5.36	120.11	111.00
1	B	164	GLU	CG-CD-OE2	-5.32	107.66	118.30
1	C	103	HIS	N-CA-CB	-5.24	101.17	110.60
1	B	168	LEU	CB-CG-CD1	-5.19	102.18	111.00
1	C	254	ASP	CB-CG-OD2	5.17	122.96	118.30
1	D	229	ARG	NE-CZ-NH2	5.16	122.88	120.30
1	D	205	ASP	CB-CG-OD2	-5.14	113.67	118.30
1	D	165	ASP	CB-CG-OD1	5.13	122.92	118.30
1	C	154	ARG	CD-NE-CZ	5.06	130.69	123.60
1	D	205	ASP	CB-CG-OD1	5.03	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2131	0	2078	16	0
1	B	2112	0	2066	11	1
1	C	2138	0	2094	16	0
1	D	2132	0	2088	26	0
2	A	6	0	8	0	0
2	B	12	0	16	0	0
2	C	12	0	16	0	0
2	D	6	0	8	0	0
3	A	58	0	0	2	0
3	B	66	0	0	0	0
3	C	54	0	0	0	0
3	D	14	0	0	0	0
All	All	8741	0	8374	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:186[B]:ARG:NH1	1:D:186[B]:ARG:HB2	2.00	0.77
1:D:185:GLN:C	1:D:185:GLN:HE21	1.88	0.76
1:A:59:GLU:O	1:A:63[A]:THR:HG23	1.93	0.68
1:D:213[A]:GLU:HA	1:D:213[A]:GLU:OE2	1.94	0.67
1:D:186[B]:ARG:HH11	1:D:186[B]:ARG:HB2	1.59	0.67
1:C:157:LEU:CD1	1:C:209:ILE:HD11	2.27	0.65
1:D:163[A]:ILE:HD12	1:D:164:GLU:OE1	2.03	0.57
1:C:103:HIS:O	1:C:103:HIS:CD2	2.58	0.57
1:A:244:GLU:HG3	3:A:447:HOH:O	2.06	0.55
1:C:157:LEU:HD12	1:C:209:ILE:HD11	1.89	0.55
1:D:137:THR:HG22	1:D:142:THR:HG22	1.89	0.54
1:A:12:ILE:HD11	1:A:146:MET:HE1	1.90	0.53
1:D:125:PHE:CZ	1:D:209:ILE:HD12	2.44	0.52
1:A:104:ASN:ND2	1:A:215:GLN:HE22	2.08	0.51
1:C:9:ILE:HD13	1:C:86[B]:ILE:CD1	2.41	0.51
1:B:18:ILE:CD1	1:B:239:LEU:HD11	2.41	0.50
1:A:80:SER:HA	1:A:105:ALA:HA	1.93	0.50
1:C:103:HIS:HD2	1:C:103:HIS:O	1.92	0.49
1:B:52:ASN:ND2	1:B:138:SER:OG	2.44	0.49
1:D:185:GLN:O	1:D:185:GLN:NE2	2.34	0.49
1:C:49:GLU:HA	1:C:136:TYR:CE1	2.48	0.49
1:C:103:HIS:HE1	1:C:238:GLN:HB3	1.78	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:90:ALA:HA	1:D:97:ILE:HD12	1.94	0.48
1:A:163:ILE:HD11	3:A:455:HOH:O	2.12	0.48
1:D:10:HIS:CD2	1:D:15:GLY:HA2	2.48	0.47
1:D:213[A]:GLU:CA	1:D:213[A]:GLU:OE2	2.62	0.47
1:D:8:LEU:HB3	1:D:19:TRP:CE2	2.50	0.47
1:A:104:ASN:HD22	1:A:215:GLN:HE22	1.62	0.47
1:C:103:HIS:C	1:C:103:HIS:CD2	2.88	0.47
1:B:226:LYS:HE3	1:B:228:TYR:CZ	2.49	0.47
1:D:201:TYR:CE2	1:D:219:ILE:HD11	2.49	0.47
1:C:47:GLN:OE1	1:D:45:PRO:HB3	2.15	0.46
1:C:9:ILE:HD13	1:C:86[B]:ILE:HD11	1.98	0.46
1:A:24:PRO:HD3	1:B:167:GLU:HG2	1.98	0.45
1:A:90:ALA:HA	1:A:97:ILE:HD12	1.98	0.45
1:D:11:THR:HB	1:D:80:SER:HB3	1.99	0.45
1:C:9:ILE:CD1	1:C:86[B]:ILE:HD11	2.47	0.45
1:B:18:ILE:HD11	1:B:239:LEU:HD11	1.98	0.45
1:C:209:ILE:HG22	1:C:210:PHE:CD1	2.52	0.44
1:D:11:THR:CB	1:D:80:SER:HB3	2.48	0.44
1:A:163:ILE:N	1:A:163:ILE:CD1	2.81	0.44
1:C:12:ILE:HD11	1:C:146:MET:HE2	1.99	0.44
1:D:103:HIS:ND1	1:D:103:HIS:O	2.46	0.44
1:D:201:TYR:CD2	1:D:219:ILE:HD11	2.52	0.43
1:D:83:GLY:HA2	1:D:86:ILE:HD12	2.00	0.43
1:A:114:PRO:HG2	1:A:186:ARG:HG3	2.00	0.43
1:A:103:HIS:C	1:A:103:HIS:CD2	2.92	0.42
1:B:55:ASP:OD2	1:B:186:ARG:NH1	2.48	0.42
1:B:94:PRO:HG3	1:B:193:GLY:HA2	2.00	0.42
1:D:128:TRP:HB3	1:D:131:THR:HB	2.00	0.42
1:D:103:HIS:ND1	1:D:103:HIS:C	2.73	0.42
1:D:103:HIS:HD1	1:D:103:HIS:C	2.23	0.42
1:A:61:LEU:O	1:A:65:MET:HG2	2.19	0.42
1:C:157:LEU:HG	1:C:209:ILE:CD1	2.50	0.42
1:B:90:ALA:HA	1:B:97:ILE:HD12	2.01	0.42
1:B:86:ILE:HG22	1:B:87:ALA:N	2.33	0.42
1:D:90:ALA:HA	1:D:97:ILE:CD1	2.50	0.42
1:D:115:SER:OG	1:D:184:ALA:HA	2.20	0.41
1:D:92[B]:LYS:O	1:D:92[B]:LYS:HG3	2.21	0.41
1:A:153:MET:HG2	1:A:157:LEU:HD22	2.02	0.41
1:A:209:ILE:O	1:A:211:PRO:HD3	2.20	0.41
1:B:130:ASP:OD1	1:B:149:GLY:HA3	2.20	0.41
1:D:185:GLN:C	1:D:185:GLN:NE2	2.66	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:125:PHE:CD1	1:C:126:PRO:HD2	2.55	0.41
1:B:260:LEU:N	1:B:260:LEU:HD23	2.36	0.40
1:A:39:ALA:HB3	1:A:57:TYR:HA	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:137:THR:OG1	1:B:139:ASN:O[7_555]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/272 (97%)	253 (96%)	10 (4%)	0	100	100
1	B	261/272 (96%)	247 (95%)	14 (5%)	0	100	100
1	C	265/272 (97%)	254 (96%)	11 (4%)	0	100	100
1	D	263/272 (97%)	252 (96%)	11 (4%)	0	100	100
All	All	1052/1088 (97%)	1006 (96%)	46 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/236 (98%)	220 (95%)	11 (5%)	30	14
1	B	230/236 (98%)	218 (95%)	12 (5%)	27	12
1	C	231/236 (98%)	225 (97%)	6 (3%)	51	40
1	D	231/236 (98%)	218 (94%)	13 (6%)	25	11
All	All	923/944 (98%)	881 (95%)	42 (5%)	35	16

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63[A]	THR
1	A	63[B]	THR
1	A	70	GLN
1	A	86[A]	ILE
1	A	86[B]	ILE
1	A	103	HIS
1	A	123	GLU
1	A	148	LEU
1	A	163	ILE
1	A	192	GLU
1	A	209	ILE
1	B	25	LEU
1	B	55	ASP
1	B	98[A]	SER
1	B	98[B]	SER
1	B	103	HIS
1	B	139	ASN
1	B	140	ASN
1	B	142	THR
1	B	146	MET
1	B	148	LEU
1	B	163	ILE
1	B	260	LEU
1	C	55	ASP
1	C	91	ASP
1	C	103	HIS
1	C	148	LEU
1	C	180	GLN
1	C	209	ILE
1	D	25	LEU
1	D	55	ASP
1	D	62	LEU
1	D	73	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	92[A]	LYS
1	D	92[B]	LYS
1	D	103	HIS
1	D	146	MET
1	D	163[A]	ILE
1	D	163[B]	ILE
1	D	185	GLN
1	D	208	LYS
1	D	209	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	103	HIS
1	A	104	ASN
1	A	185	GLN
1	B	50	GLN
1	B	52	ASN
1	C	52	ASN
1	C	103	HIS
1	C	104	ASN
1	D	52	ASN
1	D	139	ASN
1	D	185	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	300	-	5,5,5	0.87	0	5,5,5	0.86	0
2	GOL	B	300	-	5,5,5	0.69	0	5,5,5	1.12	1 (20%)
2	GOL	B	301	-	5,5,5	0.94	0	5,5,5	0.94	0
2	GOL	C	300	-	5,5,5	1.43	0	5,5,5	1.41	1 (20%)
2	GOL	C	301	-	5,5,5	0.84	0	5,5,5	0.98	0
2	GOL	D	300	-	5,5,5	0.53	0	5,5,5	1.14	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	300	-	-	0/4/4/4	0/0/0/0
2	GOL	B	300	-	-	0/4/4/4	0/0/0/0
2	GOL	B	301	-	-	0/4/4/4	0/0/0/0
2	GOL	C	300	-	-	0/4/4/4	0/0/0/0
2	GOL	C	301	-	-	0/4/4/4	0/0/0/0
2	GOL	D	300	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	300	GOL	O1-C1-C2	2.14	120.85	110.07
2	B	300	GOL	O3-C3-C2	2.41	122.21	110.07
2	C	300	GOL	O1-C1-C2	2.92	124.77	110.07

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/272 (95%)	0.01	13 (5%) 30 40	20, 33, 51, 70	0
1	B	259/272 (95%)	0.00	11 (4%) 37 47	20, 29, 49, 86	0
1	C	265/272 (97%)	0.10	15 (5%) 24 34	22, 33, 56, 69	0
1	D	260/272 (95%)	0.58	36 (13%) 3 5	25, 48, 80, 108	0
All	All	1044/1088 (95%)	0.17	75 (7%) 16 25	20, 34, 65, 108	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	140	ASN	6.4
1	D	261	ALA	5.2
1	B	140	ASN	4.8
1	D	81	CYS	4.6
1	D	139	ASN	4.4
1	D	106	LEU	4.3
1	C	266	GLU	4.3
1	D	137	THR	4.3
1	D	142	THR	4.3
1	A	260	LEU	4.2
1	C	86[A]	ILE	4.1
1	D	138	SER	3.7
1	D	83	GLY	3.6
1	D	82	GLY	3.5
1	D	192	GLU	3.4
1	A	126	PRO	3.4
1	D	70	GLN	3.4
1	D	141	GLU	3.3
1	D	80	SER	3.3
1	D	110	THR	3.3
1	A	261	ALA	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	105	ALA	3.2
1	D	11	THR	3.1
1	D	126	PRO	3.0
1	B	260	LEU	3.0
1	D	77	VAL	3.0
1	D	84	LEU	3.0
1	D	9	ILE	2.9
1	D	86	ILE	2.9
1	C	84	LEU	2.9
1	D	217	TRP	2.8
1	A	125	PHE	2.8
1	D	111	GLU	2.8
1	D	69	PRO	2.7
1	C	9	ILE	2.7
1	D	191	GLU	2.7
1	A	70	GLN	2.7
1	C	82	GLY	2.7
1	C	137	THR	2.7
1	A	86[A]	ILE	2.6
1	C	81	CYS	2.6
1	B	258	ASP	2.6
1	C	77	VAL	2.5
1	D	125	PHE	2.5
1	B	137	THR	2.5
1	C	126	PRO	2.5
1	A	11	THR	2.4
1	C	136	TYR	2.4
1	D	136	TYR	2.4
1	B	11	THR	2.4
1	C	105	ALA	2.3
1	D	78	GLY	2.3
1	C	80	SER	2.3
1	C	78	GLY	2.3
1	D	116	TYR	2.3
1	C	139	ASN	2.3
1	B	101	VAL	2.3
1	B	106	LEU	2.3
1	A	127	ASP	2.3
1	C	11	THR	2.2
1	A	140	ASN	2.2
1	D	112	HIS	2.2
1	A	18	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	81	CYS	2.2
1	B	80	SER	2.1
1	D	103	HIS	2.1
1	D	143	ILE	2.1
1	D	102	PHE	2.1
1	A	83	GLY	2.1
1	B	102	PHE	2.1
1	A	81	CYS	2.1
1	D	71	GLY	2.1
1	D	12	ILE	2.0
1	B	105	ALA	2.0
1	A	84	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GOL	B	301	6/6	0.65	0.29	23.40	51,57,60,63	0
2	GOL	A	300	6/6	0.85	0.33	4.08	51,57,60,63	0
2	GOL	B	300	6/6	0.91	0.22	1.73	36,58,65,69	0
2	GOL	C	300	6/6	0.89	0.22	1.53	43,48,53,54	0
2	GOL	D	300	6/6	0.85	0.22	0.79	52,55,56,56	0
2	GOL	C	301	6/6	0.80	0.28	-	39,44,46,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.