



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 02:26 am GMT

PDB ID : 3TI3  
Title : Crystal structure of 2009 pandemic H1N1 neuraminidase complexed with laninamivir  
Authors : Vavricka, C.J.; Li, Q.; Wu, Y.; Qi, J.; Wang, M.; Liu, Y.; Gao, F.; Liu, J.; Feng, E.; He, J.; Wang, J.; Liu, H.; Jiang, H.; Gao, G.F.  
Deposited on : 2011-08-20  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="width: 92%; height: 10px; background-color: green;"></div> <div style="width: 7%; height: 10px; background-color: yellow;"></div> <div style="width: 3%; height: 10px; background-color: orange;"></div> <div style="width: 1%; height: 10px; background-color: red;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> <span>%</span> <span>92%</span> <span>7%</span> <span>•</span> </div> </div>
1	B	388	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="width: 89%; height: 10px; background-color: green;"></div> <div style="width: 11%; height: 10px; background-color: yellow;"></div> <div style="width: 1%; height: 10px; background-color: orange;"></div> <div style="width: 1%; height: 10px; background-color: red;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> <span>%</span> <span>89%</span> <span>11%</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	A	1	-	-	-	X
6	GOL	A	471	-	-	-	X
6	GOL	B	471	-	-	-	X
7	ACT	A	472	-	-	-	X
7	ACT	B	1	-	-	-	X
7	ACT	B	472	-	-	-	X

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	5	0
			3012	1893	517	581	21			
1	B	387	Total	C	N	O	S	0	5	0
			3008	1890	516	581	21			

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Ca	0	0
			2	2		
4	A	3	Total	Ca	0	0
			3	3		

- Molecule 5 is 5-(ACETYLAMINO)-2,6-ANHYDRO-4-CARBAMIMIDAMIDO-3,4,5-TRIDEOXY-7-O-METHYL-D-GLYCERO-D-GALACTO-NON-2-ENONIC ACID (three-letter code: LNV) (formula: C<sub>13</sub>H<sub>22</sub>N<sub>4</sub>O<sub>7</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			24	13	4	7		
5	B	1	Total	C	N	O	0	0
			24	13	4	7		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	601	Total	O	0	0
			601	601		
8	B	668	Total	O	0	0
			668	668		



- Molecule 1: Neuraminidase





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.26Å 137.09Å 118.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.60 – 1.80 34.60 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.7 (34.60-1.80) 99.9 (34.60-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.92 (at 1.79Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, $R_{free}$	0.194 , 0.213 0.194 , 0.211	Depositor DCC
$R_{free}$ test set	4455 reflections (5.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	10.9	Xtriage
Anisotropy	0.609	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7428	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.73 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.7757e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, LNV, NAG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/3106	0.47	0/4223
1	B	0.24	0/3105	0.46	0/4221
All	All	0.25	0/6211	0.46	0/8444

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3012	0	2854	17	0
1	B	3008	0	2852	33	0
2	A	28	0	25	0	0
3	A	14	0	13	1	0
3	B	14	0	13	0	0
4	A	3	0	0	0	0
4	B	2	0	0	0	0
5	A	24	0	20	0	0
5	B	24	0	20	1	0
6	A	12	0	16	3	0
6	B	6	0	8	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	4	0	3	0	0
7	B	8	0	6	1	0
8	A	601	0	0	4	0
8	B	668	0	0	4	0
All	All	7428	0	5830	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:GLU:HA	6:A:471:GOL:H11	1.61	0.82
1:B:129:CYS:H	6:B:471:GOL:H12	1.53	0.73
1:A:83:VAL:HG12	8:A:674:HOH:O	1.89	0.71
1:B:432:LYS:HB2	7:B:1:ACT:H1	1.82	0.61
1:A:416:ASP:HB2	8:A:1015:HOH:O	2.03	0.59
1:A:432:LYS:HB2	6:A:1:GOL:H12	1.84	0.59
1:B:122[A]:ILE:HG23	1:B:423:TRP:CD1	2.38	0.59
1:A:378:TRP:HB3	1:A:392:ILE:HB	1.88	0.55
1:B:270:ALA:HB1	1:B:273:TYR:HB2	1.89	0.55
1:B:378:TRP:HB3	1:B:392:ILE:HB	1.89	0.54
1:A:155:TYR:CE1	1:B:461:GLY:HA3	2.44	0.53
1:A:270:ALA:HB1	1:A:273:TYR:HB2	1.90	0.53
1:B:406:TYR:OH	5:B:801:LVN:CAA	2.58	0.51
1:B:146:ASN:ND2	8:B:857:HOH:O	2.21	0.51
1:B:84:LYS:HD2	1:B:235:GLY:HA3	1.93	0.50
1:A:122[B]:ILE:HD12	1:A:423:TRP:HB3	1.94	0.49
1:B:122[A]:ILE:HD12	1:B:123:SER:N	2.28	0.49
1:A:187:ILE:HG23	8:A:1291:HOH:O	2.13	0.49
1:B:325:ASN:O	1:B:348:GLY:HA2	2.13	0.48
3:A:505:NAG:H83	8:A:1181:HOH:O	2.14	0.48
1:A:122[A]:ILE:HD12	1:A:123:SER:N	2.29	0.48
1:B:259:LYS:HD2	8:B:644:HOH:O	2.12	0.47
1:A:325:ASN:O	1:A:348:GLY:HA2	2.14	0.47
1:B:122[A]:ILE:HG12	1:B:410:PHE:CE2	2.50	0.46
1:B:129:CYS:H	6:B:471:GOL:C1	2.27	0.46
1:B:102:LYS:HG3	1:B:444:ILE:HG22	1.97	0.46
1:B:122[A]:ILE:HG12	1:B:410:PHE:CZ	2.50	0.46
1:B:129:CYS:N	6:B:471:GOL:H12	2.26	0.44
1:B:378:TRP:CD1	1:B:391[B]:SER:OG	2.70	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122[A]:ILE:HG23	1:B:423:TRP:HD1	1.81	0.43
1:B:218:TRP:CD1	1:B:219:ARG:HG2	2.53	0.43
1:B:122[A]:ILE:CD1	1:B:129:CYS:HB3	2.49	0.43
1:B:228:SER:HB3	1:B:350:LYS:HE2	2.01	0.42
1:B:419:ARG:HA	1:B:420:PRO:HD3	1.86	0.42
1:B:121:PHE:CG	1:B:228:SER:HA	2.54	0.42
1:B:104:ASN:O	1:B:108:ILE:HG12	2.19	0.42
1:B:412:GLN:HG3	8:B:679:HOH:O	2.19	0.42
1:A:218:TRP:CD1	1:A:219:ARG:HG2	2.54	0.42
1:B:125:SER:HB2	1:B:126:PRO:HD2	2.01	0.42
1:B:412(B):PRO:HB3	1:B:415:LEU:O	2.20	0.42
1:B:84:LYS:HD2	1:B:235:GLY:CA	2.50	0.41
1:A:121:PHE:CG	1:A:228:SER:HA	2.55	0.41
1:B:406:TYR:HB2	1:B:425:GLU:OE1	2.21	0.41
1:A:173:PHE:CZ	1:B:101:SER:HA	2.56	0.41
1:A:412(B):PRO:HB3	1:A:415:LEU:O	2.21	0.41
1:A:429:GLY:O	6:A:1:GOL:H11	2.20	0.40
1:A:320:GLY:HA3	1:A:331:LYS:O	2.21	0.40
1:B:347:ASN:HB3	8:B:657:HOH:O	2.21	0.40
1:B:125:SER:HB2	1:B:126:PRO:CD	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/388 (100%)	375 (96%)	15 (4%)	0	100	100
1	B	390/388 (100%)	374 (96%)	16 (4%)	0	100	100
All	All	780/776 (100%)	749 (96%)	31 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	339/335 (101%)	332 (98%)	7 (2%)	59	46
1	B	339/335 (101%)	333 (98%)	6 (2%)	64	53
All	All	678/670 (101%)	665 (98%)	13 (2%)	68	50

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	122[A]	ILE
1	A	122[B]	ILE
1	A	215	ILE
1	A	230	CYS
1	A	296	HIS
1	A	331	LYS
1	A	394	LYS
1	B	122[A]	ILE
1	B	122[B]	ILE
1	B	215	ILE
1	B	230	CYS
1	B	296	HIS
1	B	309	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	188	ASN
1	A	372	ASN
1	A	401	ASN
1	A	435	ASN
1	B	372	ASN
1	B	401	ASN
1	B	435	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	501	1,2	14,14,15	0.55	0	15,19,21	0.79	1 (6%)
2	NAG	A	502	2	14,14,15	0.60	0	15,19,21	1.13	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	501	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	502	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NAG	C1-O5-C5	2.20	115.19	112.17
2	A	502	NAG	C3-C4-C5	2.37	114.40	110.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

Of 15 ligands modelled in this entry, 5 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
6	GOL	A	1	-	5,5,5	0.38	0	5,5,5	0.35	0
6	GOL	A	471	-	5,5,5	0.35	0	5,5,5	0.32	0
7	ACT	A	472	-	1,3,3	1.38	0	0,3,3	0.00	-
3	NAG	A	505	1	14,14,15	0.61	0	15,19,21	0.74	0
5	LNV	A	801	-	20,24,24	4.25	6 (30%)	16,33,33	1.41	1 (6%)
7	ACT	B	1	-	1,3,3	1.37	0	0,3,3	0.00	-
6	GOL	B	471	-	5,5,5	0.34	0	5,5,5	0.28	0
7	ACT	B	472	-	1,3,3	1.34	0	0,3,3	0.00	-
3	NAG	B	505	1	14,14,15	0.58	0	15,19,21	0.65	0
5	LNV	B	801	-	20,24,24	4.19	6 (30%)	16,33,33	1.41	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	1	-	-	0/4/4/4	0/0/0/0
6	GOL	A	471	-	-	0/4/4/4	0/0/0/0
7	ACT	A	472	-	-	0/0/0/0	0/0/0/0
3	NAG	A	505	1	-	0/6/23/26	0/1/1/1
5	LNV	A	801	-	-	0/20/40/40	0/1/1/1
7	ACT	B	1	-	-	0/0/0/0	0/0/0/0
6	GOL	B	471	-	-	0/4/4/4	0/0/0/0
7	ACT	B	472	-	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	505	1	-	0/6/23/26	0/1/1/1
5	LNV	B	801	-	-	0/20/40/40	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	801	LNV	OAF-CAE	-2.38	1.41	1.46
5	B	801	LNV	OAF-CAE	-2.33	1.42	1.46
5	A	801	LNV	OAY-CAK	-2.22	1.38	1.43
5	B	801	LNV	OAY-CAK	-2.22	1.38	1.43
5	B	801	LNV	CAC-CAB	2.32	1.53	1.50
5	B	801	LNV	CBD-NBG	2.42	1.43	1.34
5	A	801	LNV	CAC-CAB	2.45	1.53	1.50
5	A	801	LNV	CBD-NBG	2.47	1.43	1.34
5	B	801	LNV	CBA-NAZ	6.23	1.44	1.33
5	A	801	LNV	CBA-NAZ	6.29	1.44	1.33
5	B	801	LNV	CAB-CAA	16.61	1.53	1.32
5	A	801	LNV	CAB-CAA	16.87	1.53	1.32

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	801	LNV	CAE-CAD-NBG	3.51	117.18	111.00
5	B	801	LNV	CAE-CAD-NBG	3.55	117.24	111.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1	GOL	2	0
6	A	471	GOL	1	0
3	A	505	NAG	1	0
7	B	1	ACT	1	0
6	B	471	GOL	3	0
5	B	801	LNV	1	0



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	387/388 (99%)	-0.06	3 (0%) 86 84	5, 9, 17, 27	0
1	B	387/388 (99%)	-0.13	3 (0%) 86 84	4, 9, 17, 26	0
All	All	774/776 (99%)	-0.09	6 (0%) 86 84	4, 9, 17, 27	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	82	SER	3.3
1	B	82	SER	2.8
1	A	122[A]	ILE	2.6
1	A	83	VAL	2.3
1	B	83	VAL	2.1
1	B	122[A]	ILE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	501	14/15	0.79	0.30	-	20,26,32,34	0
2	NAG	A	502	14/15	0.73	0.35	-	27,34,42,49	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
7	ACT	B	472	4/4	0.88	0.24	8.34	18,23,27,29	0
6	GOL	A	471	6/6	0.80	0.24	4.92	14,23,28,32	0
6	GOL	B	471	6/6	0.90	0.24	4.67	8,23,32,43	0
7	ACT	A	472	4/4	0.90	0.17	4.54	18,22,26,29	0
6	GOL	A	1	6/6	0.91	0.27	4.35	1,18,26,27	0
7	ACT	B	1	4/4	0.93	0.18	2.42	2,15,21,27	0
5	LNV	A	801	24/24	0.95	0.09	0.29	5,6,9,12	0
5	LNV	B	801	24/24	0.96	0.08	-0.08	4,6,8,13	0
4	CA	B	601	1/1	1.00	0.06	-1.09	7,7,7,7	0
4	CA	A	601	1/1	1.00	0.06	-1.30	7,7,7,7	0
4	CA	A	602	1/1	0.99	0.08	-2.33	13,13,13,13	0
4	CA	B	602	1/1	0.99	0.04	-4.69	11,11,11,11	0
4	CA	A	603	1/1	1.00	0.04	-	13,13,13,13	1
3	NAG	B	505	14/15	0.73	0.31	-	24,35,43,44	0
3	NAG	A	505	14/15	0.65	0.29	-	26,33,41,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.