



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 2, 2017 – 06:04 AM EDT

PDB ID : 3TIH
Title : Crystal structure of unliganded HIV-1 clade C strain ZM109F.PB4 gp120 core
Authors : Kwon, Y.D.; Kwong, P.D.
Deposited on : unknown
Resolution : 4.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1088 (4.40-3.60)
Clashscore	112137	1187 (4.40-3.60)
Ramachandran outliers	110173	1139 (4.40-3.60)
Sidechain outliers	110143	1126 (4.40-3.60)
RSRZ outliers	101464	1099 (4.40-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	
1	B	345	
1	C	345	
1	D	345	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 10252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HIV-1 clade C ZM109F.PB4 gp120.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	S	0	0	0
			2563	1608	446	488	21			
1	B	326	Total	C	N	O	S	0	0	0
			2563	1608	446	488	21			
1	C	326	Total	C	N	O	S	0	0	0
			2563	1608	446	488	21			
1	D	326	Total	C	N	O	S	0	0	0
			2563	1608	446	488	21			

- Molecule 1: HIV-1 clade C ZM109F.PB4 gp120



● Molecule 1: HIV-1 clade C ZM109F.PB4 gp120



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.57Å 197.21Å 83.32Å 90.00° 91.25° 90.00°	Depositor
Resolution (Å)	46.70 – 4.00 48.41 – 2.80	Depositor EDS
% Data completeness (in resolution range)	84.6 (46.70-4.00) 33.7 (48.41-2.80)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.7.2_869	Depositor
R, R_{free}	0.283 , 0.312 0.279 , 0.297	Depositor DCC
R_{free} test set	577 reflections (4.88%)	DCC
Wilson B-factor (Å ²)	59.8	Xtriage
Anisotropy	0.726	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 141.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.081 for h,-k,-l	Xtriage
F_o, F_c correlation	0.82	EDS
Total number of atoms	10252	wwPDB-VP
Average B, all atoms (Å ²)	155.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/2613	0.40	0/3535
1	B	0.23	0/2613	0.43	1/3535 (0.0%)
1	C	0.25	0/2613	0.41	0/3535
1	D	0.23	0/2613	0.40	0/3535
All	All	0.24	0/10452	0.41	1/14140 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	65	VAL	CB-CA-C	-7.23	97.67	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2563	0	2519	55	0
1	B	2563	0	2519	54	0
1	C	2563	0	2519	51	0
1	D	2563	0	2519	52	0
All	All	10252	0	10076	209	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 209 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:ASN:HD21	1:A:213:ILE:HG21	1.52	0.75
1:B:67:ASN:HD21	1:B:213:ILE:HG21	1.52	0.75
1:C:67:ASN:HD21	1:C:213:ILE:HG21	1.52	0.73
1:C:331:CYS:HB2	1:C:416:LEU:HB2	1.69	0.72
1:D:67:ASN:HD21	1:D:213:ILE:HG21	1.52	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/345 (92%)	287 (91%)	27 (8%)	2 (1%)	28	70
1	B	316/345 (92%)	287 (91%)	27 (8%)	2 (1%)	28	70
1	C	316/345 (92%)	287 (91%)	27 (8%)	2 (1%)	28	70
1	D	316/345 (92%)	287 (91%)	27 (8%)	2 (1%)	28	70
All	All	1264/1380 (92%)	1148 (91%)	108 (8%)	8 (1%)	28	70

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	262	ASN
1	B	262	ASN
1	C	262	ASN
1	D	262	ASN
1	A	473	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/303 (96%)	287 (99%)	3 (1%)	80	90
1	B	290/303 (96%)	288 (99%)	2 (1%)	87	94
1	C	290/303 (96%)	288 (99%)	2 (1%)	87	94
1	D	290/303 (96%)	287 (99%)	3 (1%)	80	90
All	All	1160/1212 (96%)	1150 (99%)	10 (1%)	82	92

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	331	CYS
1	C	65	VAL
1	D	65	VAL
1	B	120	VAL
1	C	120	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	374	HIS
1	C	67	ASN
1	D	67	ASN
1	B	229	ASN
1	C	374	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/345 (94%)	-0.33	0 100 100	63, 149, 226, 310	0
1	B	326/345 (94%)	-0.31	2 (0%) 89 84	65, 150, 228, 308	0
1	C	326/345 (94%)	-0.32	2 (0%) 89 84	65, 150, 225, 313	0
1	D	326/345 (94%)	-0.34	1 (0%) 93 91	66, 151, 225, 326	0
All	All	1304/1380 (94%)	-0.33	5 (0%) 92 88	63, 150, 226, 326	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	121	LYS	4.0
1	D	348	LYS	3.3
1	B	332	LYS	2.9
1	C	58	ALA	2.1
1	B	124	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.