



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 21, 2017 – 10:53 AM EST

PDB ID : 1TND
Title : THE 2.2 ANGSTROMS CRYSTAL STRUCTURE OF TRANSDUCIN-
ALPHA COMPLEXED WITH GTP GAMMA S
Authors : Noel, J.P.; Hamm, H.E.; Sigler, P.B.
Deposited on : unknown
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	324	
1	B	324	
1	C	324	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8599 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSDUCIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2590	1641	437	493	19			
1	B	316	Total	C	N	O	S	0	0	0
			2539	1610	428	483	18			
1	C	316	Total	C	N	O	S	0	0	0
			2539	1610	428	483	18			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		

- Molecule 3 is CACODYLATE ION (three-letter code: CAC) (formula: C₂H₆AsO₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	As 1	0	0
3	A	1	Total 1	As 1	0	0
3	B	1	Total 1	As 1	0	0
3	B	1	Total 1	As 1	0	0
3	C	1	Total 1	As 1	0	0
3	C	1	Total 1	As 1	0	0

- Molecule 4 is 5'-GUANOSINE-DIPHOSPHATE-MONOTHIOPHOSPHATE (three-letter code: GSP) (formula: C₁₀H₁₆N₅O₁₃P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total 32	C 10	N 5	O 13	P 3	S 1	0	0
4	B	1	Total 32	C 10	N 5	O 13	P 3	S 1	0	0
4	C	1	Total 32	C 10	N 5	O 13	P 3	S 1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	305	Total O 305 305	0	0
5	B	225	Total O 225 225	0	0
5	C	296	Total O 296 296	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: TRANSDUCIN



• Molecule 1: TRANSDUCIN



• Molecule 1: TRANSDUCIN





4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.50 Å 108.30 Å 79.00 Å 90.00° 112.30° 90.00°	Depositor
Resolution (Å)	6.00 – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.20)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.190 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8599	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, GSP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	1/2635 (0.0%)	0.87	2/3552 (0.1%)
1	B	0.67	1/2584 (0.0%)	0.84	2/3484 (0.1%)
1	C	0.67	0/2584	0.82	1/3484 (0.0%)
All	All	0.68	2/7803 (0.0%)	0.85	5/10520 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	62	CYS	CB-SG	-7.92	1.68	1.82
1	A	250	CYS	CB-SG	-6.99	1.70	1.82

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	32	LEU	CA-CB-CG	7.14	131.72	115.30
1	A	32	LEU	CA-CB-CG	6.75	130.83	115.30
1	B	238	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	33	LEU	CA-CB-CG	5.27	127.43	115.30
1	B	245	LEU	CA-CB-CG	5.17	127.19	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2590	0	2572	84	0
1	B	2539	0	2518	116	0
1	C	2539	0	2520	92	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	1	0
3	C	2	0	0	2	0
4	A	32	0	12	1	0
4	B	32	0	12	3	0
4	C	32	0	12	0	0
5	A	305	0	0	11	0
5	B	225	0	0	16	0
5	C	296	0	0	14	0
All	All	8599	0	7646	290	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 290 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:MET:HB3	1:A:309:ARG:HH21	1.26	1.01
1:A:308:MET:HB3	1:A:309:ARG:NH2	1.85	0.92
1:B:252:HIS:HE1	1:B:254:TYR:HD2	1.14	0.91
1:C:84:MET:HA	1:C:89:ILE:HG22	1.57	0.87
1:B:168:GLN:O	1:B:172:ARG:HG2	1.75	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/324 (99%)	294 (92%)	22 (7%)	5 (2%)	11	8
1	B	314/324 (97%)	292 (93%)	18 (6%)	4 (1%)	14	11
1	C	314/324 (97%)	301 (96%)	12 (4%)	1 (0%)	44	49
All	All	949/972 (98%)	887 (94%)	52 (6%)	10 (1%)	17	13

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	311	ASP
1	A	347	CYS
1	A	258	THR
1	B	311	ASP
1	B	312	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/288 (100%)	262 (91%)	25 (9%)	12	12
1	B	281/288 (98%)	253 (90%)	28 (10%)	9	8
1	C	281/288 (98%)	257 (92%)	24 (8%)	12	12
All	All	849/864 (98%)	772 (91%)	77 (9%)	11	11

5 of 77 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	152	LEU
1	B	238	ARG
1	C	230	LEU
1	B	161	PRO
1	B	208	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	90	GLN
1	B	247	ASN
1	C	287	ASN
1	B	133	GLN
1	A	145	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are modelled with single atom and 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GSP	A	351	2	26,34,34	1.86	9 (34%)	24,54,54	2.59	5 (20%)
4	GSP	B	351	2	26,34,34	1.96	6 (23%)	24,54,54	2.78	7 (29%)
4	GSP	C	351	2	26,34,34	1.63	6 (23%)	24,54,54	2.64	7 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GSP	A	351	2	-	0/17/38/38	0/3/3/3
4	GSP	B	351	2	-	0/17/38/38	0/3/3/3
4	GSP	C	351	2	-	0/17/38/38	0/3/3/3

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	351	GSP	C2'-C1'	-3.11	1.48	1.53
4	A	351	GSP	PG-O2G	-2.92	1.45	1.55
4	A	351	GSP	C2'-C1'	-2.84	1.49	1.53
4	C	351	GSP	PG-O2G	-2.49	1.46	1.55
4	C	351	GSP	PG-O3G	-2.30	1.47	1.55

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	351	GSP	C5-C6-N1	-9.35	110.17	123.48
4	C	351	GSP	C5-C6-N1	-8.66	111.15	123.48
4	A	351	GSP	C5-C6-N1	-8.17	111.86	123.48
4	C	351	GSP	C1'-N9-C4	-3.48	120.62	126.64
4	C	351	GSP	C4'-O4'-C1'	-3.48	106.07	109.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	351	GSP	1	0
4	B	351	GSP	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.