



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:41 am GMT

PDB ID : 3TOD
Title : Crystal Structure of C-lobe of Bovine lactoferrin Complexed with 1-Butyl-1H
-Pyrazole-5-carboxylic acid at 1.38 Å Resolution
Authors : Shukla, P.K.; Gautam, L.; Sinha, M.; Kaur, P.; Sharma, S.; Singh, T.P.
Deposited on : 2011-09-05
Resolution : 1.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2133 (1.40-1.36)
Clashscore	112137	2266 (1.40-1.36)
Ramachandran outliers	110173	2215 (1.40-1.36)
Sidechain outliers	110143	2214 (1.40-1.36)
RSRZ outliers	101464	2141 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	<div> <div>16%</div> <div> <div></div> <div>86%</div> <div>14%</div> </div> </div>
2	B	6	<div> <div>100%</div> <div> <div></div> <div>67%</div> <div>33%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	XXB	A	777	-	-	-	X
8	GOL	A	700	-	-	X	X

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 3280 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	0	0
			2560	1593	448	499	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	565	LYS	ASN	SEE REMARK 999	UNP P24627
A	608	GLU	LYS	SEE REMARK 999	UNP P24627

- Molecule 2 is a protein called peptide, LEACAF from Lactotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	6	Total	C	N	O	S	0	0	0
			45	29	6	9	1			

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 4 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Fe	0	0
			1	1		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		

- Molecule 6 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	1	3		

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is ETHANOL (three-letter code: EOH) (formula: C_2H_6O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			3	2	1		
9	A	1	Total	C	O	0	0
			3	2	1		

- Molecule 10 is 1-BUTYL-1H-PYRAZOLE-5-CARBOXYLIC ACID (three-letter code: XXB) (formula: C₈H₁₂N₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	A	1	Total	C	N	O	0	0
			12	8	2	2		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	544	Total	O	0	0
			544	544		
11	B	5	Total	O	0	0
			5	5		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lactotransferrin



• Molecule 2: peptide, LEACAF from Lactotransferrin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.41Å 49.98Å 65.29Å 90.00° 107.14° 90.00°	Depositor
Resolution (Å)	62.39 – 1.38 19.93 – 1.38	Depositor EDS
% Data completeness (in resolution range)	97.0 (62.39-1.38) 97.0 (19.93-1.38)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.78 (at 1.38Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.200 , 0.236 0.195 , 0.231	Depositor DCC
R_{free} test set	3867 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	16.6	Xtriage
Anisotropy	0.330	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3280	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, NAG, EOH, CO3, XXB, FE, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.48	13/2608 (0.5%)	1.22	9/3533 (0.3%)
2	B	1.19	0/45	1.48	1/58 (1.7%)
All	All	1.48	13/2653 (0.5%)	1.22	10/3591 (0.3%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	601	SER	CB-OG	-9.33	1.30	1.42
1	A	647	CYS	CB-SG	-6.42	1.71	1.82
1	A	388	GLU	CD-OE1	-6.39	1.18	1.25
1	A	641	PHE	CG-CD1	6.12	1.48	1.38
1	A	675	CYS	CB-SG	-5.90	1.72	1.81
1	A	573	CYS	CB-SG	-5.87	1.72	1.81
1	A	534	ALA	CA-CB	-5.60	1.40	1.52
1	A	433	TYR	CE2-CZ	-5.48	1.31	1.38
1	A	398	TYR	CD2-CE2	-5.39	1.31	1.39
1	A	415	ARG	CB-CG	-5.24	1.38	1.52
1	A	516	VAL	CB-CG1	5.15	1.63	1.52
1	A	454	LYS	CD-CE	5.14	1.64	1.51
1	A	634	SER	CB-OG	-5.05	1.35	1.42

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	415	ARG	NE-CZ-NH2	-9.33	115.64	120.30
1	A	578	ARG	NE-CZ-NH2	-6.43	117.09	120.30
2	B	681	LEU	O-C-N	5.81	132.00	122.70
1	A	589	LEU	CB-CG-CD1	-5.38	101.85	111.00
1	A	379	ASP	CB-CG-OD2	-5.33	113.50	118.30
1	A	660	TYR	CD1-CE1-CZ	-5.31	115.02	119.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	641	PHE	CG-CD1-CE1	-5.28	115.00	120.80
1	A	428	ARG	NE-CZ-NH1	-5.22	117.69	120.30
1	A	602	ASP	CB-CG-OD2	-5.21	113.61	118.30
1	A	526	TYR	CD1-CE1-CZ	-5.16	115.16	119.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2560	0	2480	22	0
2	B	45	0	39	2	0
3	A	84	0	75	0	0
4	A	1	0	0	0	0
5	A	2	0	0	0	0
6	A	4	0	0	0	0
7	A	5	0	0	0	0
8	A	12	0	14	5	0
9	A	6	0	12	1	0
10	A	12	0	11	0	0
11	A	544	0	0	4	0
11	B	5	0	0	0	0
All	All	3280	0	2631	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:681:LEU:HD12	2:B:681:LEU:O	1.86	0.75
1:A:395:ASP:N	8:A:700:GOL:H32	2.04	0.71
1:A:395:ASP:H	8:A:700:GOL:H32	1.59	0.67
1:A:395:ASP:HA	1:A:595:HIS:CD2	2.30	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:623:LYS:HB3	1:A:623:LYS:HZ2	1.68	0.58
1:A:617:LEU:HD23	1:A:623:LYS:HE3	1.86	0.57
1:A:395:ASP:H	8:A:700:GOL:C3	2.18	0.56
1:A:497:PRO:O	1:A:514:LYS:NZ	2.34	0.54
1:A:565:LYS:HG3	1:A:567:GLU:H	1.71	0.54
1:A:421:SER:HB2	11:A:791:HOH:O	2.07	0.54
8:A:700:GOL:H12	11:A:20:HOH:O	2.08	0.54
2:B:681:LEU:O	2:B:682:GLU:HB3	2.11	0.50
1:A:653:GLY:O	1:A:654:ARG:C	2.51	0.48
1:A:671:ASN:HA	1:A:674:LYS:HE2	1.96	0.47
1:A:553:ASN:HD21	1:A:565:LYS:NZ	2.13	0.46
1:A:626:PRO:HA	1:A:630:CYS:SG	2.56	0.46
1:A:566:ARG:HG2	1:A:581:VAL:CG2	2.46	0.45
1:A:632:PHE:O	1:A:633:LYS:HD3	2.17	0.45
9:A:701:EOH:H21	11:A:120:HOH:O	2.17	0.44
1:A:520:LYS:HD2	1:A:520:LYS:HA	1.79	0.44
1:A:566:ARG:HG2	1:A:581:VAL:HG21	1.99	0.43
1:A:669:ILE:CG2	1:A:673:LYS:HE2	2.48	0.43
1:A:567:GLU:O	1:A:567:GLU:HG3	2.18	0.42
1:A:514:LYS:HE3	1:A:514:LYS:HB3	1.83	0.42
8:A:700:GOL:C1	11:A:20:HOH:O	2.68	0.42
1:A:403:GLY:HA3	1:A:657:TYR:CG	2.55	0.42
1:A:455:LYS:HB3	1:A:504:LEU:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/335 (99%)	324 (97%)	9 (3%)	0	100	100
2	B	4/6 (67%)	3 (75%)	0	1 (25%)	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	337/341 (99%)	327 (97%)	9 (3%)	1 (0%)	44 17

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	682	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	278/278 (100%)	277 (100%)	1 (0%)	93 81
2	B	4/4 (100%)	4 (100%)	0	100 100
All	All	282/282 (100%)	281 (100%)	1 (0%)	93 81

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	515	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	360	GLN
1	A	489	GLN
1	A	551	ASN
1	A	585	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	A	687	1,3	14,14,15	1.15	2 (14%)	15,19,21	1.56	2 (13%)
3	NAG	A	688	3	14,14,15	0.82	0	15,19,21	1.82	3 (20%)
3	NAG	A	689	1,3	14,14,15	0.84	0	15,19,21	2.13	4 (26%)
3	NAG	A	690	3	14,14,15	0.83	1 (7%)	15,19,21	1.39	2 (13%)
3	NAG	A	692	1,3	14,14,15	0.90	0	15,19,21	1.86	6 (40%)
3	NAG	A	693	3	14,14,15	0.70	0	15,19,21	2.07	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	687	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	688	3	-	0/6/23/26	0/1/1/1
3	NAG	A	689	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	690	3	-	0/6/23/26	0/1/1/1
3	NAG	A	692	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	693	3	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	687	NAG	O5-C1	-2.59	1.39	1.43
3	A	690	NAG	O5-C1	-2.06	1.40	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	687	NAG	O7-C7	2.21	1.28	1.23

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	693	NAG	C3-C4-C5	-5.71	100.15	110.22
3	A	689	NAG	C1-C2-N2	-4.95	102.03	110.49
3	A	687	NAG	O5-C1-C2	-3.98	105.93	111.47
3	A	688	NAG	C1-O5-C5	-3.80	106.92	112.17
3	A	689	NAG	O5-C1-C2	-3.22	106.99	111.47
3	A	689	NAG	C4-C3-C2	-2.92	106.73	111.02
3	A	692	NAG	O5-C1-C2	-2.76	107.64	111.47
3	A	690	NAG	O7-C7-C8	-2.51	117.48	122.06
3	A	693	NAG	O6-C6-C5	-2.48	103.01	111.34
3	A	692	NAG	O3-C3-C4	-2.37	105.20	110.36
3	A	687	NAG	C6-C5-C4	-2.22	107.80	113.00
3	A	692	NAG	C2-N2-C7	-2.15	119.81	122.94
3	A	692	NAG	O4-C4-C5	-2.01	104.23	109.28
3	A	689	NAG	O7-C7-N2	2.04	125.85	121.92
3	A	693	NAG	O4-C4-C5	2.66	115.98	109.28
3	A	692	NAG	C4-C3-C2	2.69	114.97	111.02
3	A	690	NAG	C1-O5-C5	2.88	116.13	112.17
3	A	688	NAG	C3-C4-C5	3.26	115.97	110.22
3	A	692	NAG	C1-O5-C5	3.52	117.01	112.17
3	A	688	NAG	C4-C3-C2	3.76	116.53	111.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	CO3	A	697	4	0,3,3	0.00	-	0,3,3	0.00	-
7	SO4	A	698	-	4,4,4	0.31	0	6,6,6	0.48	0
8	GOL	A	699	-	5,5,5	0.30	0	5,5,5	0.36	0
8	GOL	A	700	-	5,5,5	2.13	2 (40%)	5,5,5	1.49	1 (20%)
9	EOH	A	701	-	2,2,2	0.52	0	1,1,1	0.48	0
9	EOH	A	703	-	2,2,2	0.46	0	1,1,1	0.17	0
10	XXB	A	777	-	8,12,12	0.55	0	8,15,15	1.99	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	CO3	A	697	4	-	0/0/0/0	0/0/0/0
7	SO4	A	698	-	-	0/0/0/0	0/0/0/0
8	GOL	A	699	-	-	0/4/4/4	0/0/0/0
8	GOL	A	700	-	-	0/4/4/4	0/0/0/0
9	EOH	A	701	-	-	0/0/0/0	0/0/0/0
9	EOH	A	703	-	-	0/0/0/0	0/0/0/0
10	XXB	A	777	-	-	0/4/8/8	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	700	GOL	O3-C3	-3.85	1.26	1.42
8	A	700	GOL	O1-C1	-2.07	1.33	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	700	GOL	C3-C2-C1	-2.69	100.82	111.52
10	A	777	XXB	C5-N1-N2	-2.47	114.74	120.61
10	A	777	XXB	C4-C3-C2	4.33	108.30	104.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	700	GOL	5	0
9	A	701	EOH	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	335/335 (100%)	1.27	52 (15%) 2 3	9, 16, 33, 58	1 (0%)
2	B	6/6 (100%)	9.50	6 (100%) 0 0	33, 38, 65, 67	0
All	All	341/341 (100%)	1.41	58 (17%) 2 2	9, 16, 35, 67	1 (0%)

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	681	LEU	21.4
2	B	682	GLU	12.4
1	A	420	HIS	8.9
1	A	342	TYR	8.6
1	A	419	LYS	8.0
1	A	418	SER	6.5
1	A	422	SER	6.5
2	B	686	PHE	6.1
2	B	683	ALA	6.0
1	A	511	GLY	5.8
2	B	684	CYS	5.6
2	B	685	ALA	5.5
1	A	421	SER	5.4
1	A	627	ASP	4.7
1	A	429	PRO	4.7
1	A	510	GLN	4.6
1	A	621	ASN	4.3
1	A	635	GLU	4.2
1	A	676	SER	4.2
1	A	626	PRO	4.1
1	A	423	LEU	4.1
1	A	509	ASP	3.7
1	A	559	ASP	3.6
1	A	417	SER	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	373	THR	3.5
1	A	438	VAL	3.4
1	A	424	ASP	3.0
1	A	439	VAL	3.0
1	A	581	VAL	3.0
1	A	648	LEU	2.9
1	A	399	ILE	2.9
1	A	661	LEU	2.9
1	A	664	GLU	2.8
1	A	631	LEU	2.7
1	A	623	LYS	2.7
1	A	416	LYS	2.7
1	A	572	LEU	2.6
1	A	609	GLN	2.6
1	A	557	THR	2.6
1	A	344	ARG	2.6
1	A	512	LEU	2.5
1	A	363	GLN	2.5
1	A	427	LEU	2.5
1	A	640	LEU	2.5
1	A	663	THR	2.4
1	A	632	PHE	2.3
1	A	430	THR	2.3
1	A	548	VAL	2.3
1	A	473	LEU	2.2
1	A	667	THR	2.2
1	A	475	VAL	2.2
1	A	675	CYS	2.2
1	A	636	THR	2.1
1	A	481	CYS	2.1
1	A	597	VAL	2.1
1	A	530	PHE	2.0
1	A	669	ILE	2.0
1	A	651	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	687	14/15	0.79	0.17	0.29	22,27,35,39	0
3	NAG	A	689	14/15	0.79	0.13	-0.10	23,29,38,40	0
3	NAG	A	693	14/15	0.85	0.17	-0.34	27,35,40,42	0
3	NAG	A	692	14/15	0.90	0.08	-0.71	17,23,26,28	0
3	NAG	A	690	14/15	0.88	0.20	-	35,40,43,51	0
3	NAG	A	688	14/15	0.69	0.42	-	51,59,62,64	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
8	GOL	A	700	6/6	0.91	0.29	9.89	15,28,31,36	0
10	XXB	A	777	12/12	0.49	0.29	4.92	32,35,40,41	12
6	CO3	A	697	4/4	0.95	0.12	1.24	7,9,10,10	0
9	EOH	A	703	3/3	0.81	0.11	0.42	31,31,32,34	0
9	EOH	A	701	3/3	0.93	0.14	0.02	24,24,30,31	0
5	ZN	A	695	1/1	0.99	0.08	-1.39	14,14,14,14	0
4	FE	A	694	1/1	1.00	0.09	-1.73	9,9,9,9	0
7	SO4	A	698	5/5	0.93	0.21	-	32,34,37,42	0
8	GOL	A	699	6/6	0.34	0.27	-	43,50,51,51	0
5	ZN	A	696	1/1	0.98	0.07	-	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.