



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 10:54 pm GMT

PDB ID : 1TPA  
Title : THE GEOMETRY OF THE REACTIVE SITE AND OF THE PEPTIDE GROUPS IN TRYPSIN, TRYPSINOGEN AND ITS COMPLEXES WITH INHIBITORS  
Authors : Huber, R.; Bode, W.; Deisenhofer, J.  
Deposited on : 1982-09-27  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	223	<div> <div>2%</div> <div>69%</div> <div>28%</div> <div>.</div> </div>
2	I	58	<div> <div>69%</div> <div>24%</div> <div>.</div> <div>.</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2242 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANHYDRO-TRYPSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	223	Total	C	N	O	S	36	0	0
			1628	1012	279	323	14			

- Molecule 2 is a protein called BOVINE PANCREATIC TRYPSIN INHIBITOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	58	Total	C	N	O	S	35	0	0
			454	284	84	79	7			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	1	Total	Ca	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	123	Total	O	0	0
			123	123		
4	I	36	Total	O	0	0
			36	36		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ANHYDRO-TRYPSIN



- Molecule 2: BOVINE PANCREATIC TRYPSIN INHIBITOR



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.50Å 84.40Å 122.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.80 – 1.90 30.72 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.80-1.90) 68.7 (30.72-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	unknown	Depositor
R, $R_{free}$	0.175 , (Not available) 0.174 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	16.6	Xtriage
Anisotropy	0.343	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 69.7	EDS
L-test for twinning <sup>1</sup>	$\langle  L  \rangle = 0.38$ , $\langle L^2 \rangle = 0.21$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2242	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	E	1.22	7/1659 (0.4%)	1.41	13/2249 (0.6%)
2	I	1.15	0/465	1.58	6/622 (1.0%)
All	All	1.20	7/2124 (0.3%)	1.45	19/2871 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	1	19
2	I	0	7
All	All	1	26

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	195	SER	CA-CB	-12.54	1.34	1.52
1	E	237	TRP	NE1-CE2	-7.53	1.27	1.37
1	E	215	TRP	NE1-CE2	-7.53	1.27	1.37
1	E	51	TRP	NE1-CE2	-6.58	1.28	1.37
1	E	186	GLU	CD-OE2	5.94	1.32	1.25
1	E	141	TRP	NE1-CE2	-5.71	1.30	1.37
1	E	29	TYR	CZ-OH	5.23	1.46	1.37

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	17	ARG	NE-CZ-NH1	10.46	125.53	120.30
2	I	17	ARG	NE-CZ-NH2	-9.82	115.39	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	195	SER	N-CA-CB	9.59	124.89	110.50
2	I	53	ARG	NE-CZ-NH2	-8.48	116.06	120.30
2	I	53	ARG	NE-CZ-NH1	8.26	124.43	120.30
1	E	151	TYR	CB-CG-CD1	-7.73	116.36	121.00
2	I	53	ARG	CD-NE-CZ	7.39	133.94	123.60
1	E	165	ASP	CB-CG-OD2	-6.38	112.56	118.30
1	E	156	LYS	CA-CB-CG	-5.79	100.67	113.40
1	E	154	VAL	CA-CB-CG1	-5.59	102.51	110.90
1	E	66	ARG	NE-CZ-NH2	-5.52	117.54	120.30
1	E	204	LYS	CA-CB-CG	-5.47	101.37	113.40
1	E	79	ASN	N-CA-CB	5.45	120.41	110.60
1	E	195	SER	CB-CA-C	5.33	120.22	110.10
2	I	3	ASP	N-CA-CB	5.33	120.19	110.60
1	E	151	TYR	CB-CG-CD2	5.14	124.08	121.00
1	E	97	ASN	CA-CB-CG	-5.04	102.30	113.40
1	E	130	SER	N-CA-CB	-5.03	102.95	110.50
1	E	17	VAL	CB-CA-C	-5.01	101.88	111.40

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	E	195	SER	CA

All (26) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	112	ALA	Mainchain
1	E	125	THR	Mainchain
1	E	132	ALA	Mainchain
1	E	147	SER	Mainchain
1	E	165	ASP	Sidechain
1	E	175	GLN	Sidechain
1	E	216	GLY	Mainchain
1	E	223	ASN	Sidechain
1	E	233	ASN	Sidechain,Mainchain
1	E	240	GLN	Sidechain
1	E	27	VAL	Mainchain
1	E	30	GLN	Sidechain
1	E	40	HIS	Mainchain
1	E	60	LYS	Mainchain
1	E	80	GLU	Sidechain
1	E	81	GLN	Sidechain

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	E	85	ALA	Mainchain
1	E	97	ASN	Sidechain
2	I	2	PRO	Mainchain
2	I	24	ASN	Sidechain
2	I	3	ASP	Sidechain
2	I	35	TYR	Mainchain
2	I	37	GLY	Mainchain
2	I	56	GLY	Mainchain
2	I	7	GLU	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1628	0	1585	22	2
2	I	454	0	438	6	2
3	E	1	0	0	0	0
4	E	123	0	0	0	0
4	I	36	0	0	1	0
All	All	2242	0	2023	28	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:158:LEU:HD11	1:E:188:LYS:HB3	1.81	0.61
1:E:64:GLN:HE21	1:E:66:ARG:HE	1.49	0.61
2:I:31:GLN:HB3	4:I:588:HOH:O	2.04	0.55
1:E:48:ASN:HD22	1:E:50:GLN:H	1.58	0.52
2:I:27:ALA:HB1	2:I:29:LEU:HD22	1.90	0.52
1:E:64:GLN:HE22	1:E:66:ARG:HH21	1.59	0.51
1:E:199:VAL:HG21	1:E:228:TYR:CD1	2.48	0.48
1:E:136:CYS:O	1:E:159:LYS:HA	2.14	0.47
1:E:137:LEU:HD11	1:E:157:CYS:HB3	1.96	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:17:VAL:O	1:E:188:LYS:HA	2.16	0.45
1:E:90:VAL:HA	1:E:104:MET:HB2	1.99	0.45
2:I:23:TYR:CE1	2:I:58:ALA:HB3	2.51	0.45
2:I:27:ALA:CB	2:I:29:LEU:HD22	2.46	0.45
1:E:212:ILE:HB	1:E:229:THR:HB	2.00	0.43
1:E:211:GLY:HA2	1:E:229:THR:O	2.19	0.43
1:E:142:GLY:HA2	1:E:193:GLY:HA3	2.01	0.42
1:E:56:ALA:HB1	1:E:90:VAL:HG13	2.00	0.42
2:I:13:PRO:HG3	2:I:39:ARG:HH11	1.85	0.42
1:E:124:PRO:HA	1:E:204:LYS:HD3	2.02	0.41
1:E:49:SER:HA	1:E:114:LEU:HD11	2.02	0.41
1:E:198:PRO:HB2	1:E:200:VAL:HG23	2.01	0.41
1:E:50:GLN:HG2	1:E:111:ALA:HA	2.03	0.41
1:E:48:ASN:HD22	1:E:48:ASN:C	2.24	0.41
1:E:55:ALA:O	1:E:58:CYS:HB2	2.22	0.40
1:E:165:ASP:O	1:E:169:LYS:HG3	2.21	0.40
1:E:129:ALA:HB1	1:E:162:ILE:HD12	2.04	0.40
1:E:49:SER:HB3	1:E:114:LEU:HD13	2.03	0.40
2:I:17:ARG:O	2:I:17:ARG:HD2	2.22	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:109:LYS:CG	2:I:2:PRO:CG[2_575]	1.88	0.32
1:E:109:LYS:NZ	2:I:54:THR:O[2_575]	2.10	0.10

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	E	221/223 (99%)	214 (97%)	7 (3%)	0	100 100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	56/58 (97%)	54 (96%)	1 (2%)	1 (2%)	10	2
All	All	277/281 (99%)	268 (97%)	8 (3%)	1 (0%)	38	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	I	3	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	183/184 (100%)	174 (95%)	9 (5%)	29	17
2	I	46/46 (100%)	42 (91%)	4 (9%)	12	4
All	All	229/230 (100%)	216 (94%)	13 (6%)	24	13

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	48	ASN
1	E	75	VAL
1	E	86	SER
1	E	104	MET
1	E	113	SER
1	E	127	SER
1	E	135	GLN
1	E	165	ASP
1	E	242	ILE
2	I	3	ASP
2	I	17	ARG
2	I	29	LEU
2	I	52	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	30	GLN
1	E	48	ASN
1	E	64	GLN
1	E	100	ASN
1	E	210	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	E	223/223 (100%)	-0.26	4 (1%) 69 72	8, 19, 31, 38	21 (9%)
2	I	56/58 (96%)	-0.42	0 100 100	9, 17, 28, 39	7 (12%)
All	All	279/281 (99%)	-0.29	4 (1%) 75 78	8, 19, 31, 39	28 (10%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	115	ASN	2.5
1	E	212	ILE	2.5
1	E	125	THR	2.4
1	E	147	SER	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	E	462	1/1	0.98	0.08	0.46	32,32,32,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.