



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 2, 2017 – 12:58 AM EDT

PDB ID : 3TR5
Title : Structure of a peptide chain release factor 3 (prfC) from Coxiella burnetii
Authors : Cheung, J.; Franklin, M.C.; Rudolph, M.; Cassidy, M.; Gary, E.; Burshteyn, F.; Love, J.
Deposited on : unknown
Resolution : 2.11 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4988 (2.14-2.10)
Clashscore	112137	5557 (2.14-2.10)
Ramachandran outliers	110173	5504 (2.14-2.10)
Sidechain outliers	110143	5505 (2.14-2.10)
RSRZ outliers	101464	5021 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	528	
1	B	528	
1	C	528	
1	D	528	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 17668 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptide chain release factor 3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	518	Total	C	N	O	S	Se	0	0	0
			4139	2631	720	772	4	12			
1	B	508	Total	C	N	O	S	Se	0	0	0
			4064	2588	704	756	4	12			
1	C	507	Total	C	N	O	S	Se	0	1	0
			4065	2588	704	757	4	12			
1	D	508	Total	C	N	O	S	Se	0	0	0
			4058	2584	701	757	4	12			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP Q83DC7
A	-1	ASN	-	EXPRESSION TAG	UNP Q83DC7
A	0	ALA	-	EXPRESSION TAG	UNP Q83DC7
B	-2	SER	-	EXPRESSION TAG	UNP Q83DC7
B	-1	ASN	-	EXPRESSION TAG	UNP Q83DC7
B	0	ALA	-	EXPRESSION TAG	UNP Q83DC7
C	-2	SER	-	EXPRESSION TAG	UNP Q83DC7
C	-1	ASN	-	EXPRESSION TAG	UNP Q83DC7
C	0	ALA	-	EXPRESSION TAG	UNP Q83DC7
D	-2	SER	-	EXPRESSION TAG	UNP Q83DC7
D	-1	ASN	-	EXPRESSION TAG	UNP Q83DC7
D	0	ALA	-	EXPRESSION TAG	UNP Q83DC7

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		

Continued on next page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Ca 1	0	0
2	C	1	Total 1	Ca 1	0	0

-
- The image displays the chemical structure of Guanosine Diphosphate (GDP). It consists of a guanine base (a purine derivative) linked to a ribose sugar, which is in turn linked to two phosphate groups. The guanine base is shown with its characteristic fused ring system, including the amino group at the 2-position. The ribose sugar is a five-membered ring with hydroxyl groups at the 2' and 3' positions. The two phosphate groups are connected by a pyrophosphate linkage, with the second phosphate group being the terminal one. The structure is labeled with atom names and numbers, such as N1, C2, N3, C4, N7, C8 for the guanine base, and C1', C2', C3', C4', C5' for the ribose sugar. The phosphate groups are labeled with P, O, and A, and B, indicating the phosphorus and oxygen atoms and the adenosine and butyrate groups respectively.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	363	Total O 363 363	0	0
4	B	310	Total O 310 310	0	0
4	C	250	Total O 250 250	0	0




Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	303	Total 303	O 303	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peptide chain release factor 3




- Molecule 1: Peptide chain release factor 3





- Molecule 1: Peptide chain release factor 3





• Molecule 1: Peptide chain release factor 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.25Å 141.43Å 161.87Å 90.00° 95.62° 90.00°	Depositor
Resolution (Å)	41.97 – 2.11 41.96 – 2.11	Depositor EDS
% Data completeness (in resolution range)	97.6 (41.97-2.11) 97.5 (41.96-2.11)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.7 _650	Depositor
R, R_{free}	0.215 , 0.256 0.206 , 0.248	Depositor DCC
R_{free} test set	7188 reflections (5.01%)	DCC
Wilson B-factor (Å ²)	33.9	Xtriage
Anisotropy	0.206	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 32.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	17668	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/4210	0.44	0/5660
1	B	0.26	0/4133	0.42	0/5557
1	C	0.24	0/4135	0.41	0/5561
1	D	0.26	0/4128	0.42	0/5552
All	All	0.26	0/16606	0.42	0/22330

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4139	0	4140	62	0
1	B	4064	0	4072	72	0
1	C	4065	0	4066	91	0
1	D	4058	0	4057	61	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	28	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	28	0	12	1	0
3	C	28	0	12	2	0
3	D	28	0	12	0	0
4	A	363	0	0	14	0
4	B	310	0	0	12	0
4	C	250	0	0	14	0
4	D	303	0	0	9	0
All	All	17668	0	16383	284	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 284 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:388:ASN:HD21	1:D:525:HIS:HD2	1.10	0.99
1:C:112:ALA:HB1	1:C:145:THR:HG23	1.58	0.84
1:B:149:ILE:H	1:B:149:ILE:HD12	1.46	0.80
1:C:145:THR:HG22	1:C:146:ARG:O	1.83	0.79
1:A:51:HIS:N	4:A:1025:HOH:O	2.19	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	512/528 (97%)	501 (98%)	10 (2%)	1 (0%)	51	51
1	B	500/528 (95%)	477 (95%)	21 (4%)	2 (0%)	38	35
1	C	500/528 (95%)	478 (96%)	22 (4%)	0	100	100
1	D	500/528 (95%)	482 (96%)	18 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	2012/2112 (95%)	1938 (96%)	71 (4%)	3 (0%)	55 56

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	45	SER
1	A	340	ASN
1	B	65	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	444/438 (101%)	438 (99%)	6 (1%)	71 77
1	B	437/438 (100%)	435 (100%)	2 (0%)	91 94
1	C	437/438 (100%)	433 (99%)	4 (1%)	82 86
1	D	436/438 (100%)	430 (99%)	6 (1%)	71 77
All	All	1754/1752 (100%)	1736 (99%)	18 (1%)	80 84

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	62	GLN
1	C	101	THR
1	D	54	SER
1	B	310	ARG
1	B	454	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	257	ASN
1	D	525	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	276	GLN
1	C	281	ASN
1	D	388	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GDP	A	527	-	25,30,30	1.16	2 (8%)	26,47,47	2.01	6 (23%)
3	GDP	B	527	-	25,30,30	1.16	2 (8%)	26,47,47	1.99	7 (26%)
3	GDP	C	527	-	25,30,30	1.13	2 (8%)	26,47,47	2.03	6 (23%)
3	GDP	D	527	-	25,30,30	1.19	2 (8%)	26,47,47	2.11	6 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	527	-	-	0/12/32/32	0/3/3/3
3	GDP	B	527	-	-	0/12/32/32	0/3/3/3
3	GDP	C	527	-	-	0/12/32/32	0/3/3/3
3	GDP	D	527	-	-	0/12/32/32	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	527	GDP	C5-C4	2.95	1.47	1.40
3	A	527	GDP	C5-C4	3.00	1.47	1.40
3	B	527	GDP	C5-C4	3.07	1.47	1.40
3	D	527	GDP	C5-C4	3.11	1.47	1.40
3	C	527	GDP	C6-C5	3.72	1.48	1.41

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	527	GDP	C5-C6-N1	-4.19	117.52	123.48
3	A	527	GDP	C6-C5-C4	-4.16	116.70	120.84
3	C	527	GDP	C6-C5-C4	-4.15	116.72	120.84
3	D	527	GDP	C6-C5-C4	-4.03	116.84	120.84
3	C	527	GDP	C5-C6-N1	-4.00	117.79	123.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	527	GDP	1	0
3	C	527	GDP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	506/528 (95%)	-0.35	2 (0%) 92 93	15, 29, 56, 82	0
1	B	496/528 (93%)	-0.21	11 (2%) 62 67	17, 37, 72, 100	0
1	C	495/528 (93%)	0.03	14 (2%) 53 60	19, 45, 82, 122	0
1	D	496/528 (93%)	-0.23	10 (2%) 65 70	18, 34, 62, 90	0
All	All	1993/2112 (94%)	-0.19	37 (1%) 67 71	15, 36, 72, 122	0

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	63	ARG	4.9
1	C	476	ALA	4.0
1	D	62	GLN	3.5
1	C	3	VAL	3.3
1	D	193	GLY	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GDP	C	527	28/28	0.98	0.10	-0.02	27,41,43,44	0
3	GDP	D	527	28/28	0.97	0.10	-0.03	18,37,42,44	0
3	GDP	B	527	28/28	0.98	0.10	-0.07	27,47,50,54	0
3	GDP	A	527	28/28	0.99	0.09	-0.24	19,25,31,36	0
2	CA	B	526	1/1	0.12	0.21	-	93,93,93,93	0
2	CA	C	526	1/1	0.79	0.14	-	81,81,81,81	0
2	CA	D	526	1/1	0.77	0.12	-	65,65,65,65	0
2	CA	A	526	1/1	0.99	0.06	-	47,47,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.