



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 17, 2017 – 06:29 AM EDT

PDB ID : 5TRY  
Title : Structure of Mycobacterium tuberculosis proteasome in complex with N,C-capped dipeptide PKS2206  
Authors : Hsu, H.-C.; Fan, H.; Singh, P.K.; Wang, R.; Sukenick, G.; Nathan, C.; Lin, G.; Li, H.  
Deposited on : unknown  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20029824
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.


























Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1692 (3.00-3.00)
Clashscore	112137	2037 (3.00-3.00)
Ramachandran outliers	110173	1973 (3.00-3.00)
Sidechain outliers	110143	1976 (3.00-3.00)
RSRZ outliers	101464	1716 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	
1	B	240	
1	C	240	
1	D	240	
1	E	240	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	240	
1	G	240	
1	O	240	
1	P	240	
1	Q	240	
1	R	240	
1	S	240	
1	T	240	
1	U	240	
2	H	240	
2	I	240	
2	J	240	
2	K	240	
2	L	240	
2	M	240	
2	N	240	
2	V	240	
2	W	240	
2	X	240	
2	Y	240	
2	Z	240	
2	a	240	
2	b	240	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 47136 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	218	Total	C	N	O	S	0	0	0
			1677	1050	306	317	4			
1	B	215	Total	C	N	O	S	0	0	0
			1660	1041	303	312	4			
1	C	216	Total	C	N	O	S	0	0	0
			1664	1043	304	313	4			
1	D	217	Total	C	N	O	S	0	0	0
			1670	1046	305	315	4			
1	E	217	Total	C	N	O	S	0	0	0
			1671	1047	305	315	4			
1	F	218	Total	C	N	O	S	0	0	0
			1678	1050	306	318	4			
1	G	216	Total	C	N	O	S	0	0	0
			1662	1040	304	314	4			
1	O	216	Total	C	N	O	S	0	0	0
			1662	1040	304	314	4			
1	P	217	Total	C	N	O	S	0	0	0
			1674	1050	305	315	4			
1	Q	218	Total	C	N	O	S	0	0	0
			1679	1051	306	318	4			
1	R	217	Total	C	N	O	S	0	0	0
			1671	1047	305	315	4			
1	S	219	Total	C	N	O	S	0	0	0
			1683	1053	307	319	4			
1	T	216	Total	C	N	O	S	0	0	0
			1664	1043	304	313	4			
1	U	216	Total	C	N	O	S	0	0	0
			1664	1043	304	313	4			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	MET	-	initiating methionine	UNP A5U4D5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	9	MET	-	initiating methionine	UNP A5U4D5
C	9	MET	-	initiating methionine	UNP A5U4D5
D	9	MET	-	initiating methionine	UNP A5U4D5
E	9	MET	-	initiating methionine	UNP A5U4D5
F	9	MET	-	initiating methionine	UNP A5U4D5
G	9	MET	-	initiating methionine	UNP A5U4D5
O	9	MET	-	initiating methionine	UNP A5U4D5
P	9	MET	-	initiating methionine	UNP A5U4D5
Q	9	MET	-	initiating methionine	UNP A5U4D5
R	9	MET	-	initiating methionine	UNP A5U4D5
S	9	MET	-	initiating methionine	UNP A5U4D5
T	9	MET	-	initiating methionine	UNP A5U4D5
U	9	MET	-	initiating methionine	UNP A5U4D5

- Molecule 2 is a protein called Proteasome subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	222	Total	C	N	O	S	0	0	0
			1638	1027	282	324	5			
2	I	222	Total	C	N	O	S	0	0	0
			1638	1027	282	324	5			
2	J	222	Total	C	N	O	S	0	0	0
			1638	1027	282	324	5			
2	K	223	Total	C	N	O	S	0	0	0
			1642	1029	283	325	5			
2	L	223	Total	C	N	O	S	0	0	0
			1642	1029	283	325	5			
2	M	222	Total	C	N	O	S	0	0	0
			1638	1027	282	324	5			
2	N	223	Total	C	N	O	S	0	0	0
			1642	1029	283	325	5			
2	V	223	Total	C	N	O	S	0	0	0
			1642	1029	283	325	5			
2	W	223	Total	C	N	O	S	0	0	0
			1642	1029	283	325	5			
2	X	222	Total	C	N	O	S	0	0	0
			1638	1027	282	324	5			
2	Y	223	Total	C	N	O	S	0	0	0
			1642	1029	283	325	5			
2	Z	222	Total	C	N	O	S	0	0	0
			1638	1027	282	324	5			
2	a	223	Total	C	N	O	S	0	0	0
			1642	1029	283	325	5			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	b	223	Total	C	N	O	S	0	0	0
			1642	1029	283	325	5			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	235	HIS	-	expression tag	UNP A5U4D6
H	236	HIS	-	expression tag	UNP A5U4D6
H	237	HIS	-	expression tag	UNP A5U4D6
H	238	HIS	-	expression tag	UNP A5U4D6
H	239	HIS	-	expression tag	UNP A5U4D6
H	240	HIS	-	expression tag	UNP A5U4D6
I	235	HIS	-	expression tag	UNP A5U4D6
I	236	HIS	-	expression tag	UNP A5U4D6
I	237	HIS	-	expression tag	UNP A5U4D6
I	238	HIS	-	expression tag	UNP A5U4D6
I	239	HIS	-	expression tag	UNP A5U4D6
I	240	HIS	-	expression tag	UNP A5U4D6
J	235	HIS	-	expression tag	UNP A5U4D6
J	236	HIS	-	expression tag	UNP A5U4D6
J	237	HIS	-	expression tag	UNP A5U4D6
J	238	HIS	-	expression tag	UNP A5U4D6
J	239	HIS	-	expression tag	UNP A5U4D6
J	240	HIS	-	expression tag	UNP A5U4D6
K	235	HIS	-	expression tag	UNP A5U4D6
K	236	HIS	-	expression tag	UNP A5U4D6
K	237	HIS	-	expression tag	UNP A5U4D6
K	238	HIS	-	expression tag	UNP A5U4D6
K	239	HIS	-	expression tag	UNP A5U4D6
K	240	HIS	-	expression tag	UNP A5U4D6
L	235	HIS	-	expression tag	UNP A5U4D6
L	236	HIS	-	expression tag	UNP A5U4D6
L	237	HIS	-	expression tag	UNP A5U4D6
L	238	HIS	-	expression tag	UNP A5U4D6
L	239	HIS	-	expression tag	UNP A5U4D6
L	240	HIS	-	expression tag	UNP A5U4D6
M	235	HIS	-	expression tag	UNP A5U4D6
M	236	HIS	-	expression tag	UNP A5U4D6
M	237	HIS	-	expression tag	UNP A5U4D6
M	238	HIS	-	expression tag	UNP A5U4D6
M	239	HIS	-	expression tag	UNP A5U4D6
M	240	HIS	-	expression tag	UNP A5U4D6

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
N	235	HIS	-	expression tag	UNP A5U4D6
N	236	HIS	-	expression tag	UNP A5U4D6
N	237	HIS	-	expression tag	UNP A5U4D6
N	238	HIS	-	expression tag	UNP A5U4D6
N	239	HIS	-	expression tag	UNP A5U4D6
N	240	HIS	-	expression tag	UNP A5U4D6
V	235	HIS	-	expression tag	UNP A5U4D6
V	236	HIS	-	expression tag	UNP A5U4D6
V	237	HIS	-	expression tag	UNP A5U4D6
V	238	HIS	-	expression tag	UNP A5U4D6
V	239	HIS	-	expression tag	UNP A5U4D6
V	240	HIS	-	expression tag	UNP A5U4D6
W	235	HIS	-	expression tag	UNP A5U4D6
W	236	HIS	-	expression tag	UNP A5U4D6
W	237	HIS	-	expression tag	UNP A5U4D6
W	238	HIS	-	expression tag	UNP A5U4D6
W	239	HIS	-	expression tag	UNP A5U4D6
W	240	HIS	-	expression tag	UNP A5U4D6
X	235	HIS	-	expression tag	UNP A5U4D6
X	236	HIS	-	expression tag	UNP A5U4D6
X	237	HIS	-	expression tag	UNP A5U4D6
X	238	HIS	-	expression tag	UNP A5U4D6
X	239	HIS	-	expression tag	UNP A5U4D6
X	240	HIS	-	expression tag	UNP A5U4D6
Y	235	HIS	-	expression tag	UNP A5U4D6
Y	236	HIS	-	expression tag	UNP A5U4D6
Y	237	HIS	-	expression tag	UNP A5U4D6
Y	238	HIS	-	expression tag	UNP A5U4D6
Y	239	HIS	-	expression tag	UNP A5U4D6
Y	240	HIS	-	expression tag	UNP A5U4D6
Z	235	HIS	-	expression tag	UNP A5U4D6
Z	236	HIS	-	expression tag	UNP A5U4D6
Z	237	HIS	-	expression tag	UNP A5U4D6
Z	238	HIS	-	expression tag	UNP A5U4D6
Z	239	HIS	-	expression tag	UNP A5U4D6
Z	240	HIS	-	expression tag	UNP A5U4D6
a	235	HIS	-	expression tag	UNP A5U4D6
a	236	HIS	-	expression tag	UNP A5U4D6
a	237	HIS	-	expression tag	UNP A5U4D6
a	238	HIS	-	expression tag	UNP A5U4D6
a	239	HIS	-	expression tag	UNP A5U4D6
a	240	HIS	-	expression tag	UNP A5U4D6

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
b	235	HIS	-	expression tag	UNP A5U4D6
b	236	HIS	-	expression tag	UNP A5U4D6
b	237	HIS	-	expression tag	UNP A5U4D6
b	238	HIS	-	expression tag	UNP A5U4D6
b	239	HIS	-	expression tag	UNP A5U4D6
b	240	HIS	-	expression tag	UNP A5U4D6

- Molecule 3 is (2 {S})- {N}-[(2 {S})-3-methoxy-1-(naphthalen-1-ylmethylamino)-1-oxidanylidene-propan-2-yl]-4-oxidanylidene-2-(3-phenylpropanoylamino)-4-piperidin-1-yl-butanamide (three-letter code: 7J0) (formula: C<sub>33</sub>H<sub>40</sub>N<sub>4</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	H	1	Total	C	N	O	0	0
			42	33	4	5		
3	I	1	Total	C	N	O	0	0
			42	33	4	5		
3	J	1	Total	C	N	O	0	0
			42	33	4	5		
3	K	1	Total	C	N	O	0	0
			42	33	4	5		
3	L	1	Total	C	N	O	0	0
			42	33	4	5		
3	M	1	Total	C	N	O	0	0
			42	33	4	5		
3	N	1	Total	C	N	O	0	0
			42	33	4	5		

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	V	1	Total	C	N	O	0	0
			42	33	4	5		
3	W	1	Total	C	N	O	0	0
			42	33	4	5		
3	X	1	Total	C	N	O	0	0
			42	33	4	5		
3	Y	1	Total	C	N	O	0	0
			42	33	4	5		
3	Z	1	Total	C	N	O	0	0
			42	33	4	5		
3	a	1	Total	C	N	O	0	0
			42	33	4	5		
3	b	1	Total	C	N	O	0	0
			42	33	4	5		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	9	Total	O	0	0
			9	9		
4	B	2	Total	O	0	0
			2	2		
4	C	5	Total	O	0	0
			5	5		
4	D	3	Total	O	0	0
			3	3		
4	E	5	Total	O	0	0
			5	5		
4	F	8	Total	O	0	0
			8	8		
4	G	7	Total	O	0	0
			7	7		
4	H	7	Total	O	0	0
			7	7		
4	I	8	Total	O	0	0
			8	8		
4	J	7	Total	O	0	0
			7	7		
4	K	12	Total	O	0	0
			12	12		
4	L	5	Total	O	0	0
			5	5		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	M	12	Total 12	O 12	0	0
4	N	7	Total 7	O 7	0	0
4	O	6	Total 6	O 6	0	0
4	P	6	Total 6	O 6	0	0
4	Q	5	Total 5	O 5	0	0
4	R	8	Total 8	O 8	0	0
4	S	9	Total 9	O 9	0	0
4	T	6	Total 6	O 6	0	0
4	U	7	Total 7	O 7	0	0
4	V	9	Total 9	O 9	0	0
4	W	8	Total 8	O 8	0	0
4	X	8	Total 8	O 8	0	0
4	Y	11	Total 11	O 11	0	0
4	Z	11	Total 11	O 11	0	0
4	a	6	Total 6	O 6	0	0
4	b	8	Total 8	O 8	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha








- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha



- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha





• Molecule 2: Proteasome subunit beta

Chain H: 82% 11% 8%




• Molecule 2: Proteasome subunit beta

Chain I: 84% 9% 8%



• Molecule 2: Proteasome subunit beta

Chain J: 81% 12% 8%



• Molecule 2: Proteasome subunit beta

Chain K: 82% 11% 7%




• Molecule 2: Proteasome subunit beta



Chain L: 86% 7% 7%








● Molecule 2: Proteasome subunit beta



● Molecule 2: Proteasome subunit beta



● Molecule 2: Proteasome subunit beta





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.65Å 197.48Å 164.81Å 90.00° 103.05° 90.00°	Depositor
Resolution (Å)	51.09 – 3.00 51.09 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (51.09-3.00) 92.7 (51.09-3.00)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.25 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, $R_{free}$	0.182 , 0.234 0.182 , 0.229	Depositor DCC
$R_{free}$ test set	6856 reflections (4.93%)	DCC
Wilson B-factor (Å <sup>2</sup> )	37.5	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 44.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	47136	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7J0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/1701	0.50	0/2297
1	B	0.26	0/1684	0.46	0/2274
1	C	0.26	0/1688	0.46	0/2279
1	D	0.26	0/1694	0.48	1/2287 (0.0%)
1	E	0.29	0/1695	0.50	1/2289 (0.0%)
1	F	0.27	0/1702	0.48	0/2298
1	G	0.26	0/1686	0.48	0/2276
1	O	0.27	0/1686	0.48	0/2276
1	P	0.27	0/1699	0.49	0/2295
1	Q	0.26	0/1703	0.48	0/2300
1	R	0.29	0/1695	0.46	0/2289
1	S	0.27	0/1707	0.47	0/2305
1	T	0.27	0/1688	0.48	0/2279
1	U	0.26	0/1688	0.50	1/2279 (0.0%)
2	H	0.26	0/1662	0.49	0/2254
2	I	0.26	0/1662	0.48	0/2254
2	J	0.26	0/1662	0.50	0/2254
2	K	0.27	0/1666	0.50	0/2259
2	L	0.27	0/1666	0.50	0/2259
2	M	0.26	0/1662	0.49	0/2254
2	N	0.27	0/1666	0.50	0/2259
2	V	0.27	0/1666	0.51	0/2259
2	W	0.27	0/1666	0.50	0/2259
2	X	0.26	0/1662	0.49	0/2254
2	Y	0.27	0/1666	0.49	0/2259
2	Z	0.26	0/1662	0.49	0/2254
2	a	0.27	0/1666	0.49	0/2259
2	b	0.26	0/1666	0.48	0/2259
All	All	0.27	0/47016	0.49	3/63619 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	U	203	LEU	CA-CB-CG	7.12	131.69	115.30
1	D	40	LEU	CA-CB-CG	5.23	127.34	115.30
1	E	33	LEU	CA-CB-CG	5.09	127.00	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1677	0	1680	29	0
1	B	1660	0	1665	27	0
1	C	1664	0	1668	22	0
1	D	1670	0	1673	16	0
1	E	1671	0	1675	32	0
1	F	1678	0	1677	40	0
1	G	1662	0	1662	24	0
1	O	1662	0	1662	28	0
1	P	1674	0	1680	38	0
1	Q	1679	0	1679	36	0
1	R	1671	0	1675	21	0
1	S	1683	0	1682	21	0
1	T	1664	0	1668	25	0
1	U	1664	0	1668	26	0
2	H	1638	0	1633	19	0
2	I	1638	0	1633	14	0
2	J	1638	0	1633	21	0
2	K	1642	0	1636	15	0
2	L	1642	0	1636	12	0
2	M	1638	0	1633	13	0
2	N	1642	0	1636	22	0
2	V	1642	0	1636	15	0
2	W	1642	0	1636	20	0
2	X	1638	0	1633	20	0
2	Y	1642	0	1636	23	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Z	1638	0	1633	19	0
2	a	1642	0	1636	0	0
2	b	1642	0	1636	0	0
3	H	42	0	0	0	0
3	I	42	0	0	1	0
3	J	42	0	0	1	0
3	K	42	0	0	0	0
3	L	42	0	0	0	0
3	M	42	0	0	0	0
3	N	42	0	0	0	0
3	V	42	0	0	0	0
3	W	42	0	0	0	0
3	X	42	0	0	0	0
3	Y	42	0	0	0	0
3	Z	42	0	0	0	0
3	a	42	0	0	0	0
3	b	42	0	0	0	0
4	A	9	0	0	0	0
4	B	2	0	0	0	0
4	C	5	0	0	0	0
4	D	3	0	0	0	0
4	E	5	0	0	0	0
4	F	8	0	0	0	0
4	G	7	0	0	0	0
4	H	7	0	0	0	0
4	I	8	0	0	0	0
4	J	7	0	0	0	0
4	K	12	0	0	0	0
4	L	5	0	0	0	0
4	M	12	0	0	0	0
4	N	7	0	0	0	0
4	O	6	0	0	0	0
4	P	6	0	0	0	0
4	Q	5	0	0	0	0
4	R	8	0	0	1	0
4	S	9	0	0	0	0
4	T	6	0	0	0	0
4	U	7	0	0	0	0
4	V	9	0	0	0	0
4	W	8	0	0	1	0
4	X	8	0	0	1	0
4	Y	11	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	Z	11	0	0	0	0
4	a	6	0	0	0	0
4	b	8	0	0	0	0
All	All	47136	0	46300	544	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 544 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:9:MET:N	1:E:15:GLU:OE1	2.14	0.81
1:R:16:ARG:NH2	1:R:114:GLN:O	2.13	0.80
2:V:29:ARG:NH1	2:W:134:GLU:OE2	2.15	0.80
1:P:219:ARG:NH2	2:W:64:GLU:OE1	2.14	0.79
1:Q:140:ARG:NH2	1:Q:155:VAL:O	2.13	0.78

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/240 (89%)	207 (97%)	7 (3%)	0	100	100
1	B	211/240 (88%)	206 (98%)	5 (2%)	0	100	100
1	C	212/240 (88%)	206 (97%)	6 (3%)	0	100	100
1	D	213/240 (89%)	206 (97%)	7 (3%)	0	100	100
1	E	213/240 (89%)	206 (97%)	7 (3%)	0	100	100
1	F	214/240 (89%)	206 (96%)	8 (4%)	0	100	100
1	G	212/240 (88%)	207 (98%)	5 (2%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	O	212/240 (88%)	205 (97%)	6 (3%)	1 (0%)	32	74
1	P	213/240 (89%)	205 (96%)	8 (4%)	0	100	100
1	Q	214/240 (89%)	207 (97%)	7 (3%)	0	100	100
1	R	213/240 (89%)	205 (96%)	8 (4%)	0	100	100
1	S	215/240 (90%)	205 (95%)	10 (5%)	0	100	100
1	T	212/240 (88%)	204 (96%)	8 (4%)	0	100	100
1	U	212/240 (88%)	206 (97%)	6 (3%)	0	100	100
2	H	220/240 (92%)	216 (98%)	4 (2%)	0	100	100
2	I	220/240 (92%)	214 (97%)	6 (3%)	0	100	100
2	J	220/240 (92%)	216 (98%)	4 (2%)	0	100	100
2	K	221/240 (92%)	216 (98%)	5 (2%)	0	100	100
2	L	221/240 (92%)	217 (98%)	4 (2%)	0	100	100
2	M	220/240 (92%)	215 (98%)	4 (2%)	1 (0%)	32	74
2	N	221/240 (92%)	217 (98%)	4 (2%)	0	100	100
2	V	221/240 (92%)	217 (98%)	4 (2%)	0	100	100
2	W	221/240 (92%)	216 (98%)	5 (2%)	0	100	100
2	X	220/240 (92%)	216 (98%)	4 (2%)	0	100	100
2	Y	221/240 (92%)	215 (97%)	6 (3%)	0	100	100
2	Z	220/240 (92%)	213 (97%)	7 (3%)	0	100	100
2	a	221/240 (92%)	217 (98%)	4 (2%)	0	100	100
2	b	221/240 (92%)	217 (98%)	4 (2%)	0	100	100
All	All	6068/6720 (90%)	5903 (97%)	163 (3%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	O	205	VAL
2	M	114	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/184 (91%)	166 (99%)	1 (1%)	89	96
1	B	165/184 (90%)	164 (99%)	1 (1%)	89	96
1	C	165/184 (90%)	165 (100%)	0	100	100
1	D	166/184 (90%)	163 (98%)	3 (2%)	64	89
1	E	166/184 (90%)	165 (99%)	1 (1%)	89	96
1	F	167/184 (91%)	167 (100%)	0	100	100
1	G	165/184 (90%)	165 (100%)	0	100	100
1	O	165/184 (90%)	164 (99%)	1 (1%)	89	96
1	P	167/184 (91%)	165 (99%)	2 (1%)	75	93
1	Q	167/184 (91%)	166 (99%)	1 (1%)	89	96
1	R	166/184 (90%)	165 (99%)	1 (1%)	89	96
1	S	167/184 (91%)	166 (99%)	1 (1%)	89	96
1	T	165/184 (90%)	165 (100%)	0	100	100
1	U	165/184 (90%)	164 (99%)	1 (1%)	89	96
2	H	165/178 (93%)	165 (100%)	0	100	100
2	I	165/178 (93%)	165 (100%)	0	100	100
2	J	165/178 (93%)	165 (100%)	0	100	100
2	K	165/178 (93%)	165 (100%)	0	100	100
2	L	165/178 (93%)	165 (100%)	0	100	100
2	M	165/178 (93%)	165 (100%)	0	100	100
2	N	165/178 (93%)	165 (100%)	0	100	100
2	V	165/178 (93%)	165 (100%)	0	100	100
2	W	165/178 (93%)	165 (100%)	0	100	100
2	X	165/178 (93%)	165 (100%)	0	100	100
2	Y	165/178 (93%)	165 (100%)	0	100	100
2	Z	165/178 (93%)	165 (100%)	0	100	100
2	a	165/178 (93%)	165 (100%)	0	100	100
2	b	165/178 (93%)	165 (100%)	0	100	100
All	All	4633/5068 (91%)	4620 (100%)	13 (0%)	94	98

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	33	LEU
1	O	33	LEU
1	R	101	ASN
1	D	179	ASP
1	Q	75	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	X	90	ASN
2	a	115	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	7J0	H	301	-	45,45,45	1.88	12 (26%)	58,59,59	1.48	9 (15%)
3	7J0	I	301	-	45,45,45	1.88	12 (26%)	58,59,59	1.47	9 (15%)



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	7J0	J	301	-	45,45,45	1.88	12 (26%)	58,59,59	1.54	10 (17%)
3	7J0	K	301	-	45,45,45	1.89	12 (26%)	58,59,59	1.49	9 (15%)
3	7J0	L	301	-	45,45,45	1.86	12 (26%)	58,59,59	1.46	8 (13%)
3	7J0	M	301	-	45,45,45	1.88	11 (24%)	58,59,59	1.47	8 (13%)
3	7J0	N	301	-	45,45,45	1.88	12 (26%)	58,59,59	1.48	10 (17%)
3	7J0	V	301	-	45,45,45	1.86	12 (26%)	58,59,59	1.47	8 (13%)
3	7J0	W	301	-	45,45,45	1.89	12 (26%)	58,59,59	1.47	9 (15%)
3	7J0	X	301	-	45,45,45	1.87	12 (26%)	58,59,59	1.47	9 (15%)
3	7J0	Y	301	-	45,45,45	1.87	12 (26%)	58,59,59	1.47	8 (13%)
3	7J0	Z	301	-	45,45,45	1.87	12 (26%)	58,59,59	1.46	8 (13%)
3	7J0	a	301	-	45,45,45	1.88	12 (26%)	58,59,59	1.48	9 (15%)
3	7J0	b	301	-	45,45,45	1.87	12 (26%)	58,59,59	1.49	9 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	7J0	H	301	-	-	0/37/45/45	0/4/4/4
3	7J0	I	301	-	-	0/37/45/45	0/4/4/4
3	7J0	J	301	-	-	0/37/45/45	0/4/4/4
3	7J0	K	301	-	-	0/37/45/45	0/4/4/4
3	7J0	L	301	-	-	0/37/45/45	0/4/4/4
3	7J0	M	301	-	-	0/37/45/45	0/4/4/4
3	7J0	N	301	-	-	0/37/45/45	0/4/4/4
3	7J0	V	301	-	-	0/37/45/45	0/4/4/4
3	7J0	W	301	-	-	0/37/45/45	0/4/4/4
3	7J0	X	301	-	-	0/37/45/45	0/4/4/4
3	7J0	Y	301	-	-	0/37/45/45	0/4/4/4
3	7J0	Z	301	-	-	0/37/45/45	0/4/4/4
3	7J0	a	301	-	-	0/37/45/45	0/4/4/4
3	7J0	b	301	-	-	0/37/45/45	0/4/4/4

The worst 5 of 167 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	J	301	7J0	C19-C04	2.00	1.58	1.52
3	L	301	7J0	C19-C04	2.00	1.58	1.52
3	N	301	7J0	C19-C04	2.01	1.58	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	Z	301	7J0	C19-C04	2.01	1.58	1.52
3	a	301	7J0	C19-C04	2.01	1.58	1.52

The worst 5 of 123 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	J	301	7J0	O42-C33-C34	-2.71	116.91	122.01
3	L	301	7J0	O42-C33-C34	-2.71	116.91	122.01
3	I	301	7J0	O42-C33-C34	-2.71	116.92	122.01
3	K	301	7J0	O42-C33-C34	-2.70	116.94	122.01
3	M	301	7J0	O42-C33-C34	-2.69	116.96	122.01

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	301	7J0	1	0
3	J	301	7J0	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	218/240 (90%)	-0.41	1 (0%) 90 74	17, 36, 59, 80	0
1	B	215/240 (89%)	-0.23	0 100 100	21, 45, 77, 92	0
1	C	216/240 (90%)	-0.22	0 100 100	19, 49, 81, 101	0
1	D	217/240 (90%)	-0.28	0 100 100	21, 45, 75, 83	0
1	E	217/240 (90%)	-0.41	0 100 100	16, 42, 65, 95	0
1	F	218/240 (90%)	-0.23	3 (1%) 75 49	19, 47, 81, 113	0
1	G	216/240 (90%)	-0.45	0 100 100	15, 37, 66, 82	0
1	O	216/240 (90%)	-0.09	1 (0%) 90 74	20, 51, 87, 102	0
1	P	217/240 (90%)	-0.24	2 (0%) 84 61	20, 47, 77, 111	0
1	Q	218/240 (90%)	-0.27	3 (1%) 75 49	23, 44, 75, 115	0
1	R	217/240 (90%)	-0.32	1 (0%) 90 74	17, 45, 71, 92	0
1	S	219/240 (91%)	-0.37	3 (1%) 75 49	19, 38, 76, 101	0
1	T	216/240 (90%)	-0.20	1 (0%) 90 74	21, 46, 76, 85	0
1	U	216/240 (90%)	-0.31	1 (0%) 90 74	20, 41, 73, 101	0
2	H	222/240 (92%)	-0.51	0 100 100	19, 29, 50, 81	0
2	I	222/240 (92%)	-0.53	0 100 100	16, 27, 47, 65	0
2	J	222/240 (92%)	-0.57	0 100 100	19, 30, 54, 86	0
2	K	223/240 (92%)	-0.61	0 100 100	18, 30, 53, 70	0
2	L	223/240 (92%)	-0.56	0 100 100	17, 27, 51, 72	0
2	M	222/240 (92%)	-0.51	0 100 100	18, 30, 53, 82	0
2	N	223/240 (92%)	-0.53	0 100 100	18, 32, 59, 85	0
2	V	223/240 (92%)	-0.54	0 100 100	19, 28, 48, 64	0
2	W	223/240 (92%)	-0.51	0 100 100	18, 31, 57, 73	0
2	X	222/240 (92%)	-0.54	0 100 100	19, 30, 54, 77	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
2	Y	223/240 (92%)	-0.56	0 100 100	17, 29, 55, 87	0
2	Z	222/240 (92%)	-0.57	0 100 100	16, 28, 53, 76	0
2	a	223/240 (92%)	-0.51	0 100 100	19, 32, 58, 77	0
2	b	223/240 (92%)	-0.43	0 100 100	18, 30, 56, 77	0
All	All	6152/6720 (91%)	-0.41	16 (0%) 93 82	15, 35, 71, 115	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	169	GLU	3.6
1	S	193	ALA	3.3
1	Q	236	ASP	3.3
1	P	201	PRO	3.2
1	F	236	ASP	3.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	7J0	J	301	42/42	0.95	0.21	1.92	19,26,60,81	0
3	7J0	N	301	42/42	0.94	0.23	1.72	20,29,53,59	0
3	7J0	M	301	42/42	0.94	0.20	1.51	14,24,49,63	0
3	7J0	Z	301	42/42	0.95	0.19	1.46	19,25,38,50	0
3	7J0	L	301	42/42	0.95	0.19	1.30	14,24,39,52	0

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	7J0	H	301	42/42	0.95	0.21	1.28	15,25,45,66	0
3	7J0	Y	301	42/42	0.96	0.20	1.15	14,24,53,75	0
3	7J0	a	301	42/42	0.95	0.20	1.08	21,27,48,62	0
3	7J0	b	301	42/42	0.95	0.21	0.94	16,24,38,41	0
3	7J0	W	301	42/42	0.95	0.19	0.93	14,24,43,55	0
3	7J0	I	301	42/42	0.95	0.19	0.79	16,26,46,64	0
3	7J0	X	301	42/42	0.95	0.19	0.78	15,25,41,47	0
3	7J0	K	301	42/42	0.94	0.20	0.73	21,31,52,64	0
3	7J0	V	301	42/42	0.95	0.20	0.54	21,30,51,62	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.