



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 3, 2017 – 04:37 PM EDT

PDB ID : 5TUI
Title : Crystal structure of tetracycline deconstructase Tet(50) in complex with chlortetracycline
Authors : Park, J.; Tolia, N.H.
Deposited on : unknown
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1762 (1.76-1.76)
Clashscore	112137	1889 (1.76-1.76)
Ramachandran outliers	110173	1868 (1.76-1.76)
Sidechain outliers	110143	1868 (1.76-1.76)
RSRZ outliers	101464	1770 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	409	<div> <div>4%</div> <div>91%</div> <div>5%</div> </div>
1	B	409	<div> <div>4%</div> <div>83%</div> <div>7%</div> <div>10%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	403	-	-	-	X
2	SO4	B	402	-	-	-	X
2	SO4	B	403	-	-	-	X
4	CTC	B	405	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 12688 atoms, of which 6049 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tetracycline destructase Tet(50).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	388	Total	C	H	N	O	S	0	3	0
			6139	1964	3055	521	582	17			
1	B	370	Total	C	H	N	O	S	0	6	0
			5854	1873	2912	497	556	16			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP A0A059WYP6
A	-19	GLY	-	expression tag	UNP A0A059WYP6
A	-18	SER	-	expression tag	UNP A0A059WYP6
A	-17	SER	-	expression tag	UNP A0A059WYP6
A	-16	HIS	-	expression tag	UNP A0A059WYP6
A	-15	HIS	-	expression tag	UNP A0A059WYP6
A	-14	HIS	-	expression tag	UNP A0A059WYP6
A	-13	HIS	-	expression tag	UNP A0A059WYP6
A	-12	HIS	-	expression tag	UNP A0A059WYP6
A	-11	HIS	-	expression tag	UNP A0A059WYP6
A	-10	SER	-	expression tag	UNP A0A059WYP6
A	-9	SER	-	expression tag	UNP A0A059WYP6
A	-8	GLY	-	expression tag	UNP A0A059WYP6
A	-7	LEU	-	expression tag	UNP A0A059WYP6
A	-6	VAL	-	expression tag	UNP A0A059WYP6
A	-5	PRO	-	expression tag	UNP A0A059WYP6
A	-4	ARG	-	expression tag	UNP A0A059WYP6
A	-3	GLY	-	expression tag	UNP A0A059WYP6
A	-2	SER	-	expression tag	UNP A0A059WYP6
A	-1	HIS	-	expression tag	UNP A0A059WYP6
A	0	MET	-	expression tag	UNP A0A059WYP6
B	-20	MET	-	expression tag	UNP A0A059WYP6
B	-19	GLY	-	expression tag	UNP A0A059WYP6
B	-18	SER	-	expression tag	UNP A0A059WYP6
B	-17	SER	-	expression tag	UNP A0A059WYP6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	expression tag	UNP A0A059WYP6
B	-15	HIS	-	expression tag	UNP A0A059WYP6
B	-14	HIS	-	expression tag	UNP A0A059WYP6
B	-13	HIS	-	expression tag	UNP A0A059WYP6
B	-12	HIS	-	expression tag	UNP A0A059WYP6
B	-11	HIS	-	expression tag	UNP A0A059WYP6
B	-10	SER	-	expression tag	UNP A0A059WYP6
B	-9	SER	-	expression tag	UNP A0A059WYP6
B	-8	GLY	-	expression tag	UNP A0A059WYP6
B	-7	LEU	-	expression tag	UNP A0A059WYP6
B	-6	VAL	-	expression tag	UNP A0A059WYP6
B	-5	PRO	-	expression tag	UNP A0A059WYP6
B	-4	ARG	-	expression tag	UNP A0A059WYP6
B	-3	GLY	-	expression tag	UNP A0A059WYP6
B	-2	SER	-	expression tag	UNP A0A059WYP6
B	-1	HIS	-	expression tag	UNP A0A059WYP6
B	0	MET	-	expression tag	UNP A0A059WYP6

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	0	0
			84	27	31	9	15	2		
3	B	1	Total	C	H	N	O	P	0	0
			84	27	31	9	15	2		

- Molecule 4 is 7-CHLOROTETRACYCLINE (three-letter code: CTC) (formula: C₂₂H₂₃ClN₂O₈).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total	C	Cl	H	N	O	0	0
			53	22	1	20	2	8		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	234	Total O 234 234	0	0
5	B	210	Total O 210 210	0	0

- Molecule 1: Tetracycline destructase Tet(50)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.10Å 107.22Å 152.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.77 – 1.75 19.77 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.8 (19.77-1.75) 98.8 (19.77-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 1.76Å)	Xtriage
Refinement program	PHENIX (1.10 _2155: ???)	Depositor
R, R_{free}	0.178 , 0.219 0.177 , 0.219	Depositor DCC
R_{free} test set	3010 reflections (3.57%)	DCC
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12688	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FAD, CTC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	2/3152 (0.1%)	0.71	0/4250
1	B	0.55	1/3015 (0.0%)	0.69	3/4063 (0.1%)
All	All	0.56	3/6167 (0.0%)	0.70	3/8313 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	4
All	All	0	7

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	74	CYS	CB-SG	-7.69	1.69	1.82
1	A	292	TYR	CB-CG	-5.27	1.43	1.51
1	A	74	CYS	CB-SG	-5.16	1.73	1.81

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	205	LEU	CA-CB-CG	6.12	129.38	115.30
1	B	49	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	B	156	ASP	CB-CG-OD1	5.29	123.06	118.30

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	155	ALA	Mainchain
1	A	188	ASN	Mainchain
1	A	291	ALA	Mainchain
1	B	155	ALA	Mainchain
1	B	188	ASN	Mainchain
1	B	291	ALA	Mainchain
1	B	71	GLN	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3084	3055	3067	9	1
1	B	2942	2912	2921	18	0
2	A	15	0	0	0	0
2	B	15	0	0	0	0
3	A	53	31	31	0	0
3	B	53	31	31	0	0
4	B	33	20	20	3	0
5	A	234	0	0	1	0
5	B	210	0	0	5	0
All	All	6639	6049	6070	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:254:GLN:NE2	5:B:502:HOH:O	2.19	0.76
1:A:5:ILE:O	5:A:501:HOH:O	2.07	0.72
1:B:205:LEU:HG	4:B:405:CTC:O10	1.98	0.64
4:B:405:CTC:CL7	4:B:405:CTC:H6'1	2.38	0.60
1:B:87:GLU:OE1	5:B:501:HOH:O	2.15	0.60
1:B:40:ARG:NH1	5:B:503:HOH:O	2.38	0.57
1:B:169:LYS:HD2	5:B:578:HOH:O	2.05	0.56
1:B:158:ILE:HD11	1:B:272:THR:OG1	2.06	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:300:GLN:NE2	1:B:375:ILE:HD11	2.22	0.54
1:B:127:VAL:HG13	1:B:139[A]:VAL:HG13	1.91	0.53
1:A:200:GLU:CD	1:A:359:VAL:HG11	2.30	0.52
1:A:51[B]:ILE:HD13	1:A:382:ILE:HG22	1.92	0.52
1:B:372:LEU:O	1:B:375:ILE:HG22	2.12	0.50
1:A:142:LYS:HG2	5:B:510:HOH:O	2.13	0.48
1:B:297:LEU:HD13	1:B:349[B]:SER:HB2	1.96	0.48
1:A:51[B]:ILE:HD11	1:A:307:VAL:HG21	1.97	0.47
1:B:176[B]:ASN:OD1	1:B:272:THR:HG22	2.15	0.46
1:A:51[B]:ILE:HD13	1:A:382:ILE:CG2	2.49	0.43
1:B:221:PHE:N	4:B:405:CTC:H9	2.33	0.43
1:B:344:PHE:CE2	1:B:375:ILE:HD13	2.54	0.43
1:B:375:ILE:HD12	1:B:378:VAL:CG2	2.48	0.42
1:A:12:VAL:O	1:A:12:VAL:HG12	2.20	0.42
1:B:292:TYR:O	1:B:293:CYS:C	2.57	0.41
1:A:51[A]:ILE:HD13	1:A:382:ILE:HG23	2.01	0.41
1:B:198:LEU:HG	1:B:205:LEU:HD11	2.02	0.41
1:B:300:GLN:CD	1:B:375:ILE:HD11	2.40	0.41
1:A:292:TYR:O	1:A:293:CYS:C	2.60	0.40
1:B:6:LYS:HE2	1:B:149:TYR:CE1	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:GLN:OE1	1:A:361:LYS:HZ3[3_646]	1.57	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	389/409 (95%)	384 (99%)	5 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	370/409 (90%)	363 (98%)	7 (2%)	0	100	100
All	All	759/818 (93%)	747 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	332/347 (96%)	331 (100%)	1 (0%)	94	91
1	B	317/347 (91%)	317 (100%)	0	100	100
All	All	649/694 (94%)	648 (100%)	1 (0%)	94	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	401	-	4,4,4	0.13	0	6,6,6	0.29	0
2	SO4	A	402	-	4,4,4	0.23	0	6,6,6	0.29	0
2	SO4	A	403	-	4,4,4	0.23	0	6,6,6	0.20	0
3	FAD	A	404	-	50,58,58	1.63	11 (22%)	53,89,89	2.05	12 (22%)
2	SO4	B	401	-	4,4,4	0.27	0	6,6,6	0.23	0
2	SO4	B	402	-	4,4,4	0.19	0	6,6,6	0.10	0
2	SO4	B	403	-	4,4,4	0.26	0	6,6,6	0.35	0
3	FAD	B	404	-	50,58,58	1.51	8 (16%)	53,89,89	2.31	14 (26%)
4	CTC	B	405	-	34,36,36	1.61	5 (14%)	40,60,60	1.89	10 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	401	-	-	0/0/0/0	0/0/0/0
2	SO4	A	402	-	-	0/0/0/0	0/0/0/0
2	SO4	A	403	-	-	0/0/0/0	0/0/0/0
3	FAD	A	404	-	-	0/28/50/50	0/6/6/6
2	SO4	B	401	-	-	0/0/0/0	0/0/0/0
2	SO4	B	402	-	-	0/0/0/0	0/0/0/0
2	SO4	B	403	-	-	0/0/0/0	0/0/0/0
3	FAD	B	404	-	-	0/28/50/50	0/6/6/6
4	CTC	B	405	-	-	0/8/74/74	0/4/4/4

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	405	CTC	C4B-C1	-2.70	1.51	1.55
3	B	404	FAD	C1'-N10	-2.70	1.45	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	404	FAD	C1'-N10	-2.29	1.46	1.48
3	B	404	FAD	C10-N1	2.09	1.36	1.33
3	A	404	FAD	C5'-C4'	2.13	1.54	1.51
3	A	404	FAD	C10-N1	2.16	1.36	1.33
4	B	405	CTC	C7-CL7	2.18	1.78	1.73
3	A	404	FAD	O4B-C1B	2.33	1.44	1.41
3	A	404	FAD	C9A-N10	2.54	1.42	1.38
3	A	404	FAD	C4X-N5	2.83	1.37	1.33
3	A	404	FAD	C5A-C4A	2.93	1.47	1.40
3	B	404	FAD	C5A-C4A	2.97	1.47	1.40
3	B	404	FAD	C4X-N5	2.99	1.37	1.33
4	B	405	CTC	C6B-C6A	3.11	1.48	1.41
3	A	404	FAD	C8-C7	3.22	1.49	1.41
3	B	404	FAD	C8-C7	3.23	1.49	1.41
4	B	405	CTC	C6B-C10	3.29	1.47	1.41
3	B	404	FAD	C4-C4X	3.44	1.47	1.41
3	A	404	FAD	C9A-C5X	3.46	1.49	1.42
3	B	404	FAD	C9A-C5X	3.53	1.49	1.42
3	B	404	FAD	C4X-C10	4.30	1.48	1.41
3	A	404	FAD	C4-C4X	4.46	1.49	1.41
3	A	404	FAD	C4X-C10	4.57	1.49	1.41
4	B	405	CTC	C7-C6A	5.47	1.51	1.40

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	404	FAD	N3A-C2A-N1A	-8.86	121.14	128.86
3	A	404	FAD	N3A-C2A-N1A	-6.46	123.23	128.86
4	B	405	CTC	C2'-C2-C1	-5.82	114.27	120.89
4	B	405	CTC	C4A-C4B-C1	-3.81	106.68	111.05
3	B	404	FAD	C4'-C3'-C2'	-3.77	105.28	113.41
3	B	404	FAD	C4B-O4B-C1B	-3.30	106.26	109.77
3	B	404	FAD	O5'-P-O1P	-2.87	97.66	109.25
3	B	404	FAD	C1'-C2'-C3'	-2.86	101.64	109.82
3	A	404	FAD	C1'-N10-C10	-2.84	115.59	118.50
4	B	405	CTC	C8-C7-CL7	-2.74	112.70	118.39
3	B	404	FAD	C1'-N10-C10	-2.73	115.71	118.50
3	B	404	FAD	C4X-C10-N10	-2.71	118.64	120.52
4	B	405	CTC	O12-C12-C5B	-2.65	121.36	123.86
3	A	404	FAD	C4B-O4B-C1B	-2.62	106.98	109.77
4	B	405	CTC	O1-C1-C2	-2.60	118.34	123.63
3	B	404	FAD	C1B-N9A-C4A	-2.48	122.35	126.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	404	FAD	O5'-P-O1P	-2.48	99.24	109.25
3	A	404	FAD	O2'-C2'-C1'	-2.38	104.28	109.79
3	A	404	FAD	C4'-C3'-C2'	-2.19	108.69	113.41
4	B	405	CTC	C4B-C12-C5B	-2.00	120.85	123.05
3	A	404	FAD	O2A-PA-O1A	2.05	122.88	112.28
4	B	405	CTC	C2-C2'-N2'	2.26	123.63	119.07
3	B	404	FAD	C5X-C9A-N10	2.59	119.58	117.66
3	B	404	FAD	O2P-P-O1P	2.69	126.21	112.28
3	A	404	FAD	N6A-C6A-N1A	2.70	124.11	118.77
4	B	405	CTC	C4B-C4A-C4	2.70	115.33	111.64
4	B	405	CTC	C4B-C1-C2	2.90	120.49	116.01
4	B	405	CTC	O12-C12-C4B	2.91	117.74	113.34
3	B	404	FAD	C2A-N1A-C6A	2.93	123.89	118.77
3	A	404	FAD	O2P-P-O1P	3.00	127.80	112.28
3	B	404	FAD	C4X-N5-C5X	3.41	120.36	116.76
3	A	404	FAD	C4X-N5-C5X	3.68	120.65	116.76
3	A	404	FAD	C2-N1-C10	4.43	119.17	114.90
3	B	404	FAD	C2-N1-C10	5.46	120.16	114.90
3	B	404	FAD	C1'-N10-C9A	6.39	124.21	118.35
3	A	404	FAD	C1'-N10-C9A	7.21	124.95	118.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	405	CTC	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/409 (94%)	0.13	17 (4%) 35 42	13, 20, 39, 68	0
1	B	370/409 (90%)	0.07	18 (4%) 30 36	13, 21, 40, 63	0
All	All	758/818 (92%)	0.10	35 (4%) 33 39	13, 20, 40, 68	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	7.4
1	B	4	HIS	7.1
1	A	356	ASP	6.1
1	A	354	LEU	4.9
1	A	3	LYS	4.8
1	A	230	GLU	4.5
1	A	264	ASP	4.5
1	A	228	VAL	4.2
1	B	264	ASP	4.2
1	A	355	LYS	4.1
1	A	4	HIS	3.9
1	A	357	ASP	3.8
1	A	226	LYS	3.3
1	B	5	ILE	3.3
1	A	170	ASN	3.2
1	B	362	GLU	3.2
1	B	134	GLU	3.1
1	B	320	ASP	3.0
1	A	145	ARG	2.9
1	B	361	LYS	2.9
1	B	367	ARG	2.7
1	B	262	GLU	2.6
1	A	2	THR	2.5
1	B	347	TRP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	40	ARG	2.5
1	A	134	GLU	2.4
1	B	135	ASP	2.3
1	B	351	SER	2.3
1	B	40	ARG	2.2
1	A	142	LYS	2.2
1	B	363	ILE	2.2
1	B	118	ASP	2.2
1	B	202	ASN	2.2
1	B	117	ALA	2.0
1	B	277	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SO4	B	402	5/5	0.93	0.22	8.23	39,40,53,59	0
2	SO4	A	403	5/5	0.92	0.32	7.94	31,38,58,59	0
2	SO4	B	403	5/5	0.92	0.24	3.18	42,49,52,65	0
4	CTC	B	405	33/33	0.88	0.17	3.13	24,38,47,52	0
3	FAD	A	404	53/53	0.95	0.12	0.95	17,25,34,39	0
3	FAD	B	404	53/53	0.95	0.10	0.03	13,21,31,32	0
2	SO4	A	401	5/5	0.96	0.18	-	37,42,50,53	0
2	SO4	B	401	5/5	0.92	0.26	-	32,53,60,64	0
2	SO4	A	402	5/5	0.95	0.17	-	28,47,53,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.