



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 04:02 am GMT

PDB ID : 3TY8
Title : Crystal Structure of C. Thermocellum PNKP Ligase Domain Apo Form
Authors : Smith, P.; Wang, L.; Shuman, S.
Deposited on : 2011-09-23
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	413	 3% 66% 24% • 8%
1	B	413	 4% 61% 29% • 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	911	-	-	X	X
2	GOL	B	903	-	-	-	X

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polynucleotide 2',3'-cyclic phosphate phosphodiesterase / polynucleotide 5'-hydroxyl-kinase / polynucleotide 3'-phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	379	3019	1914	523	559	23	0	0	0
1	B	379	3064	1943	529	570	22	0	5	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	458	MET	-	EXPRESSION TAG	UNP A3DJ38
A	459	GLY	-	EXPRESSION TAG	UNP A3DJ38
A	460	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	461	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	462	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	463	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	464	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	465	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	466	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	467	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	468	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	469	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	470	SER	-	EXPRESSION TAG	UNP A3DJ38
A	471	SER	-	EXPRESSION TAG	UNP A3DJ38
A	472	GLY	-	EXPRESSION TAG	UNP A3DJ38
A	473	HIS	-	EXPRESSION TAG	UNP A3DJ38
A	474	ILE	-	EXPRESSION TAG	UNP A3DJ38
A	475	GLU	-	EXPRESSION TAG	UNP A3DJ38
A	476	GLY	-	EXPRESSION TAG	UNP A3DJ38
A	477	ARG	-	EXPRESSION TAG	UNP A3DJ38
A	478	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	458	MET	-	EXPRESSION TAG	UNP A3DJ38
B	459	GLY	-	EXPRESSION TAG	UNP A3DJ38
B	460	HIS	-	EXPRESSION TAG	UNP A3DJ38

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	461	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	462	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	463	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	464	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	465	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	466	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	467	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	468	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	469	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	470	SER	-	EXPRESSION TAG	UNP A3DJ38
B	471	SER	-	EXPRESSION TAG	UNP A3DJ38
B	472	GLY	-	EXPRESSION TAG	UNP A3DJ38
B	473	HIS	-	EXPRESSION TAG	UNP A3DJ38
B	474	ILE	-	EXPRESSION TAG	UNP A3DJ38
B	475	GLU	-	EXPRESSION TAG	UNP A3DJ38
B	476	GLY	-	EXPRESSION TAG	UNP A3DJ38
B	477	ARG	-	EXPRESSION TAG	UNP A3DJ38
B	478	HIS	-	EXPRESSION TAG	UNP A3DJ38

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is D(-)-TARTARIC ACID (three-letter code: TAR) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	4	6		
3	B	1	Total	C	O	0	0
			10	4	6		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	73	Total	O	0	0
			73	73		
4	B	42	Total	O	0	0
			42	42		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.14Å 94.16Å 76.03Å 90.00° 95.78° 90.00°	Depositor
Resolution (Å)	37.82 – 2.60 37.82 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.1 (37.82-2.60) 96.0 (37.82-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.61Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.195 , 0.241 0.187 , 0.234	Depositor DCC
R_{free} test set	1249 reflections (4.99%)	DCC
Wilson B-factor (Å ²)	45.7	Xtrriage
Anisotropy	0.572	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6236	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, TAR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/3084	0.47	0/4166
1	B	0.32	0/3130	0.47	0/4227
All	All	0.32	0/6214	0.47	0/8393

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3019	0	2959	110	0
1	B	3064	0	3008	123	0
2	A	12	0	16	5	0
2	B	6	0	8	0	0
3	A	10	0	4	1	0
3	B	10	0	4	0	0
4	A	73	0	0	0	0
4	B	42	0	0	0	0
All	All	6236	0	5999	226	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 226 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:810:ASP:HA	1:A:812:ASN:H	1.08	1.17
1:A:489:LEU:HD21	1:A:855:VAL:HG11	1.24	1.11
1:A:489:LEU:CD2	1:A:855:VAL:HG11	1.87	1.04
1:A:477:ARG:HA	1:A:859:LEU:HD13	1.39	1.03
1:A:810:ASP:CA	1:A:812:ASN:H	1.70	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/413 (91%)	361 (96%)	13 (4%)	1 (0%)	44	70
1	B	380/413 (92%)	364 (96%)	15 (4%)	1 (0%)	44	70
All	All	755/826 (91%)	725 (96%)	28 (4%)	2 (0%)	44	70

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	808	THR
1	A	732	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/361 (89%)	312 (97%)	10 (3%)	45	73
1	B	329/361 (91%)	317 (96%)	12 (4%)	40	68
All	All	651/722 (90%)	629 (97%)	22 (3%)	44	69

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	864	GLU
1	B	532[B]	HIS
1	B	786	LEU
1	B	501	THR
1	B	532[A]	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	532	HIS
1	A	643	GLN
1	A	719	HIS
1	B	719	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	910	-	5,5,5	0.30	0	5,5,5	0.26	0
2	GOL	A	911	-	5,5,5	0.32	0	5,5,5	0.34	0
3	TAR	A	912	-	3,9,9	1.83	1 (33%)	6,12,12	1.13	1 (16%)
3	TAR	B	901	-	3,9,9	1.64	0	6,12,12	0.91	0
2	GOL	B	903	-	5,5,5	0.37	0	5,5,5	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	910	-	-	0/4/4/4	0/0/0/0
2	GOL	A	911	-	-	0/4/4/4	0/0/0/0
3	TAR	A	912	-	-	0/4/12/12	0/0/0/0
3	TAR	B	901	-	-	0/4/12/12	0/0/0/0
2	GOL	B	903	-	-	0/4/4/4	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	912	TAR	O3-C3	-2.31	1.38	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	912	TAR	C4-C3-C2	-2.60	107.50	113.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	911	GOL	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	912	TAR	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	379/413 (91%)	-0.12	13 (3%) 46 38	27, 48, 108, 142	0
1	B	379/413 (91%)	-0.03	16 (4%) 37 29	33, 60, 107, 162	0
All	All	758/826 (91%)	-0.08	29 (3%) 41 33	27, 54, 108, 162	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	665	ILE	7.2
1	A	647	ASN	4.0
1	A	646	LEU	3.9
1	B	480	SER	3.8
1	A	813	ILE	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GOL	A	911	6/6	0.61	0.43	7.70	58,60,62,62	0
2	GOL	B	903	6/6	0.73	0.41	5.76	70,74,74,75	0
3	TAR	A	912	10/10	0.80	0.35	1.75	80,82,85,85	0
3	TAR	B	901	10/10	0.79	0.21	0.96	80,82,84,84	0
2	GOL	A	910	6/6	0.84	0.44	-	71,72,72,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.