



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 7, 2017 – 01:46 PM EST

PDB ID : 5TYI
Title : Grb7 SH2 with bicyclic peptide containing pY mimetic
Authors : Watson, G.M.; Wilce, M.C.J.; Wilce, J.A.
Deposited on : unknown
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1170 (2.16-2.16)
Clashscore	112137	1278 (2.16-2.16)
Ramachandran outliers	110173	1256 (2.16-2.16)
Sidechain outliers	110143	1255 (2.16-2.16)
RSRZ outliers	101464	1175 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	120	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 77%, yellow 77%, yellow 90%, green 90%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> .% 77% 9% 13% </div> </div>
1	B	120	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 3%, orange 3%, orange 79%, yellow 79%, yellow 88%, green 88%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 3% 79% 8% 13% </div> </div>
1	C	120	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 0%, green 77%, yellow 77%, yellow 90%, green 90%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 77% 10% 13% </div> </div>
1	D	120	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 73%, yellow 73%, yellow 85%, green 85%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> .% 73% 12% 15% </div> </div>
2	L	9	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 0%, green 89%, yellow 89%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 89% 11% </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	M	9	 100%
2	N	9	 78%22%
2	P	9	 67%33%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3651 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Growth factor receptor-bound protein 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	104	Total	C	N	O	S	0	0	0
			802	511	145	141	5			
1	B	105	Total	C	N	O	S	0	0	0
			817	522	148	142	5			
1	C	104	Total	C	N	O	S	0	0	0
			813	517	148	143	5			
1	D	102	Total	C	N	O	S	0	0	0
			789	503	142	139	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	413	GLY	-	expression tag	UNP Q14451
A	414	SER	-	expression tag	UNP Q14451
B	413	GLY	-	expression tag	UNP Q14451
B	414	SER	-	expression tag	UNP Q14451
C	413	GLY	-	expression tag	UNP Q14451
C	414	SER	-	expression tag	UNP Q14451
D	413	GLY	-	expression tag	UNP Q14451
D	414	SER	-	expression tag	UNP Q14451

- Molecule 2 is a protein called Peptide inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	9	Total	C	N	O	S	0	0	0
			77	48	12	16	1			
2	M	9	Total	C	N	O	S	0	0	0
			77	48	12	16	1			
2	N	9	Total	C	N	O	S	0	0	0
			77	48	12	16	1			
2	P	9	Total	C	N	O	S	0	0	0
			77	48	12	16	1			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	26	Total 26	O 26	0	0
3	B	24	Total 24	O 24	0	0
3	C	25	Total 25	O 25	0	0
3	D	23	Total 23	O 23	0	0
3	L	4	Total 4	O 4	0	0
3	M	6	Total 6	O 6	0	0
3	N	4	Total 4	O 4	0	0
3	P	10	Total 10	O 10	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Growth factor receptor-bound protein 7




- Molecule 1: Growth factor receptor-bound protein 7



- Molecule 1: Growth factor receptor-bound protein 7



- Molecule 1: Growth factor receptor-bound protein 7



- Molecule 2: Peptide inhibitor





- Molecule 2: Peptide inhibitor



There are no outlier residues recorded for this chain.

- Molecule 2: Peptide inhibitor

Chain N:  78% 22%



- Molecule 2: Peptide inhibitor

Chain P:  67% 33%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.07Å 107.61Å 48.01Å 90.00° 101.38° 90.00°	Depositor
Resolution (Å)	44.18 – 2.15 44.18 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.5 (44.18-2.15) 98.6 (44.18-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.207 , 0.246 0.207 , 0.221	Depositor DCC
R_{free} test set	1163 reflections (4.84%)	DCC
Wilson B-factor (Å ²)	22.1	Xtriage
Anisotropy	0.512	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 62.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3651	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.01 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.8522e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 48V, 99Y

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/817	0.64	0/1104
1	B	0.49	0/834	0.60	0/1127
1	C	0.54	0/829	0.62	0/1119
1	D	0.44	0/806	0.59	0/1090
2	L	0.60	0/52	0.61	0/65
2	M	0.44	0/52	0.52	0/65
2	N	0.43	0/52	0.51	0/65
2	P	0.43	0/52	0.53	0/65
All	All	0.48	0/3494	0.61	0/4700

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	802	0	759	7	0
1	B	817	0	778	10	0
1	C	813	0	775	10	0
1	D	789	0	735	11	0
2	L	77	0	46	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	M	77	0	46	0	0
2	N	77	0	47	3	0
2	P	77	0	47	3	0
3	A	26	0	0	0	0
3	B	24	0	0	0	0
3	C	25	0	0	0	0
3	D	23	0	0	1	0
3	L	4	0	0	0	0
3	M	6	0	0	0	0
3	N	4	0	0	0	0
3	P	10	0	0	0	0
All	All	3651	0	3233	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:438:ARG:NH2	2:N:4:GLY:O	2.17	0.78
1:B:426:HIS:NE2	1:B:504:ASP:HB2	2.02	0.74
1:D:487:GLU:O	1:D:489:GLY:N	2.22	0.72
1:C:487:GLU:HG3	1:C:487:GLU:O	1.94	0.68
1:C:432:PHE:O	1:C:529:ARG:HD3	1.94	0.68
1:D:478:LYS:NZ	3:D:601:HOH:O	2.29	0.66
1:B:426:HIS:CE1	1:B:504:ASP:CG	2.72	0.63
1:B:426:HIS:NE2	1:B:504:ASP:CB	2.62	0.62
1:B:426:HIS:CE1	1:B:506:LEU:HB3	2.40	0.57
1:A:516:ARG:HH21	1:A:520:PRO:HA	1.71	0.56
1:D:445:ILE:HG23	1:D:454:LEU:HD23	1.88	0.56
1:A:497:ASP:HB3	1:C:474:LEU:HD12	1.88	0.55
1:C:487:GLU:C	1:C:489:GLY:N	2.61	0.54
1:B:448:GLN:HB2	1:B:454:LEU:HD21	1.89	0.54
1:D:462:ARG:HH22	2:P:4:GLY:HA3	1.73	0.52
1:B:426:HIS:ND1	1:B:506:LEU:HB2	2.23	0.52
1:D:448:GLN:HB2	1:D:454:LEU:HD21	1.93	0.50
1:B:460:SER:HB3	1:B:463:ASN:O	2.12	0.49
1:C:429:GLN:O	1:C:529:ARG:NH1	2.45	0.49
1:D:506:LEU:O	1:D:510:GLU:HG3	2.12	0.49
1:C:445:ILE:HG23	1:C:454:LEU:HD23	1.95	0.48
1:A:438:ARG:NH1	2:L:4:GLY:O	2.47	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:426:HIS:CD2	1:B:504:ASP:HB2	2.50	0.46
1:B:425:ILE:O	1:B:428:THR:OG1	2.33	0.45
1:D:473:HIS:CE1	1:D:474:LEU:HG	2.54	0.43
1:C:496:ASP:HB2	2:N:9:48V:H8	1.99	0.43
1:D:497:ASP:OD1	2:P:9:48V:H9	2.19	0.43
1:A:486:GLU:HA	1:A:490:ARG:O	2.20	0.42
1:C:496:ASP:CB	2:N:9:48V:H8	2.49	0.42
1:A:441:SER:HB2	1:A:456:LEU:HD11	2.02	0.41
1:D:426:HIS:CG	1:D:505:LEU:HD12	2.55	0.41
1:D:462:ARG:HH21	2:P:5:99Y:C07	2.33	0.41
1:B:426:HIS:CE1	1:B:506:LEU:CB	3.03	0.41
1:A:485:SER:O	1:A:491:LEU:HA	2.21	0.40
1:D:426:HIS:CD2	1:D:505:LEU:HD12	2.57	0.40
1:A:432:PHE:CZ	1:A:434:GLY:HA2	2.57	0.40
1:C:506:LEU:O	1:C:510:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/120 (83%)	97 (97%)	3 (3%)	0	100	100
1	B	101/120 (84%)	100 (99%)	1 (1%)	0	100	100
1	C	100/120 (83%)	99 (99%)	1 (1%)	0	100	100
1	D	98/120 (82%)	95 (97%)	3 (3%)	0	100	100
2	L	3/9 (33%)	3 (100%)	0	0	100	100
2	M	3/9 (33%)	3 (100%)	0	0	100	100
2	N	3/9 (33%)	3 (100%)	0	0	100	100
2	P	3/9 (33%)	3 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	411/516 (80%)	403 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	82/106 (77%)	81 (99%)	1 (1%)	75	80
1	B	84/106 (79%)	83 (99%)	1 (1%)	75	80
1	C	85/106 (80%)	85 (100%)	0	100	100
1	D	81/106 (76%)	81 (100%)	0	100	100
2	L	4/6 (67%)	4 (100%)	0	100	100
2	M	4/6 (67%)	4 (100%)	0	100	100
2	N	4/6 (67%)	4 (100%)	0	100	100
2	P	4/6 (67%)	4 (100%)	0	100	100
All	All	348/448 (78%)	346 (99%)	2 (1%)	89	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	491	LEU
1	B	500	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	507	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	99Y	L	5	2	11,14,15	0.46	0	14,18,20	1.11	1 (7%)
2	48V	L	9	2	9,9,10	1.03	0	7,10,12	0.74	0
2	99Y	M	5	2	11,14,15	0.78	0	14,18,20	1.76	3 (21%)
2	48V	M	9	2	9,9,10	1.39	1 (11%)	7,10,12	0.58	0
2	99Y	N	5	2	11,14,15	0.60	0	14,18,20	1.47	3 (21%)
2	48V	N	9	2	9,9,10	2.13	2 (22%)	7,10,12	2.16	2 (28%)
2	99Y	P	5	2	11,14,15	0.50	0	14,18,20	1.78	4 (28%)
2	48V	P	9	2	9,9,10	0.78	0	7,10,12	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	99Y	L	5	2	-	0/4/10/12	0/1/1/1
2	48V	L	9	2	-	0/7/9/10	0/0/0/0
2	99Y	M	5	2	-	0/4/10/12	0/1/1/1
2	48V	M	9	2	-	0/7/9/10	0/0/0/0
2	99Y	N	5	2	-	0/4/10/12	0/1/1/1
2	48V	N	9	2	-	0/7/9/10	0/0/0/0
2	99Y	P	5	2	-	0/4/10/12	0/1/1/1
2	48V	P	9	2	-	0/7/9/10	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	N	9	48V	C06-C02	3.30	1.61	1.53
2	M	9	48V	C08-S07	3.61	1.84	1.81
2	N	9	48V	C08-S07	4.79	1.85	1.81

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	5	99Y	C06-C07-C08	-3.52	115.72	120.45
2	M	5	99Y	C03-C04-C12	-2.61	115.63	120.91
2	M	5	99Y	C11-C07-C08	-2.59	116.97	120.45
2	P	5	99Y	C03-C04-C05	-2.49	115.89	120.91
2	N	5	99Y	C11-C07-C08	-2.11	117.62	120.45
2	N	5	99Y	C03-C04-C12	-2.04	116.78	120.91
2	P	5	99Y	C04-C03-C02	2.54	119.42	114.29
2	L	5	99Y	C03-C02-C13	2.75	116.71	111.41
2	N	5	99Y	C04-C03-C02	3.13	120.61	114.29
2	P	5	99Y	C11-C07-C08	3.20	124.75	120.45
2	N	9	48V	C02-C06-S07	3.79	122.49	113.00
2	M	5	99Y	C04-C03-C02	3.85	122.07	114.29
2	N	9	48V	C06-C02-C03	4.13	118.83	109.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	N	9	48V	2	0
2	P	5	99Y	1	0
2	P	9	48V	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	104/120 (86%)	0.14	1 (0%) 82 86	14, 25, 47, 58	0
1	B	105/120 (87%)	0.20	4 (3%) 41 48	14, 26, 44, 49	0
1	C	104/120 (86%)	0.12	0 100 100	17, 27, 44, 57	0
1	D	102/120 (85%)	0.23	1 (0%) 82 86	18, 28, 46, 49	0
2	L	7/9 (77%)	0.15	0 100 100	20, 20, 29, 30	0
2	M	7/9 (77%)	-0.29	0 100 100	18, 22, 28, 29	0
2	N	7/9 (77%)	0.03	0 100 100	20, 23, 26, 30	0
2	P	7/9 (77%)	-0.37	0 100 100	20, 22, 24, 26	0
All	All	443/516 (85%)	0.15	6 (1%) 75 80	14, 26, 46, 58	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	423	ALA	3.1
1	B	424	ALA	2.6
1	B	425	ILE	2.5
1	D	464	PRO	2.2
1	B	426	HIS	2.0
1	A	526	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	48V	N	9	10/11	0.91	0.12	-	27,31,36,38	0
2	99Y	P	5	14/15	0.93	0.14	-	16,20,30,33	0
2	99Y	N	5	14/15	0.95	0.13	-	15,20,25,26	0
2	48V	M	9	10/11	0.91	0.12	-	23,28,34,35	0
2	99Y	M	5	14/15	0.95	0.11	-	17,20,27,28	0
2	99Y	L	5	14/15	0.95	0.12	-	15,17,21,23	0
2	48V	L	9	10/11	0.96	0.12	-	22,24,29,30	0
2	48V	P	9	10/11	0.96	0.14	-	18,23,24,24	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.