



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 03:00 pm GMT

PDB ID : 3TZN
Title : Crystal Structure of the Yersinia pestis Dihydropteroate synthase.
Authors : Wu, Y.
Deposited on : 2011-09-27
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4955 (2.10-2.06)
Clashscore	112137	5537 (2.10-2.06)
Ramachandran outliers	110173	5483 (2.10-2.06)
Sidechain outliers	110143	5484 (2.10-2.06)
RSRZ outliers	101464	4991 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 1%; height: 10px; background-color: red;"></div> <div style="width: 66%; height: 10px; background-color: green;"></div> <div style="width: 18%; height: 10px; background-color: yellow;"></div> <div style="width: 13%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%; text-align: center;"> 67% 18% 13% </div> </div> </div>
1	B	280	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 9%; height: 10px; background-color: red;"></div> <div style="width: 46%; height: 10px; background-color: green;"></div> <div style="width: 19%; height: 10px; background-color: yellow;"></div> <div style="width: 24%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%; text-align: center;"> 9% 55% 19% 24% </div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3561 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 7,8-dihydropteroate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	243	Total	C	N	O	S	0	1	0
			1860	1171	339	336	14			
1	B	212	Total	C	N	O	S	0	0	0
			1617	1023	296	285	13			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP Q7CKJ1
A	-1	SER	-	EXPRESSION TAG	UNP Q7CKJ1
A	0	HIS	-	EXPRESSION TAG	UNP Q7CKJ1
B	-2	GLY	-	EXPRESSION TAG	UNP Q7CKJ1
B	-1	SER	-	EXPRESSION TAG	UNP Q7CKJ1
B	0	HIS	-	EXPRESSION TAG	UNP Q7CKJ1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	63	Total	O	0	0
			63	63		
2	B	21	Total	O	0	0
			21	21		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 7,8-dihydropteroate synthase



• Molecule 1: 7,8-dihydropteroate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.43Å 49.94Å 75.51Å 90.00° 90.24° 90.00°	Depositor
Resolution (Å)	41.65 – 2.08 41.65 – 2.08	Depositor EDS
% Data completeness (in resolution range)	85.0 (41.65-2.08) 85.0 (41.65-2.08)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.68 (at 2.08Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, R_{free}	0.208 , 0.250 0.199 , 0.245	Depositor DCC
R_{free} test set	1386 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	45.2	Xtriage
Anisotropy	0.183	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.030 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3561	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	6/1894 (0.3%)	0.93	1/2563 (0.0%)
1	B	0.78	0/1640	0.77	0/2217
All	All	0.99	6/3534 (0.2%)	0.86	1/4780 (0.0%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	198	TYR	CE1-CZ	-8.36	1.27	1.38
1	A	263	VAL	CB-CG1	6.52	1.66	1.52
1	A	163	ARG	CG-CD	5.94	1.66	1.51
1	A	231	PRO	CA-C	5.45	1.63	1.52
1	A	259	VAL	CB-CG1	5.37	1.64	1.52
1	A	158	MET	CB-CG	5.05	1.67	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	48	LEU	CB-CG-CD1	-5.56	101.54	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1860	0	1885	28	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1617	0	1666	52	0
2	A	63	0	0	1	0
2	B	21	0	0	2	0
All	All	3561	0	3551	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:LEU:HD13	1:B:55:ILE:HD11	1.46	0.95
1:B:139:MET:HG2	1:B:140:HIS:H	1.34	0.91
1:B:230:VAL:CG2	1:B:234:GLN:HB2	2.03	0.88
1:B:225:GLY:CA	1:B:235:ARG:HE	1.89	0.86
1:B:222:SER:O	1:B:226:GLN:HG2	1.75	0.84
1:B:230:VAL:HG22	1:B:234:GLN:HB2	1.57	0.84
1:B:99:LYS:HB2	1:B:102:VAL:HG22	1.63	0.79
1:B:230:VAL:CG2	1:B:234:GLN:CB	2.61	0.78
1:B:14:ARG:HG3	1:B:14:ARG:HH11	1.49	0.77
1:B:48:LEU:CD1	1:B:55:ILE:HD11	2.16	0.73
1:A:120:LEU:HD11	1:A:136:VAL:HG11	1.71	0.72
1:A:38:ASP:O	1:A:42:GLN:HG3	1.89	0.72
1:B:193:ASN:OD1	1:B:196:HIS:ND1	2.24	0.70
1:B:212:LEU:O	2:B:280:HOH:O	2.13	0.65
1:B:14:ARG:CG	1:B:14:ARG:HH11	2.12	0.63
1:B:225:GLY:HA2	1:B:235:ARG:HE	1.63	0.62
1:A:186:PRO:HB2	1:A:200:LEU:HD21	1.82	0.62
1:B:139:MET:CG	1:B:140:HIS:H	2.11	0.61
1:B:3:LEU:HD11	1:B:183:LEU:HD11	1.83	0.60
1:B:139:MET:HE3	1:B:187:GLY:HA2	1.83	0.60
1:B:139:MET:CE	1:B:187:GLY:HA2	2.33	0.58
1:B:167:HIS:O	1:B:170:GLU:HB2	2.04	0.58
1:B:97:THR:OG1	1:B:102:VAL:HG21	2.04	0.58
1:B:230:VAL:CG2	1:B:234:GLN:HB3	2.34	0.58
1:A:83:GLU:O	1:A:87:GLN:HG3	2.04	0.57
1:A:78:VAL:O	1:A:81:VAL:HG13	2.05	0.57
1:B:139:MET:HG2	1:B:140:HIS:N	2.14	0.56
1:A:142:GLN:HE22	1:A:153:TYR:H	1.52	0.56
1:A:199:GLN:NE2	2:A:285:HOH:O	2.38	0.56
1:B:6:ARG:CZ	1:B:110:GLY:HA2	2.37	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:209:HIS:CE1	1:B:210:PHE:CE2	2.94	0.55
1:B:99:LYS:O	1:B:103:ILE:HG13	2.06	0.55
1:B:225:GLY:HA3	1:B:235:ARG:HE	1.72	0.55
1:A:119:SER:O	1:A:171:ARG:NH2	2.41	0.54
1:B:96:ASP:HA	1:B:115:ASN:O	2.07	0.54
1:B:230:VAL:HG21	1:B:234:GLN:HB3	1.88	0.54
1:A:22:ASN:OD1	1:A:58:GLY:O	2.26	0.53
1:A:275:ALA:HB1	1:B:199:GLN:HG2	1.90	0.53
1:B:12:LEU:HD13	1:B:213:PRO:HB3	1.90	0.52
1:B:114:ILE:CD1	1:B:134:LEU:HD12	2.41	0.51
1:B:114:ILE:HD11	1:B:134:LEU:HD12	1.91	0.51
1:B:157:LEU:O	1:B:161:ILE:HG12	2.10	0.51
1:A:96:ASP:HA	1:A:115:ASN:O	2.11	0.50
1:A:112:HIS:O	1:A:135:PRO:HD2	2.12	0.50
1:B:193:ASN:OD1	1:B:196:HIS:CG	2.64	0.49
1:B:81:VAL:HG12	1:B:85:LEU:HD12	1.94	0.49
1:B:209:HIS:CE1	1:B:210:PHE:CD2	3.01	0.49
1:B:231:PRO:HG2	1:B:234:GLN:OE1	2.13	0.49
1:A:128:ALA:O	1:A:132:THR:HG23	2.13	0.49
1:B:162:ASN:OD1	1:B:210:PHE:HE2	1.95	0.49
1:A:48:LEU:CD2	1:A:91:VAL:HG21	2.43	0.49
1:B:167:HIS:CD2	1:B:167:HIS:C	2.86	0.48
1:B:79:VAL:HG23	1:B:80:PRO:HD3	1.96	0.46
1:B:84:ALA:O	1:B:88:ARG:HG2	2.16	0.46
1:B:48:LEU:HD11	1:B:91:VAL:HG21	1.96	0.46
1:A:45:GLN:NE2	1:A:89:PHE:HE1	2.14	0.44
1:B:97:THR:OG1	1:B:102:VAL:CG2	2.65	0.44
1:A:-1:SER:HB3	1:A:1:MET:H	1.81	0.44
1:A:39:GLN:HA	1:A:42:GLN:HE21	1.82	0.44
1:B:209:HIS:ND1	1:B:210:PHE:N	2.66	0.43
1:B:54:LEU:N	1:B:54:LEU:HD12	2.33	0.43
1:A:219:SER:HB2	1:A:256:VAL:HA	2.00	0.42
1:B:100:ALA:HA	1:B:103:ILE:HD12	2.01	0.42
1:A:84:ALA:HA	1:A:88:ARG:HH21	1.85	0.42
1:A:139:MET:HA	1:A:185:ASP:O	2.20	0.42
1:A:160:ASP:OD1	1:A:163:ARG:NH2	2.52	0.42
1:B:237:ILE:HG12	2:B:290:HOH:O	2.19	0.42
1:B:16:GLN:HA	1:B:16:GLN:HE21	1.85	0.42
1:B:185:ASP:HA	1:B:215:LEU:O	2.20	0.41
1:A:179:LYS:HG2	1:A:212:LEU:HD21	2.01	0.41
1:B:225:GLY:HA3	1:B:235:ARG:HH11	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:ILE:HG13	1:A:118:ARG:N	2.34	0.41
1:A:231:PRO:HA	1:A:232:PRO:HD3	1.93	0.41
1:A:101:ALA:O	1:A:105:GLU:HG2	2.21	0.41
1:A:75:LEU:O	1:A:79:VAL:HB	2.21	0.41
1:A:162:ASN:HA	1:A:162:ASN:HD22	1.72	0.40
1:B:10:LEU:HB2	1:B:92:TRP:CE2	2.56	0.40
1:B:48:LEU:HD11	1:B:91:VAL:HG11	2.02	0.40
1:A:74:GLU:O	1:A:77:ARG:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/280 (84%)	232 (98%)	4 (2%)	0	100	100
1	B	202/280 (72%)	194 (96%)	8 (4%)	0	100	100
All	All	438/560 (78%)	426 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/227 (88%)	197 (99%)	2 (1%)	80	84

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	174/227 (77%)	164 (94%)	10 (6%)	24	20
All	All	373/454 (82%)	361 (97%)	12 (3%)	44	45

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-1	SER
1	A	81	VAL
1	B	14	ARG
1	B	16	GLN
1	B	36	ASN
1	B	79	VAL
1	B	115	ASN
1	B	116	ASP
1	B	117	ILE
1	B	170	GLU
1	B	233	GLN
1	B	258	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
1	A	39	GLN
1	A	42	GLN
1	A	45	GLN
1	A	142	GLN
1	A	162	ASN
1	A	249	GLN
1	B	16	GLN
1	B	39	GLN
1	B	42	GLN
1	B	43	HIS
1	B	115	ASN
1	B	167	HIS
1	B	233	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	243/280 (86%)	0.14	4 (1%) 72 75	30, 46, 85, 101	0
1	B	212/280 (75%)	0.75	26 (12%) 5 6	33, 82, 125, 139	0
All	All	455/560 (81%)	0.43	30 (6%) 19 23	30, 60, 118, 139	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	75	LEU	6.8
1	B	101	ALA	6.0
1	B	97	THR	5.9
1	B	169	ILE	5.2
1	B	172	CYS	5.2
1	B	161	ILE	4.5
1	A	78	VAL	3.6
1	B	177	ILE	3.6
1	B	102	VAL	3.5
1	A	76	ASP	3.3
1	B	165	PHE	3.3
1	A	59	GLY	3.2
1	B	196	HIS	3.2
1	B	168	HIS	3.1
1	B	77	ARG	3.0
1	B	8	LEU	3.0
1	B	7	GLY	2.9
1	B	57	ILE	2.7
1	B	133	GLY	2.6
1	B	164	PHE	2.6
1	B	167	HIS	2.5
1	B	190	PHE	2.5
1	B	137	CYS	2.5
1	B	38	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	187	GLY	2.3
1	B	175	ALA	2.3
1	B	6	ARG	2.2
1	B	157	LEU	2.2
1	B	98	SER	2.2
1	B	178	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.