



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 06:05 am GMT

PDB ID : 1U1Y
Title : Crystal structure of a complex between WT bacteriophage MS2 coat protein and an F5 aptamer RNA stemloop with 2aminopurine substituted at the-10 position
Authors : Horn, W.T.; Convery, M.A.; Stonehouse, N.J.; Adams, C.J.; Liljas, L.; Phillips, S.E.; Stockley, P.G.
Deposited on : 2004-07-16
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2749 (2.90-2.82)
Ramachandran outliers	110173	2687 (2.90-2.82)
Sidechain outliers	110143	2690 (2.90-2.82)
RNA backbone	2435	1033 (3.20-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	R	17	
1	S	17	
2	A	129	
2	B	129	
2	C	129	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3755 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*CP*CP*GP*GP*(2PR)P*GP*GP*AP*UP*CP*A P*CP*CP*AP*CP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	R	17	Total	C	N	O	P	0	0	0
			359	162	70	111	16			
1	S	16	Total	C	N	O	P	0	0	0
			345	154	67	108	16			

- Molecule 2 is a protein called Coat protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	129	Total	C	N	O	S	0	0	0
			964	606	165	189	4			
2	B	129	Total	C	N	O	S	0	0	0
			964	606	165	189	4			
2	C	129	Total	C	N	O	S	0	0	0
			964	606	165	189	4			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	38	Total	O	0	0
			38	38		
3	B	51	Total	O	0	0
			51	51		
3	C	43	Total	O	0	0
			43	43		
3	R	15	Total	O	0	0
			15	15		
3	S	12	Total	O	0	0
			12	12		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: 5'-R(*CP*CP*GP*GP*(2PR)P*GP*GP*AP*UP*CP*AP*CP*CP*AP*CP*GP*G)-3'




- Molecule 1: 5'-R(*CP*CP*GP*GP*(2PR)P*GP*GP*AP*UP*CP*AP*CP*CP*AP*CP*GP*G)-3'




- Molecule 2: Coat protein



- Molecule 2: Coat protein



- Molecule 2: Coat protein



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	288.61Å 288.61Å 654.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.85	Depositor
% Data completeness (in resolution range)	71.2 (30.00-2.85)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.188 , 0.196	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3755	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2PR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	R	0.28	0/376	0.82	1/581 (0.2%)
1	S	0.96	2/361 (0.6%)	2.14	7/559 (1.3%)
2	A	0.33	0/981	0.62	0/1337
2	B	0.33	0/981	0.64	0/1337
2	C	0.33	0/981	0.61	0/1337
All	All	0.43	2/3680 (0.1%)	0.93	8/5151 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	S	115	C	C4'-C3'	11.27	1.65	1.53
1	S	115	C	C3'-O3'	6.29	1.50	1.42

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	S	115	C	N1-C1'-C2'	-40.17	61.78	114.00
1	S	115	C	C1'-O4'-C4'	-8.31	103.25	109.90
1	S	115	C	C6-N1-C2	-7.63	117.25	120.30
1	S	115	C	O4'-C1'-C2'	7.17	114.05	107.60
1	S	115	C	C4'-C3'-O3'	6.85	126.70	113.00
1	S	115	C	C5'-C4'-O4'	-6.17	101.70	109.10
1	R	11	A	N9-C1'-C2'	5.50	121.15	114.00
1	S	115	C	O4'-C1'-N1	5.42	112.54	108.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	359	0	183	1	0
1	S	345	0	175	19	0
2	A	964	0	964	21	0
2	B	964	0	964	15	0
2	C	964	0	964	13	0
3	A	38	0	0	5	0
3	B	51	0	0	6	0
3	C	43	0	0	8	0
3	R	15	0	0	0	0
3	S	12	0	0	5	0
All	All	3755	0	3250	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:S:115:C:O2'	1:S:116:G:C5'	1.76	1.32
1:S:115:C:O3'	3:S:122:HOH:O	1.63	1.12
1:S:115:C:O2'	1:S:116:G:H5'	1.36	1.11
1:S:115:C:O2'	1:S:116:G:O5'	1.63	1.11
2:C:20:VAL:HG13	2:C:32:TRP:HB3	1.37	1.04
2:B:20:VAL:HG13	2:B:32:TRP:HB3	1.41	1.02
1:S:115:C:HO2'	1:S:116:G:C5'	1.65	1.00
2:A:20:VAL:HG13	2:A:32:TRP:HB3	1.53	0.88
1:S:115:C:C3'	3:S:122:HOH:O	2.24	0.80
1:S:111:A:H2'	3:S:119:HOH:O	1.95	0.65
2:C:20:VAL:CG1	2:C:32:TRP:HB3	2.23	0.62
1:S:115:C:HO2'	1:S:116:G:P	2.19	0.62
2:C:61:LYS:HG2	2:C:87:ASN:HD22	1.65	0.61
2:A:49:ARG:HH12	2:A:57:LYS:HE2	1.65	0.61
2:C:56:ARG:HD2	3:C:161:HOH:O	2.00	0.61
2:C:16:GLY:HA3	3:C:168:HOH:O	2.01	0.60
2:A:56:ARG:HD2	2:B:38:ARG:HD3	1.83	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:27:ASN:HB2	3:B:151:HOH:O	1.99	0.60
2:C:87:ASN:HB3	3:C:149:HOH:O	2.01	0.60
2:B:15:THR:HG23	3:B:166:HOH:O	2.02	0.60
2:A:57:LYS:HE3	2:A:89:GLU:CD	2.24	0.58
2:A:57:LYS:HD3	3:A:163:HOH:O	2.03	0.57
2:A:1:ALA:HB3	3:A:158:HOH:O	2.06	0.56
2:C:49:ARG:HD2	3:C:155:HOH:O	2.06	0.56
1:S:109:U:H5'	1:S:110:C:C5	2.43	0.54
2:A:52:SER:OG	2:A:53:ALA:N	2.41	0.53
1:S:108:A:H2'	1:S:110:C:C5	2.43	0.53
1:S:109:U:C5'	1:S:110:C:H5	2.20	0.53
2:A:33:ILE:HD13	2:A:38:ARG:NH1	2.24	0.53
2:B:20:VAL:HG13	2:B:32:TRP:CB	2.27	0.52
1:S:106:G:H5''	3:S:124:HOH:O	2.08	0.52
2:B:3:ASN:H	2:B:3:ASN:ND2	2.08	0.52
1:S:115:C:O2'	1:S:116:G:P	2.67	0.51
2:A:54:GLN:HB2	3:A:132:HOH:O	2.09	0.51
2:B:1:ALA:O	2:B:2:SER:CB	2.58	0.51
2:C:125:ASN:HA	3:C:158:HOH:O	2.10	0.50
2:A:86:LEU:HD21	2:A:88:MET:CE	2.42	0.50
2:C:56:ARG:NH1	3:C:166:HOH:O	2.44	0.50
2:B:83:ARG:HD2	3:B:165:HOH:O	2.10	0.50
2:A:49:ARG:NH1	2:A:57:LYS:HE2	2.27	0.49
2:A:33:ILE:HD13	2:A:38:ARG:HH11	1.78	0.48
2:B:56:ARG:NE	3:B:141:HOH:O	2.40	0.47
2:A:8:VAL:HA	2:A:19:THR:HA	1.95	0.47
1:S:104:G:C6	1:S:106:G:N7	2.83	0.47
2:A:57:LYS:HE3	2:A:89:GLU:OE1	2.16	0.46
2:B:67:VAL:O	2:B:70:GLN:HB3	2.17	0.45
2:C:57:LYS:NZ	3:C:164:HOH:O	2.49	0.45
1:S:112:C:O2'	1:S:113:C:H5'	2.16	0.45
2:A:103:LEU:HD12	2:A:103:LEU:HA	1.84	0.45
2:B:62:VAL:N	3:B:176:HOH:O	2.51	0.43
2:C:70:GLN:O	2:C:76:GLU:HA	2.19	0.43
1:S:109:U:H5'	1:S:110:C:H5	1.80	0.43
2:A:60:ILE:HB	2:A:88:MET:HG2	2.00	0.43
2:A:56:ARG:HD3	3:A:160:HOH:O	2.18	0.42
2:C:20:VAL:HG13	2:C:32:TRP:CB	2.28	0.42
2:A:61:LYS:HZ3	2:A:87:ASN:HD21	1.67	0.42
1:S:109:U:C5'	1:S:110:C:C5	3.00	0.42
2:B:28:GLY:HA3	3:C:157:HOH:O	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:61:LYS:NZ	2:A:87:ASN:HD21	2.19	0.41
2:B:87:ASN:HB2	3:B:164:HOH:O	2.19	0.41
2:B:30:ALA:O	2:B:45:THR:HA	2.20	0.41
1:R:16:G:O2'	1:R:17:G:H5'	2.20	0.41
1:S:112:C:H5'	3:S:121:HOH:O	2.20	0.41
2:C:15:THR:C	2:C:17:ASP:H	2.24	0.40
2:A:43:LYS:HE2	3:A:138:HOH:O	2.21	0.40
1:S:109:U:H5''	1:S:110:C:H5	1.87	0.40
2:B:118:ILE:HB	2:B:119:PRO:HD3	2.04	0.40
2:A:31:GLU:HG2	2:A:32:TRP:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	127/129 (98%)	120 (94%)	7 (6%)	0	100	100
2	B	127/129 (98%)	123 (97%)	3 (2%)	1 (1%)	22	52
2	C	127/129 (98%)	113 (89%)	14 (11%)	0	100	100
All	All	381/387 (98%)	356 (93%)	24 (6%)	1 (0%)	44	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	2	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	106/106 (100%)	98 (92%)	8 (8%)	16	39
2	B	106/106 (100%)	96 (91%)	10 (9%)	10	27
2	C	106/106 (100%)	95 (90%)	11 (10%)	8	22
All	All	318/318 (100%)	289 (91%)	29 (9%)	11	29

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	4	PHE
2	A	8	VAL
2	A	19	THR
2	A	20	VAL
2	A	25	PHE
2	A	48	VAL
2	A	52	SER
2	A	128	ILE
2	B	3	ASN
2	B	4	PHE
2	B	20	VAL
2	B	25	PHE
2	B	35	SER
2	B	38	ARG
2	B	48	VAL
2	B	56	ARG
2	B	77	LEU
2	B	114	ASP
2	C	15	THR
2	C	20	VAL
2	C	23	SER
2	C	35	SER
2	C	38	ARG
2	C	48	VAL
2	C	49	ARG
2	C	54	GLN
2	C	56	ARG
2	C	99	SER
2	C	102	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	87	ASN
2	B	3	ASN
2	B	36	ASN
2	B	70	GLN
2	C	6	GLN
2	C	87	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	R	14/17 (82%)	1 (7%)	0
1	S	14/17 (82%)	5 (35%)	0
All	All	28/34 (82%)	6 (21%)	0

All (6) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	R	6	G
1	S	106	G
1	S	109	U
1	S	112	C
1	S	115	C
1	S	116	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	2PR	R	5	1	16,23,24	0.64	0	17,33,36	1.40	3 (17%)
1	2PR	S	105	1	16,23,24	0.66	0	17,33,36	1.40	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	2PR	R	5	1	-	0/3/21/22	0/3/3/3
1	2PR	S	105	1	-	0/3/21/22	0/3/3/3

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	S	105	2PR	C2-N3-C4	-3.36	111.23	115.16
1	R	5	2PR	C2-N3-C4	-3.34	111.26	115.16
1	S	105	2PR	N2-C2-N1	-2.56	114.93	117.39
1	R	5	2PR	N2-C2-N1	-2.49	115.00	117.39
1	R	5	2PR	O4'-C1'-N9	-2.26	103.97	107.78
1	S	105	2PR	O3'-C3'-C2'	-2.06	103.33	110.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.