



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:58 am GMT

PDB ID : 3U2C  
Title : Aldose reductase in complex with NSAID-type inhibitor at 1.0 Å resolution  
Authors : Steuber, H.  
Deposited on : 2011-10-03  
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1400 (1.10-0.90)
Clashscore	112137	1005 (1.08-0.92)
Ramachandran outliers	110173	1411 (1.10-0.90)
Sidechain outliers	110143	1410 (1.10-0.90)
RSRZ outliers	101464	1410 (1.10-0.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PG5	A	1803	-	-	-	X
4	PG5	A	1805	-	-	X	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CIT	A	2000	-	-	-	X
6	SUZ	A	2001	-	-	-	X
7	PG4	A	2011	-	-	-	X

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 3040 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aldose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	315	2594	1672	432	477	13	0	14	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	4	ILE	LEU	CONFLICT	UNP P15121

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C<sub>21</sub>H<sub>28</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is 1-METHOXY-2-[2-(2-METHOXY-ETHOXY)]-ETHANE (three-letter code: PG5) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	8	4		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	12	8	4	0	0

- Molecule 5 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	13	6	7	0	0

- Molecule 6 is [(1Z)-5-FLUORO-2-METHYL-1-{4-[METHYLSULFINYL]BENZYLIDENE}-1H-INDEN-3-YL]ACETIC ACID (three-letter code: SUZ) (formula: C<sub>20</sub>H<sub>17</sub>FO<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	O	S		
6	A	1	25	20	1	3	1	0	0

- Molecule 7 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	A	1	13	8	5	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	309	Total	O	0	0
			309	309		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldose reductase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.09Å 67.21Å 49.42Å 90.00° 92.38° 90.00°	Depositor
Resolution (Å)	15.00 – 1.00 33.60 – 1.00	Depositor EDS
% Data completeness (in resolution range)	90.5 (15.00-1.00) 95.3 (33.60-1.00)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.69 (at 1.00Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.113 , 0.138 0.113 , 0.136	Depositor DCC
$R_{free}$ test set	7792 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	7.4	Xtrriage
Anisotropy	0.180	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 62.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.003 for l,k,-h 0.026 for h,-k,-l 0.017 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3040	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, PG4, PG5, CIT, PEG, SUZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.77	0/2698	1.31	42/3660 (1.1%)

There are no bond length outliers.

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	293	ARG	NE-CZ-NH1	20.21	130.41	120.30
1	A	284	ASP	CB-CG-OD2	13.69	130.62	118.30
1	A	284	ASP	CA-CB-CG	9.28	133.81	113.40
1	A	1	ALA	O-C-N	-9.19	107.99	122.70
1	A	306	HIS	CG-ND1-CE1	9.14	120.99	108.20
1	A	293	ARG	NE-CZ-NH2	-8.60	116.00	120.30
1	A	193	GLU	OE1-CD-OE2	8.39	133.37	123.30
1	A	3[A]	ARG	NE-CZ-NH1	8.32	124.46	120.30
1	A	3[B]	ARG	NE-CZ-NH1	8.32	124.46	120.30
1	A	115	PHE	CB-CG-CD2	-8.19	115.07	120.80
1	A	115	PHE	CB-CG-CD1	7.77	126.24	120.80
1	A	125	ASP	CB-CG-OD1	7.02	124.62	118.30
1	A	125	ASP	C-N-CA	6.87	138.88	121.70
1	A	1	ALA	CB-CA-C	6.86	120.38	110.10
1	A	134[A]	ASP	CB-CG-OD2	-6.68	112.29	118.30
1	A	134[B]	ASP	CB-CG-OD2	-6.68	112.29	118.30
1	A	2	SER	N-CA-CB	6.61	120.41	110.50
1	A	250	ARG	NE-CZ-NH1	-6.23	117.19	120.30
1	A	60[A]	GLU	OE1-CD-OE2	-6.18	115.88	123.30
1	A	60[B]	GLU	OE1-CD-OE2	-6.18	115.88	123.30
1	A	125	ASP	CA-C-N	6.11	130.63	117.20
1	A	307	LYS	C-N-CA	6.05	136.82	121.70
1	A	284	ASP	CB-CG-OD1	-5.73	113.14	118.30
1	A	1	ALA	CA-C-O	5.63	131.92	120.10

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	125	ASP	CA-C-O	-5.54	108.47	120.10
1	A	217	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	A	209	TYR	CB-CG-CD1	5.50	124.30	121.00
1	A	82	TYR	CB-CG-CD1	-5.46	117.73	121.00
1	A	134[A]	ASP	CB-CG-OD1	5.45	123.21	118.30
1	A	134[B]	ASP	CB-CG-OD1	5.45	123.21	118.30
1	A	129[A]	ASN	CB-CG-ND2	-5.42	103.68	116.70
1	A	129[B]	ASN	CB-CG-ND2	-5.42	103.68	116.70
1	A	273	PHE	CB-CG-CD1	5.41	124.59	120.80
1	A	307	LYS	O-C-N	-5.29	114.24	122.70
1	A	293	ARG	NH1-CZ-NH2	-5.28	113.59	119.40
1	A	129[A]	ASN	CA-C-O	5.25	131.13	120.10
1	A	129[B]	ASN	CA-C-O	5.25	131.13	120.10
1	A	209	TYR	CA-CB-CG	5.24	123.35	113.40
1	A	129[A]	ASN	CA-CB-CG	-5.23	101.90	113.40
1	A	129[B]	ASN	CA-CB-CG	-5.23	101.90	113.40
1	A	296	ARG	CG-CD-NE	5.21	122.75	111.80
1	A	82	TYR	CB-CG-CD2	5.04	124.02	121.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2594	0	2645	21	0
2	A	48	0	25	1	0
3	A	14	0	20	0	0
4	A	24	0	36	9	0
5	A	13	0	5	0	0
6	A	25	0	17	0	0
7	A	13	0	18	3	0
8	A	309	0	0	10	0
All	All	3040	0	2766	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:LYS:HE2	8:A:1288:HOH:O	1.55	1.05
1:A:52:ASN:HB2	4:A:1805:PG5:H61	1.39	1.01
4:A:1805:PG5:C8	8:A:1030:HOH:O	2.33	0.76
1:A:307:LYS:HE3	8:A:1062:HOH:O	1.89	0.71
4:A:1805:PG5:H83	8:A:1030:HOH:O	1.92	0.70
1:A:70:GLU:HG3	8:A:1330:HOH:O	1.94	0.68
4:A:1805:PG5:H72	8:A:1318:HOH:O	2.01	0.60
1:A:127:SER:HB3	1:A:129[A]:ASN:OD1	2.03	0.58
1:A:274:LYS:HE3	8:A:1199:HOH:O	2.04	0.57
1:A:126:GLU:HA	1:A:126:GLU:OE1	2.06	0.55
1:A:274:LYS:HD2	8:A:1279:HOH:O	2.07	0.55
1:A:185:GLU:OE2	1:A:187:HIS:HE1	1.92	0.53
1:A:52:ASN:HB2	4:A:1805:PG5:C6	2.28	0.51
1:A:232:ARG:HG2	4:A:1803:PG5:H32	1.92	0.51
4:A:1805:PG5:H71	8:A:1305:HOH:O	2.13	0.49
1:A:187:HIS:HD2	1:A:189:TYR:H	1.60	0.48
1:A:46:HIS:O	1:A:49[B]:GLN:NE2	2.48	0.46
1:A:20:TRP:CE3	1:A:21:LYS:HE3	2.50	0.45
1:A:262:LYS:O	2:A:400:NAP:H8A	2.17	0.44
1:A:98:ASP:OD1	4:A:1805:PG5:H62	2.19	0.43
4:A:1803:PG5:H52	4:A:1803:PG5:H31	1.80	0.43
1:A:125:ASP:OD1	1:A:129[A]:ASN:ND2	2.50	0.42
1:A:41:HIS:HA	1:A:73:PHE:O	2.20	0.42
7:A:2011:PG4:H82	8:A:1332:HOH:O	2.19	0.42
1:A:129[A]:ASN:H	1:A:129[A]:ASN:ND2	2.09	0.41
7:A:2011:PG4:H62	7:A:2011:PG4:H81	1.89	0.41
1:A:242:LYS:HE2	1:A:278:PHE:CD1	2.55	0.41
1:A:181:VAL:HA	1:A:205:VAL:O	2.20	0.41
7:A:2011:PG4:H41	7:A:2011:PG4:H62	1.75	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	327/316 (104%)	322 (98%)	5 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	294/281 (105%)	286 (97%)	8 (3%)	50 12

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	126	GLU
1	A	129[A]	ASN
1	A	129[B]	ASN
1	A	182	ASN
1	A	234	LYS
1	A	241	ASN
1	A	301	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	93	GLN
1	A	136	ASN
1	A	182	ASN
1	A	187	HIS
1	A	200	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	283	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PEG	A	1800	-	6,6,6	0.54	0	5,5,5	1.06	1 (20%)
4	PG5	A	1803	-	11,11,11	1.06	0	10,10,10	1.01	0
4	PG5	A	1805	-	11,11,11	0.46	0	10,10,10	0.24	0
5	CIT	A	2000	-	3,12,12	1.27	0	3,17,17	2.45	2 (66%)
6	SUZ	A	2001	-	24,27,27	1.61	4 (16%)	30,39,39	2.02	6 (20%)
3	PEG	A	2010	-	6,6,6	0.61	0	5,5,5	0.95	0
7	PG4	A	2011	-	12,12,12	0.46	0	11,11,11	0.29	0
2	NAP	A	400	-	44,52,52	1.23	6 (13%)	51,80,80	1.50	4 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	1800	-	-	0/4/4/4	0/0/0/0
4	PG5	A	1803	-	-	0/9/9/9	0/0/0/0
4	PG5	A	1805	-	-	0/9/9/9	0/0/0/0
5	CIT	A	2000	-	-	0/6/16/16	0/0/0/0
6	SUZ	A	2001	-	-	0/10/28/28	0/3/3/3
3	PEG	A	2010	-	-	0/4/4/4	0/0/0/0
7	PG4	A	2011	-	-	0/10/10/10	0/0/0/0
2	NAP	A	400	-	-	0/27/67/67	0/5/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	400	NAP	C2N-C3N	-3.87	1.33	1.39
2	A	400	NAP	C8A-N7A	-2.89	1.29	1.34
6	A	2001	SUZ	C11-C10	-2.66	1.48	1.51
6	A	2001	SUZ	C20-C2	2.01	1.41	1.38
2	A	400	NAP	C4N-C3N	2.12	1.42	1.39
2	A	400	NAP	C2A-N3A	2.14	1.35	1.32
2	A	400	NAP	P2B-O2B	2.31	1.63	1.59
6	A	2001	SUZ	C18-C7	2.61	1.50	1.45
2	A	400	NAP	C6N-N1N	2.79	1.42	1.35
6	A	2001	SUZ	C13-C10	4.17	1.51	1.44

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	2001	SUZ	C3-C4-C5	-4.08	115.93	121.28
6	A	2001	SUZ	C17-C18-C13	-3.49	117.21	121.09
2	A	400	NAP	N3A-C2A-N1A	-3.43	125.87	128.86
6	A	2001	SUZ	C14-C13-C10	-3.31	127.35	132.87
6	A	2001	SUZ	C20-C2-C3	-2.87	116.99	120.85
5	A	2000	CIT	C4-C3-C2	-2.73	102.93	109.75
2	A	400	NAP	C3N-C7N-N7N	-2.26	115.20	117.77
2	A	400	NAP	C2N-C3N-C7N	2.17	125.66	119.34
6	A	2001	SUZ	C3-C2-S	2.26	126.74	119.62
3	A	1800	PEG	O2-C2-C1	2.30	120.76	110.15
5	A	2000	CIT	C3-C2-C1	3.08	119.76	114.95
6	A	2001	SUZ	C4-C3-C2	6.02	124.85	119.42
2	A	400	NAP	C3N-C2N-N1N	6.91	127.38	120.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1803	PG5	2	0
4	A	1805	PG5	7	0
7	A	2011	PG4	3	0
2	A	400	NAP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	315/316 (99%)	-0.11	5 (1%) 72 63	4, 9, 27, 50	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	ALA	8.8
1	A	127	SER	4.2
1	A	128	GLY	3.0
1	A	222	PRO	2.9
1	A	126	GLU	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	PG5	A	1803	12/12	0.82	0.15	8.01	15,23,43,47	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	PG4	A	2011	13/13	0.83	0.19	7.12	23,36,48,57	0
4	PG5	A	1805	12/12	0.69	0.26	4.56	32,43,47,49	0
5	CIT	A	2000	13/13	0.85	0.14	3.72	19,29,47,58	0
6	SUZ	A	2001	25/25	0.94	0.12	3.18	10,13,44,69	0
3	PEG	A	1800	7/7	0.79	0.19	1.73	22,31,34,53	0
3	PEG	A	2010	7/7	0.50	0.23	1.19	46,56,61,67	0
2	NAP	A	400	48/48	0.99	0.05	-0.73	4,5,9,10	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.