



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 24, 2018 – 06:43 AM EST

PDB ID : 1UA2
Title : Crystal Structure of Human CDK7
Authors : Lolli, G.; Lowe, E.D.; Brown, N.R.; Johnson, L.N.
Deposited on : 2004-08-11
Resolution : 3.02 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1924 (3.04-3.00)
Clashscore	112137	2279 (3.04-3.00)
Ramachandran outliers	110173	2207 (3.04-3.00)
Sidechain outliers	110143	2210 (3.04-3.00)
RSRZ outliers	101464	1948 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	346	<div> <div>34%</div> <div>38%</div> <div>9%</div> <div>17%</div> </div>
1	B	346	<div> <div>29%</div> <div>37%</div> <div>15%</div> <div>17%</div> </div>
1	C	346	<div> <div>4%</div> <div>36%</div> <div>36%</div> <div>10%</div> <div>17%</div> </div>
1	D	346	<div> <div>2%</div> <div>35%</div> <div>35%</div> <div>12%</div> <div>17%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell division protein kinase 7.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	287	Total	C	N	O	P	S	0	0	0
			2291	1480	387	412	1	11			
1	B	287	Total	C	N	O	P	S	0	0	0
			2291	1480	387	412	1	11			
1	C	287	Total	C	N	O	P	S	0	0	0
			2291	1480	387	412	1	11			
1	D	287	Total	C	N	O	P	S	0	0	0
			2291	1480	387	412	1	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	170	TPO	THR	MODIFIED RESIDUE	UNP P50613
B	170	TPO	THR	MODIFIED RESIDUE	UNP P50613
C	170	TPO	THR	MODIFIED RESIDUE	UNP P50613
D	170	TPO	THR	MODIFIED RESIDUE	UNP P50613

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	16	Total	O	0	0
			16	16		
3	B	13	Total	O	0	0
			13	13		
3	C	12	Total	O	0	0
			12	12		
3	D	7	Total	O	0	0
			7	7		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cell division protein kinase 7




• Molecule 1: Cell division protein kinase 7




PRO
LYS
LYS
LEU
ILE
PHE

• Molecule 1: Cell division protein kinase 7



• Molecule 1: Cell division protein kinase 7



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.50 Å 191.63 Å 75.79 Å 90.00° 94.40° 90.00°	Depositor
Resolution (Å)	49.45 – 3.02 29.42 – 2.96	Depositor EDS
% Data completeness (in resolution range)	95.5 (49.45-3.02) 91.2 (29.42-2.96)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 2.95 Å)	Xtriage
Refinement program	REFMAC 5.2	Depositor
R, R_{free}	0.213 , 0.288 0.210 , 0.285	Depositor DCC
R_{free} test set	1742 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	52.7	Xtriage
Anisotropy	0.419	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 60.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	9336	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/2336	1.04	8/3169 (0.3%)
1	B	0.76	0/2336	1.05	16/3169 (0.5%)
1	C	0.76	0/2336	1.00	11/3169 (0.3%)
1	D	0.68	0/2336	0.99	11/3169 (0.3%)
All	All	0.75	0/9344	1.02	46/12676 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	3
1	C	0	2
All	All	0	9

There are no bond length outliers.

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	195	ASP	CB-CG-OD2	9.33	126.70	118.30
1	A	92	ASP	CB-CG-OD2	7.60	125.14	118.30
1	A	216	ASP	CB-CG-OD2	7.57	125.11	118.30
1	B	270	ASP	CB-CG-OD2	7.17	124.75	118.30
1	D	195	ASP	CB-CG-OD2	7.12	124.70	118.30

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	112	SER	Peptide
1	A	164	SER	Peptide
1	A	297	ASN	Peptide
1	A	304	GLY	Peptide
1	B	34	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2291	0	2300	182	0
1	B	2291	0	2300	169	0
1	C	2291	0	2300	134	0
1	D	2291	0	2300	138	0
2	A	31	0	12	2	0
2	B	31	0	12	3	0
2	C	31	0	12	1	0
2	D	31	0	12	0	0
3	A	16	0	0	9	0
3	B	13	0	0	5	0
3	C	12	0	0	1	0
3	D	7	0	0	2	0
All	All	9336	0	9248	625	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 625 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:GLU:OE2	1:A:238:PRO:HA	1.21	1.34
1:A:43:ILE:CD1	1:A:160:LYS:HE2	1.70	1.21
1:A:32:LYS:NZ	1:A:32:LYS:HB2	1.49	1.20
1:A:24:ALA:CB	1:A:160:LYS:NZ	2.06	1.18
1:A:103:LYS:O	1:A:105:ASN:N	1.75	1.17

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/346 (82%)	242 (86%)	22 (8%)	18 (6%)	1	8
1	B	282/346 (82%)	231 (82%)	26 (9%)	25 (9%)	1	4
1	C	282/346 (82%)	237 (84%)	30 (11%)	15 (5%)	2	13
1	D	282/346 (82%)	227 (80%)	37 (13%)	18 (6%)	1	8
All	All	1128/1384 (82%)	937 (83%)	115 (10%)	76 (7%)	1	7

5 of 76 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	ASP
1	A	22	GLN
1	A	84	LYS
1	A	104	ASP
1	A	112	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/298 (84%)	213 (86%)	36 (14%)	4	16
1	B	249/298 (84%)	200 (80%)	49 (20%)	1	8
1	C	249/298 (84%)	207 (83%)	42 (17%)	2	11
1	D	249/298 (84%)	197 (79%)	52 (21%)	1	6
All	All	996/1192 (84%)	817 (82%)	179 (18%)	2	10

5 of 179 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	251	SER
1	C	132	TRP
1	D	203	LEU
1	B	292	MET
1	C	40	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	131	HIS
1	C	22	GLN
1	D	135	HIS
1	B	141	ASN
1	B	142	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	A	170	1	9,10,11	2.93	4 (44%)	10,14,16	1.56	1 (10%)
1	TPO	B	170	1	9,10,11	3.03	5 (55%)	10,14,16	1.16	1 (10%)
1	TPO	C	170	1	9,10,11	3.18	5 (55%)	10,14,16	1.20	1 (10%)
1	TPO	D	170	1	9,10,11	3.20	5 (55%)	10,14,16	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	170	1	-	1/8/11/13	0/0/0/0
1	TPO	B	170	1	-	1/8/11/13	0/0/0/0
1	TPO	C	170	1	-	0/8/11/13	0/0/0/0
1	TPO	D	170	1	-	0/8/11/13	0/0/0/0

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	170	TPO	P-OG1	2.06	1.63	1.59
1	B	170	TPO	CA-C	2.13	1.53	1.50
1	B	170	TPO	P-OG1	2.33	1.63	1.59
1	D	170	TPO	P-OG1	2.50	1.63	1.59
1	A	170	TPO	P-OG1	2.67	1.64	1.59

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	170	TPO	C-CA-N	-2.35	105.11	109.86
1	B	170	TPO	C-CA-N	2.21	114.32	109.86
1	C	170	TPO	C-CA-N	2.71	115.32	109.86

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	170	TPO	OG1-CB-CA-N
1	B	170	TPO	OG1-CB-CA-N

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	170	TPO	2	0
1	B	170	TPO	1	0
1	C	170	TPO	1	0
1	D	170	TPO	3	0

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ATP	A	381	-	27,33,33	1.01	2 (7%)	25,52,52	1.58	4 (16%)
2	ATP	B	382	-	27,33,33	1.01	2 (7%)	25,52,52	1.74	4 (16%)
2	ATP	C	383	-	27,33,33	1.05	2 (7%)	25,52,52	1.72	4 (16%)
2	ATP	D	384	-	27,33,33	0.92	1 (3%)	25,52,52	1.91	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	381	-	-	0/18/38/38	0/3/3/3
2	ATP	B	382	-	-	0/18/38/38	0/3/3/3
2	ATP	C	383	-	-	0/18/38/38	0/3/3/3
2	ATP	D	384	-	-	0/18/38/38	0/3/3/3

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	381	ATP	C2-N3	2.02	1.35	1.32
2	C	383	ATP	O4'-C1'	2.08	1.44	1.41
2	B	382	ATP	O4'-C1'	2.22	1.44	1.41
2	B	382	ATP	C5-C4	2.76	1.46	1.40
2	A	381	ATP	C5-C4	2.98	1.47	1.40

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	384	ATP	N3-C2-N1	-7.10	122.67	128.86
2	C	383	ATP	N3-C2-N1	-5.57	124.01	128.86
2	A	381	ATP	N3-C2-N1	-4.99	124.52	128.86
2	B	382	ATP	N3-C2-N1	-4.70	124.76	128.86
2	B	382	ATP	C4-C5-N7	-4.16	105.39	109.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	381	ATP	2	0
2	B	382	ATP	3	0
2	C	383	ATP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/346 (82%)	-0.39	1 (0%) 93 82	18, 36, 68, 79	0
1	B	286/346 (82%)	-0.35	2 (0%) 87 67	16, 40, 74, 82	0
1	C	286/346 (82%)	-0.17	13 (4%) 34 14	20, 37, 78, 86	0
1	D	286/346 (82%)	-0.22	7 (2%) 59 29	23, 46, 79, 87	0
All	All	1144/1384 (82%)	-0.28	23 (2%) 65 35	16, 40, 74, 87	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	104	ASP	4.1
1	C	33	ASN	3.3
1	A	165	PRO	3.2
1	C	171	HIS	3.2
1	C	84	LYS	2.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	A	170	11/12	0.87	0.26	-	64,65,66,68	0
1	TPO	B	170	11/12	0.83	0.26	-	71,72,77,77	0
1	TPO	C	170	11/12	0.82	0.38	-	76,78,83,84	0
1	TPO	D	170	11/12	0.73	0.36	-	74,76,84,84	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ATP	D	384	31/31	0.95	0.21	-0.34	59,65,70,70	0
2	ATP	B	382	31/31	0.96	0.17	-0.67	50,57,67,68	0
2	ATP	A	381	31/31	0.96	0.12	-1.25	32,35,46,47	0
2	ATP	C	383	31/31	0.94	0.15	-1.43	54,59,66,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.