



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 9, 2018 – 02:21 AM EST

PDB ID : 5UA0  
Title : Dimeric crystal structure of HTPA reductase from arabidopsis thaliana  
Authors : Keown, J.K.; Pearce, F.G.; Goldstone, D.C.  
Deposited on : 2016-12-18  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030736
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	<div> <div>7%</div> <div> <div></div> <div>78%</div> <div>•</div> <div>18%</div> </div> </div>
1	B	329	<div> <div>2%</div> <div> <div></div> <div>77%</div> <div>5%</div> <div>18%</div> </div> </div>
1	C	329	<div> <div>6%</div> <div> <div></div> <div>76%</div> <div>6%</div> <div>•</div> <div>17%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6022 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-hydroxy-tetrahydrodipicolinate reductase 2, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	271	Total	C	N	O	S	0	0	0
			1994	1269	334	378	13			
1	B	270	Total	C	N	O	S	0	0	0
			1988	1262	328	384	14			
1	C	274	Total	C	N	O	S	0	0	0
			1991	1263	331	383	14			

There are 99 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-32	MET	-	initiating methionine	UNP Q8LB01
A	-31	HIS	-	expression tag	UNP Q8LB01
A	-30	HIS	-	expression tag	UNP Q8LB01
A	-29	HIS	-	expression tag	UNP Q8LB01
A	-28	HIS	-	expression tag	UNP Q8LB01
A	-27	HIS	-	expression tag	UNP Q8LB01
A	-26	HIS	-	expression tag	UNP Q8LB01
A	-25	GLY	-	expression tag	UNP Q8LB01
A	-24	LEU	-	expression tag	UNP Q8LB01
A	-23	PRO	-	expression tag	UNP Q8LB01
A	-22	ILE	-	expression tag	UNP Q8LB01
A	-21	PRO	-	expression tag	UNP Q8LB01
A	-20	ASN	-	expression tag	UNP Q8LB01
A	-19	PRO	-	expression tag	UNP Q8LB01
A	-18	LEU	-	expression tag	UNP Q8LB01
A	-17	LEU	-	expression tag	UNP Q8LB01
A	-16	GLY	-	expression tag	UNP Q8LB01
A	-15	LEU	-	expression tag	UNP Q8LB01
A	-14	ASP	-	expression tag	UNP Q8LB01
A	-13	SER	-	expression tag	UNP Q8LB01
A	-12	THR	-	expression tag	UNP Q8LB01
A	-11	GLU	-	expression tag	UNP Q8LB01
A	-10	ASN	-	expression tag	UNP Q8LB01

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	LEU	-	expression tag	UNP Q8LB01
A	-8	TYR	-	expression tag	UNP Q8LB01
A	-7	PHE	-	expression tag	UNP Q8LB01
A	-6	GLN	-	expression tag	UNP Q8LB01
A	-5	GLY	-	expression tag	UNP Q8LB01
A	-4	ILE	-	expression tag	UNP Q8LB01
A	-3	ASP	-	expression tag	UNP Q8LB01
A	-2	PRO	-	expression tag	UNP Q8LB01
A	-1	PHE	-	expression tag	UNP Q8LB01
A	0	THR	-	expression tag	UNP Q8LB01
B	-32	MET	-	initiating methionine	UNP Q8LB01
B	-31	HIS	-	expression tag	UNP Q8LB01
B	-30	HIS	-	expression tag	UNP Q8LB01
B	-29	HIS	-	expression tag	UNP Q8LB01
B	-28	HIS	-	expression tag	UNP Q8LB01
B	-27	HIS	-	expression tag	UNP Q8LB01
B	-26	HIS	-	expression tag	UNP Q8LB01
B	-25	GLY	-	expression tag	UNP Q8LB01
B	-24	LEU	-	expression tag	UNP Q8LB01
B	-23	PRO	-	expression tag	UNP Q8LB01
B	-22	ILE	-	expression tag	UNP Q8LB01
B	-21	PRO	-	expression tag	UNP Q8LB01
B	-20	ASN	-	expression tag	UNP Q8LB01
B	-19	PRO	-	expression tag	UNP Q8LB01
B	-18	LEU	-	expression tag	UNP Q8LB01
B	-17	LEU	-	expression tag	UNP Q8LB01
B	-16	GLY	-	expression tag	UNP Q8LB01
B	-15	LEU	-	expression tag	UNP Q8LB01
B	-14	ASP	-	expression tag	UNP Q8LB01
B	-13	SER	-	expression tag	UNP Q8LB01
B	-12	THR	-	expression tag	UNP Q8LB01
B	-11	GLU	-	expression tag	UNP Q8LB01
B	-10	ASN	-	expression tag	UNP Q8LB01
B	-9	LEU	-	expression tag	UNP Q8LB01
B	-8	TYR	-	expression tag	UNP Q8LB01
B	-7	PHE	-	expression tag	UNP Q8LB01
B	-6	GLN	-	expression tag	UNP Q8LB01
B	-5	GLY	-	expression tag	UNP Q8LB01
B	-4	ILE	-	expression tag	UNP Q8LB01
B	-3	ASP	-	expression tag	UNP Q8LB01
B	-2	PRO	-	expression tag	UNP Q8LB01
B	-1	PHE	-	expression tag	UNP Q8LB01

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	THR	-	expression tag	UNP Q8LB01
C	-32	MET	-	initiating methionine	UNP Q8LB01
C	-31	HIS	-	expression tag	UNP Q8LB01
C	-30	HIS	-	expression tag	UNP Q8LB01
C	-29	HIS	-	expression tag	UNP Q8LB01
C	-28	HIS	-	expression tag	UNP Q8LB01
C	-27	HIS	-	expression tag	UNP Q8LB01
C	-26	HIS	-	expression tag	UNP Q8LB01
C	-25	GLY	-	expression tag	UNP Q8LB01
C	-24	LEU	-	expression tag	UNP Q8LB01
C	-23	PRO	-	expression tag	UNP Q8LB01
C	-22	ILE	-	expression tag	UNP Q8LB01
C	-21	PRO	-	expression tag	UNP Q8LB01
C	-20	ASN	-	expression tag	UNP Q8LB01
C	-19	PRO	-	expression tag	UNP Q8LB01
C	-18	LEU	-	expression tag	UNP Q8LB01
C	-17	LEU	-	expression tag	UNP Q8LB01
C	-16	GLY	-	expression tag	UNP Q8LB01
C	-15	LEU	-	expression tag	UNP Q8LB01
C	-14	ASP	-	expression tag	UNP Q8LB01
C	-13	SER	-	expression tag	UNP Q8LB01
C	-12	THR	-	expression tag	UNP Q8LB01
C	-11	GLU	-	expression tag	UNP Q8LB01
C	-10	ASN	-	expression tag	UNP Q8LB01
C	-9	LEU	-	expression tag	UNP Q8LB01
C	-8	TYR	-	expression tag	UNP Q8LB01
C	-7	PHE	-	expression tag	UNP Q8LB01
C	-6	GLN	-	expression tag	UNP Q8LB01
C	-5	GLY	-	expression tag	UNP Q8LB01
C	-4	ILE	-	expression tag	UNP Q8LB01
C	-3	ASP	-	expression tag	UNP Q8LB01
C	-2	PRO	-	expression tag	UNP Q8LB01
C	-1	PHE	-	expression tag	UNP Q8LB01
C	0	THR	-	expression tag	UNP Q8LB01

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	10	Total	O	0	0
			10	10		
3	B	18	Total	O	0	0
			18	18		
3	C	11	Total	O	0	0
			11	11		



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.55Å 80.62Å 92.79Å 90.00° 95.84° 90.00°	Depositor
Resolution (Å)	61.95 – 2.30 54.05 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (61.95-2.30) 100.0 (54.05-2.30)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.46 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.226 , 0.258 0.228 , 0.255	Depositor DCC
$R_{free}$ test set	2012 reflections (5.20%)	DCC
Wilson B-factor (Å <sup>2</sup> )	50.0	Xtriage
Anisotropy	0.368	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 41.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6022	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/2030	0.72	3/2751 (0.1%)
1	B	0.59	0/2024	0.73	2/2745 (0.1%)
1	C	0.64	1/2026 (0.0%)	0.81	7/2752 (0.3%)
All	All	0.61	1/6080 (0.0%)	0.75	12/8248 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	79	GLU	CG-CD	5.12	1.59	1.51

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	296	ARG	NE-CZ-NH1	-8.67	115.96	120.30
1	C	296	ARG	NE-CZ-NH2	8.63	124.61	120.30
1	B	296	ARG	NE-CZ-NH1	7.41	124.01	120.30
1	B	296	ARG	NE-CZ-NH2	-7.24	116.68	120.30
1	C	78	ARG	NE-CZ-NH2	-7.20	116.70	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1994	0	1948	6	0
1	B	1988	0	1914	10	0
1	C	1991	0	1905	13	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	A	10	0	0	0	0
3	B	18	0	0	0	0
3	C	11	0	0	0	0
All	All	6022	0	5767	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:262:GLU:OE1	1:C:296:ARG:NH1	2.19	0.74
1:C:63:VAL:HG12	1:C:65:VAL:HG23	1.77	0.67
1:A:63:VAL:HG12	1:A:65:VAL:HG23	1.78	0.65
1:B:130:TYR:CE1	1:B:283:ILE:HD11	2.38	0.58
1:C:212:ILE:HD11	1:C:217:GLN:CG	2.39	0.53

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/329 (81%)	255 (96%)	11 (4%)	1 (0%)	38	47
1	B	266/329 (81%)	257 (97%)	9 (3%)	0	100	100
1	C	268/329 (82%)	257 (96%)	10 (4%)	1 (0%)	38	47
All	All	801/987 (81%)	769 (96%)	30 (4%)	2 (0%)	51	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	185	ASP
1	C	185	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/275 (75%)	201 (97%)	6 (3%)	48	64
1	B	207/275 (75%)	201 (97%)	6 (3%)	48	64
1	C	204/275 (74%)	198 (97%)	6 (3%)	48	64
All	All	618/825 (75%)	600 (97%)	18 (3%)	48	64

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	203	SER
1	B	234	PHE
1	C	148	LYS
1	B	30	SER
1	B	148	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	179	HIS
1	C	217	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	B	301	-	4,4,4	0.41	0	6,6,6	0.21	0
2	SO4	C	301	-	4,4,4	0.45	0	6,6,6	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	B	301	-	-	0/0/0/0	0/0/0/0
2	SO4	C	301	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	271/329 (82%)	0.50	22 (8%) 13 17	39, 59, 92, 116	0
1	B	270/329 (82%)	0.30	8 (2%) 51 58	40, 58, 83, 103	0
1	C	274/329 (83%)	0.60	20 (7%) 16 21	41, 62, 104, 131	0
All	All	815/987 (82%)	0.47	50 (6%) 22 28	39, 59, 94, 131	0

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	181	ALA	6.2
1	C	207	ASP	6.1
1	C	184	LEU	6.0
1	C	182	SER	5.1
1	A	187	SER	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	C	301	5/5	0.94	0.17	-0.30	65,70,72,74	0
2	SO4	B	301	5/5	0.94	0.12	-0.97	72,77,81,87	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.