



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 03:00 am GMT

PDB ID : 1UBT  
Title : Three-dimensional Structure of The Carbon Monoxide Complex of [NiFe]hydrogenase From Desulfovibrio vulgaris Miyazaki F  
Authors : Ogata, H.; Mizoguchi, Y.; Mizuno, N.; Miki, K.; Adachi, S.; Yasuoka, N.; Yagi, T.; Yamauchi, O.; Hirota, S.; Higuchi, Y.  
Deposited on : 2003-04-04  
Resolution : 1.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2078 (1.38-1.30)
Ramachandran outliers	110173	2021 (1.38-1.30)
Sidechain outliers	110143	2021 (1.38-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	S	267	
2	L	534	

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7233 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Periplasmic [NiFe] hydrogenase Small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	S	267	Total	C	N	O	S	0	0	0
			2019	1282	342	378	17			

- Molecule 2 is a protein called Periplasmic [NiFe] hydrogenase Large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	534	Total	C	N	O	S	0	0	0
			4177	2674	725	763	15			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	514	LYS	ASN	SEE REMARK 999	UNP P21852
L	515	LEU	VAL	SEE REMARK 999	UNP P21852

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	1	Total	Mg	0	0
			1	1		

- Molecule 4 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	S	1	Total	Fe	S	0	0
			8	4	4		
4	S	1	Total	Fe	S	0	0
			8	4	4		


- Molecule 5 is FE3-S4 CLUSTER (three-letter code: F3S) (formula:  $\text{Fe}_3\text{S}_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	S	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 6 is (MU-SULPHIDO)-BIS(MU-CYS,S)-[TRICARBONYLIRON-DI-(CYS,S)NIC

KEL(II)](FE-NI) (three-letter code: FNE) (formula:  $C_3FeNiO_3S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	L	1	Total	C	Fe	Ni	O	0	0
			8	3	1	1	3		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	L	657	Total	O	0	0
			657	657		
7	S	348	Total	O	0	0
			348	348		

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Periplasmic [NiFe] hydrogenase Small subunit



- Molecule 2: Periplasmic [NiFe] hydrogenase Large subunit



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.03Å 126.01Å 66.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.34	Depositor
% Data completeness (in resolution range)	91.6 (20.00-1.34)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.123 , 0.179	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7233	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, F3S, FNE, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	S	0.67	0/2075	1.38	18/2830 (0.6%)
2	L	0.64	1/4288 (0.0%)	1.28	35/5831 (0.6%)
All	All	0.65	1/6363 (0.0%)	1.31	53/8661 (0.6%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	552	HIS	C-O	5.94	1.34	1.23

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	S	6	ARG	CD-NE-CZ	26.75	161.06	123.60
1	S	100	ARG	NE-CZ-NH1	13.01	126.80	120.30
2	L	506	LEU	CA-CB-CG	12.30	143.59	115.30
1	S	100	ARG	CD-NE-CZ	11.86	140.21	123.60
2	L	340	ARG	CG-CD-NE	11.38	135.71	111.80

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	2019	0	1949	67	0
2	L	4177	0	4129	73	0
3	L	1	0	0	0	0
4	S	16	0	0	0	0
5	S	7	0	0	0	0
6	L	8	0	0	0	0
7	L	657	0	0	14	0
7	S	348	0	0	8	0
All	All	7233	0	6078	121	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 121 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:S:126:ASN:HD21	1:S:130:ALA:H	1.14	0.89
1:S:26:ARG:HH21	2:L:233:ASN:HD21	1.14	0.89
1:S:230:ASN:HB2	1:S:233:LYS:HE2	1.54	0.88
1:S:2:MET:HA	2:L:182:GLN:HG2	1.58	0.86
1:S:2:MET:HG2	2:L:182:GLN:HE21	1.38	0.86

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	265/267 (99%)	257 (97%)	7 (3%)	1 (0%)	38	12
2	L	532/534 (100%)	518 (97%)	13 (2%)	1 (0%)	51	20
All	All	797/801 (100%)	775 (97%)	20 (2%)	2 (0%)	44	17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	S	4	PRO
2	L	20	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	S	213/213 (100%)	192 (90%)	21 (10%)	9	0
2	L	438/438 (100%)	405 (92%)	33 (8%)	16	1
All	All	651/651 (100%)	597 (92%)	54 (8%)	13	0

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	43	VAL
2	L	161	LYS
2	L	445	LEU
2	L	49	LYS
2	L	132	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	75	HIS
2	L	132	HIS
2	L	468	GLN
2	L	113	ASN
2	L	188	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	FNE	L	1004	2	3,7,9	2.97	2 (66%)	0,9,15	0.00	-
4	SF4	S	1001	1	0,12,12	0.00	-	0,24,24	0.00	-
4	SF4	S	1002	1	0,12,12	0.00	-	0,24,24	0.00	-
5	F3S	S	1003	1	0,9,9	0.00	-	0,15,15	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	FNE	L	1004	2	-	0/0/9/18	0/0/0/1
4	SF4	S	1001	1	-	0/0/48/48	2/6/5/5
4	SF4	S	1002	1	-	0/0/48/48	2/6/5/5
5	F3S	S	1003	1	-	0/0/24/24	0/0/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	L	1004	FNE	O2-C2	-4.25	1.09	1.17
6	L	1004	FNE	O3-C3	-2.26	1.13	1.17

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

All (4) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	S	1001	SF4	FE1-FE2-S3-S4
4	S	1001	SF4	FE3-FE4-S1-S2
4	S	1002	SF4	FE1-FE2-S3-S4
4	S	1002	SF4	FE3-FE4-S1-S2

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.