



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 10:27 am GMT

PDB ID : 3UET
Title : Crystal structure of alpha-1,3/4-fucosidase from Bifidobacterium longum subsp. infantis D172A/E217A mutant complexed with lacto-N-fucopentaose II
Authors : Sakurama, H.; Fushinobu, S.; Yoshida, E.; Honda, Y.; Hidaka, M.; Ashida, H.; Kitaoka, M.; Katayama, T.; Yamamoto, K.; Kumagai, H.
Deposited on : 2011-10-31
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	
1	B	478	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	505	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-1,3/4-fucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total	C	N	O	S	0	0	0
			3569	2234	644	678	13			
1	B	456	Total	C	N	O	S	0	0	0
			3569	2234	644	678	13			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	172	ALA	ASP	ENGINEERED MUTATION	UNP B7GNN8
A	217	ALA	GLU	ENGINEERED MUTATION	UNP B7GNN8
B	172	ALA	ASP	ENGINEERED MUTATION	UNP B7GNN8
B	217	ALA	GLU	ENGINEERED MUTATION	UNP B7GNN8

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Na	0	0
			1	1		
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	3	Total	C	N	O	0	0
			36	20	1	15		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SUGAR (ALPHA-L-FUCOSE) (three-letter code: FUC) (formula: $C_6H_{12}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	2	Total	C	N	O	0	0
			26	14	1	11		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	218	Total	O	0	0
			218	218		
7	B	245	Total	O	0	0
			245	245		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-1,3/4-fucosidase



- Molecule 1: Alpha-1,3/4-fucosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.59Å 105.97Å 120.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.05 – 2.10 32.05 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.0 (32.05-2.10) 96.1 (32.05-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.173 , 0.223 0.173 , 0.223	Depositor DCC
R_{free} test set	3042 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	25.8	Xtriage
Anisotropy	0.139	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7683	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NA, GAL, NAG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.04	5/3656 (0.1%)	1.04	11/4974 (0.2%)
1	B	1.09	8/3656 (0.2%)	1.03	10/4974 (0.2%)
All	All	1.07	13/7312 (0.2%)	1.03	21/9948 (0.2%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	47	TRP	CD2-CE2	6.83	1.49	1.41
1	B	47	TRP	CG-CD1	6.59	1.46	1.36
1	A	135	TRP	CD2-CE2	6.38	1.49	1.41
1	A	47	TRP	CD2-CE2	6.00	1.48	1.41
1	B	290	TRP	CD2-CE2	5.96	1.48	1.41
1	B	106	TRP	CD2-CE2	5.83	1.48	1.41
1	A	290	TRP	CD2-CE2	5.60	1.48	1.41
1	A	277	TRP	CD2-CE2	5.56	1.48	1.41
1	B	312	SER	CB-OG	-5.37	1.35	1.42
1	B	226	TRP	CD2-CE2	5.08	1.47	1.41
1	A	170	TRP	CD2-CE2	5.06	1.47	1.41
1	B	188	TRP	CD2-CE2	5.01	1.47	1.41
1	B	310	TRP	CD2-CE2	5.00	1.47	1.41

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	351	ARG	NE-CZ-NH1	15.34	127.97	120.30
1	B	149	ASP	CB-CG-OD1	8.57	126.02	118.30
1	A	115	ARG	NE-CZ-NH1	8.24	124.42	120.30
1	B	222	ARG	NE-CZ-NH2	-8.07	116.26	120.30
1	A	351	ARG	NE-CZ-NH2	-7.10	116.75	120.30
1	B	362	ARG	NE-CZ-NH1	6.83	123.71	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	149	ASP	CB-CG-OD2	-6.71	112.27	118.30
1	B	115	ARG	NE-CZ-NH1	5.95	123.28	120.30
1	B	190	ARG	NE-CZ-NH2	-5.95	117.33	120.30
1	B	351	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	A	187	ASP	CB-CG-OD1	5.80	123.52	118.30
1	A	319	SER	CB-CA-C	5.74	121.00	110.10
1	A	422	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	B	123	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	B	311	LEU	CB-CG-CD1	-5.38	101.85	111.00
1	A	232	ARG	NE-CZ-NH2	-5.29	117.66	120.30
1	A	234	ARG	NE-CZ-NH1	-5.24	117.68	120.30
1	A	115	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	A	49	LEU	CB-CG-CD1	5.14	119.73	111.00
1	A	422	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	B	196	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3569	0	3429	30	0
1	B	3569	0	3430	26	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	36	0	33	0	0
4	A	4	0	6	0	0
4	B	4	0	6	0	0
5	B	10	0	10	1	0
6	B	26	0	24	1	0
7	A	218	0	0	4	0
7	B	245	0	0	3	0
All	All	7683	0	6938	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:ASN:HB3	7:A:653:HOH:O	1.71	0.89
1:B:362:ARG:HH11	1:B:362:ARG:HG3	1.35	0.88
1:B:232:ARG:HH22	1:B:254:THR:HG22	1.59	0.67
1:B:362:ARG:NH1	1:B:362:ARG:HG3	2.09	0.67
1:B:362:ARG:HH11	1:B:362:ARG:CG	2.09	0.66
1:A:82:THR:HA	1:A:131:TYR:HB3	1.79	0.65
1:B:232:ARG:NH2	1:B:254:THR:HG22	2.13	0.64
1:B:42:MET:HE3	7:B:586:HOH:O	1.97	0.63
1:B:95:ARG:HH22	1:B:162:GLN:HE22	1.45	0.63
1:A:183:THR:CG2	1:A:185:TYR:HE2	2.14	0.61
1:B:233:LEU:HD21	1:B:255:VAL:HG12	1.83	0.58
1:A:183:THR:HG22	1:A:185:TYR:CE2	2.37	0.58
1:A:30:GLU:OE1	1:A:351:ARG:NH2	2.29	0.58
1:A:183:THR:CG2	1:A:185:TYR:CE2	2.87	0.58
1:B:35:LEU:HD11	1:B:79:VAL:HG21	1.84	0.58
1:B:95:ARG:HH22	1:B:162:GLN:NE2	2.02	0.57
1:A:204:ILE:HB	1:A:209:PRO:HD2	1.86	0.56
1:A:351:ARG:NH2	7:A:578:HOH:O	2.37	0.56
1:A:224:ASN:OD1	1:A:439:GLY:HA3	2.06	0.55
1:A:406:ILE:C	1:A:406:ILE:HD12	2.27	0.55
1:A:183:THR:HG21	1:A:185:TYR:HE2	1.71	0.54
1:A:215:GLY:O	1:A:258:GLN:HG3	2.08	0.53
5:B:502:FUC:C1	6:B:503:NAG:C4	2.84	0.52
1:B:35:LEU:CD1	1:B:79:VAL:HG21	2.39	0.52
1:B:90:CYS:HB2	1:B:100:THR:HG22	1.93	0.51
1:B:353:ALA:O	1:B:357:VAL:HG22	2.10	0.51
1:B:353:ALA:HB1	1:B:476:VAL:CG2	2.41	0.50
1:A:472:ARG:HG3	1:A:473:ALA:N	2.27	0.48
1:B:476:VAL:HG22	1:B:477:ARG:N	2.29	0.47
1:A:37:PHE:CD1	1:A:326:PRO:HB2	2.50	0.46
1:B:180:ASN:ND2	1:B:182:LYS:HD3	2.31	0.46
1:B:421:LEU:HD11	1:B:459:LEU:HB3	1.97	0.46
1:B:95:ARG:NH2	7:B:603:HOH:O	2.49	0.45
1:A:67:TRP:O	1:A:71:LEU:HG	2.17	0.45
1:A:73:ALA:O	1:A:344:GLY:HA3	2.16	0.45
1:A:60:ARG:HG2	7:A:619:HOH:O	2.17	0.45
1:A:91:LEU:HD22	1:A:91:LEU:N	2.31	0.45
1:B:351:ARG:NH2	7:B:539:HOH:O	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:425:GLY:O	1:A:432:GLU:HA	2.17	0.45
1:A:358:ARG:O	1:A:398:PRO:HD3	2.17	0.44
1:B:170:TRP:CZ2	1:B:206:VAL:HG21	2.54	0.43
1:B:34:PHE:HB3	1:B:323:ASN:HB2	2.01	0.43
1:B:472:ARG:HG2	1:B:473:ALA:N	2.34	0.43
1:A:401:THR:O	1:A:453:GLU:HA	2.19	0.43
1:A:334:GLU:N	1:A:335:PRO:CD	2.83	0.42
1:B:354:LEU:HD12	1:B:354:LEU:HA	1.84	0.42
1:A:90:CYS:HB2	1:A:100:THR:HG22	2.02	0.41
1:A:177:GLU:OE1	1:A:182:LYS:N	2.49	0.41
1:B:235:SER:HA	1:B:238:LEU:HG	2.03	0.41
1:A:159:LEU:HA	1:A:159:LEU:HD23	1.91	0.41
1:B:327:SER:HB2	1:B:328:PRO:HD2	2.01	0.41
1:A:287:ARG:HD3	7:A:486:HOH:O	2.21	0.41
1:A:329:GLU:OE1	1:A:331:LEU:HD12	2.21	0.41
1:A:55:ALA:HA	1:A:105:PRO:HD3	2.02	0.40
1:A:46:GLU:O	1:A:47:TRP:HD1	2.03	0.40
1:A:34:PHE:HB3	1:A:323:ASN:HB2	2.04	0.40
1:B:9:ILE:HG23	1:B:9:ILE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	452/478 (95%)	434 (96%)	17 (4%)	1 (0%)	51	52
1	B	452/478 (95%)	437 (97%)	14 (3%)	1 (0%)	51	52
All	All	904/956 (95%)	871 (96%)	31 (3%)	2 (0%)	51	52

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	206	VAL
1	B	206	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/391 (96%)	360 (96%)	14 (4%)	39	39
1	B	374/391 (96%)	358 (96%)	16 (4%)	33	32
All	All	748/782 (96%)	718 (96%)	30 (4%)	36	36

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	PHE
1	A	49	LEU
1	A	96	LEU
1	A	182	LYS
1	A	256	SER
1	A	319	SER
1	A	336	ASP
1	A	358	ARG
1	A	362	ARG
1	A	422	ARG
1	A	431	THR
1	A	447	LEU
1	A	472	ARG
1	A	476	VAL
1	B	11	LEU
1	B	34	PHE
1	B	49	LEU
1	B	95	ARG
1	B	96	LEU
1	B	131	TYR
1	B	140	GLU
1	B	182	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	253	THR
1	B	260	ASP
1	B	336	ASP
1	B	357	VAL
1	B	362	ARG
1	B	398	PRO
1	B	427	LEU
1	B	459	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	10	ASN
1	B	162	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

5 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FUC	A	502	3	9,10,11	1.40	1 (11%)	13,14,16	1.33	2 (15%)
3	NAG	A	503	3	15,15,15	1.27	1 (6%)	21,21,21	1.85	6 (28%)
3	GAL	A	504	3	11,11,12	1.07	1 (9%)	13,15,17	1.54	3 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	B	503	5,6	15,15,15	1.05	1 (6%)	21,21,21	2.00	5 (23%)
6	GAL	B	504	6	11,11,12	0.80	1 (9%)	13,15,17	1.97	3 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FUC	A	502	3	-	0/0/17/20	0/1/1/1
3	NAG	A	503	3	-	0/6/26/26	0/1/1/1
3	GAL	A	504	3	-	0/2/19/22	0/1/1/1
6	NAG	B	503	5,6	-	0/6/26/26	0/1/1/1
6	GAL	B	504	6	-	0/2/19/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	504	GAL	O5-C1	-2.20	1.40	1.43
6	B	504	GAL	O5-C1	-2.20	1.40	1.43
6	B	503	NAG	C1-C2	2.80	1.56	1.52
3	A	502	FUC	C2-C3	3.48	1.57	1.52
3	A	503	NAG	C1-C2	3.49	1.57	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	504	GAL	C1-C2-C3	-4.24	104.27	109.65
6	B	504	GAL	C1-O5-C5	-3.91	106.78	112.17
3	A	504	GAL	O2-C2-C1	-3.86	101.33	109.18
6	B	503	NAG	C1-C2-C3	-3.38	105.94	110.54
6	B	503	NAG	O6-C6-C5	-3.34	100.11	111.34
3	A	503	NAG	O6-C6-C5	-3.28	100.29	111.34
3	A	503	NAG	C1-C2-C3	-2.52	107.11	110.54
6	B	503	NAG	O7-C7-C8	-2.43	117.64	122.06
3	A	503	NAG	O3-C3-C4	-2.07	105.86	110.36
3	A	504	GAL	C1-C2-C3	-2.06	107.04	109.65
6	B	504	GAL	O3-C3-C2	2.19	114.02	110.02
3	A	502	FUC	C1-O5-C5	2.28	117.44	112.39
3	A	504	GAL	O5-C1-C2	2.29	114.38	110.79
3	A	503	NAG	C2-N2-C7	2.29	129.00	123.19
3	A	502	FUC	O5-C5-C4	2.57	113.86	109.62

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	503	NAG	O3-C3-C2	3.13	115.97	109.61
6	B	503	NAG	C1-C2-N2	3.65	114.96	110.73
3	A	503	NAG	O1-C1-C2	4.67	118.93	109.22
6	B	503	NAG	O1-C1-C2	5.38	120.39	109.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	503	NAG	1	0

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	505	-	3,3,3	0.37	0	2,2,2	0.33	0
5	FUC	B	502	6	9,10,11	0.89	0	13,14,16	2.18	2 (15%)
4	EDO	B	505	-	3,3,3	0.23	0	2,2,2	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	505	-	-	0/1/1/1	0/0/0/0
5	FUC	B	502	6	-	0/0/17/20	0/1/1/1
4	EDO	B	505	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	502	FUC	O3-C3-C2	-2.27	105.90	110.02
5	B	502	FUC	C1-O5-C5	6.81	127.44	112.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	502	FUC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	456/478 (95%)	0.11	27 (5%) 23 29	18, 27, 53, 83	0
1	B	456/478 (95%)	0.01	16 (3%) 44 51	17, 26, 47, 98	0
All	All	912/956 (95%)	0.06	43 (4%) 32 38	17, 27, 50, 98	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	253	THR	10.9
1	A	178	GLY	5.3
1	B	181	GLY	5.1
1	A	399	GLN	5.0
1	A	254	THR	5.0
1	A	239	THR	4.9
1	B	269	ALA	4.3
1	A	398	PRO	4.2
1	A	179	LYS	4.2
1	A	428	PRO	4.0
1	B	182	LYS	3.9
1	B	178	GLY	3.9
1	B	179	LYS	3.8
1	A	185	TYR	3.7
1	A	430	GLY	3.7
1	B	254	THR	3.6
1	B	180	ASN	3.5
1	A	60	ARG	3.5
1	A	386	ALA	3.4
1	A	180	ASN	3.4
1	A	387	ASP	3.2
1	A	433	ARG	3.2
1	A	429	ASP	3.0
1	A	431	THR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	389	ALA	2.8
1	A	427	LEU	2.7
1	B	176	GLY	2.7
1	A	255	VAL	2.7
1	B	9	ILE	2.7
1	B	386	ALA	2.6
1	A	181	GLY	2.6
1	B	185	TYR	2.6
1	A	131	TYR	2.5
1	B	177	GLU	2.5
1	A	183	THR	2.4
1	A	368	ALA	2.3
1	A	132	LEU	2.3
1	A	401	THR	2.3
1	A	61	ASN	2.3
1	A	400	PRO	2.2
1	B	399	GLN	2.2
1	A	432	GLU	2.0
1	B	387	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	A	503	15/15	0.86	0.16	1.32	25,33,58,60	0
3	FUC	A	502	10/11	0.96	0.13	-0.29	20,25,28,29	0
6	GAL	B	504	11/12	0.96	0.10	-0.50	26,29,33,36	0
3	GAL	A	504	11/12	0.98	0.10	-1.18	22,25,28,29	0
6	NAG	B	503	15/15	0.89	0.17	-	27,37,52,58	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	EDO	A	505	4/4	0.96	0.14	2.94	33,36,40,42	0
4	EDO	B	505	4/4	0.95	0.12	0.21	31,32,33,36	0
2	NA	B	501	1/1	0.99	0.08	-0.57	26,26,26,26	0
5	FUC	B	502	10/11	0.97	0.11	-0.69	21,24,25,27	0
2	NA	A	501	1/1	0.98	0.07	-1.80	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.