



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 02:07 am GMT

PDB ID : 1UF3  
Title : Crystal structure of TT1561 of thermus thermophilus HB8  
Authors : Kato-Murayama, M.; Shirouzu, M.; Terada, T.; Murayama, K.; Kuramitsu, S.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2003-05-23  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	<div> <div>3%</div> <div>81% 18%</div> </div>
1	B	228	<div> <div>4%</div> <div>82% 15% ..</div> </div>
1	C	228	<div> <div>5%</div> <div>75% 23% ..</div> </div>
1	D	228	<div> <div>4%</div> <div>76% 24%</div> </div>
1	E	228	<div> <div>7%</div> <div>75% 23% .</div> </div>
1	F	228	<div> <div>2%</div> <div>81% 18% .</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	228	<div><div></div><div>4%</div><div>72%</div><div>26%</div><div></div></div>
1	H	228	<div><div></div><div>5%</div><div>75%</div><div>25%</div><div></div></div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 15287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hypothetical protein TT1561.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	228	Total	C	N	O	Se	0	0	0
			1815	1168	314	327	6			
1	B	226	Total	C	N	O	Se	0	0	0
			1796	1157	309	325	5			
1	C	225	Total	C	N	O	Se	0	0	0
			1785	1151	305	324	5			
1	D	228	Total	C	N	O	Se	0	0	0
			1815	1168	314	327	6			
1	E	228	Total	C	N	O	Se	0	0	0
			1815	1168	314	327	6			
1	F	228	Total	C	N	O	Se	0	0	0
			1815	1168	314	327	6			
1	G	228	Total	C	N	O	Se	0	0	0
			1815	1168	314	327	6			
1	H	228	Total	C	N	O	Se	0	0	0
			1815	1168	314	327	6			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	2	Total	Ca	0	0
			2	2		
2	D	2	Total	Ca	0	0
			2	2		
2	E	2	Total	Ca	0	0
			2	2		
2	H	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		
2	C	2	Total	Ca	0	0
			2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		
2	F	2	Total	Ca	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	124	Total	O	0	0
			124	124		
3	B	89	Total	O	0	0
			89	89		
3	C	83	Total	O	0	0
			83	83		
3	D	84	Total	O	0	0
			84	84		
3	E	82	Total	O	0	0
			82	82		
3	F	140	Total	O	0	0
			140	140		
3	G	102	Total	O	0	0
			102	102		
3	H	98	Total	O	0	0
			98	98		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: hypothetical protein TT1561




#### • Molecule 1: hypothetical protein TT1561




#### • Molecule 1: hypothetical protein TT1561






#### • Molecule 1: hypothetical protein TT1561









- Molecule 1: hypothetical protein TT1561




- Molecule 1: hypothetical protein TT1561



- Molecule 1: hypothetical protein TT1561



- Molecule 1: hypothetical protein TT1561



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	212.38Å 77.34Å 147.80Å 90.00° 127.21° 90.00°	Depositor
Resolution (Å)	29.97 – 2.10 29.96 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.2 (29.97-2.10) 98.0 (29.96-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.54 (at 2.10Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.195 , 0.245 0.196 , 0.244	Depositor DCC
$R_{free}$ test set	10967 reflections (10.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.3	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 51.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	15287	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.51	0/1861	0.75	2/2520 (0.1%)
1	B	0.44	0/1842	0.72	1/2496 (0.0%)
1	C	0.42	0/1831	0.68	0/2482
1	D	0.44	0/1861	0.72	0/2520
1	E	0.44	0/1861	0.72	2/2520 (0.1%)
1	F	0.51	0/1861	0.76	3/2520 (0.1%)
1	G	0.46	0/1861	0.72	2/2520 (0.1%)
1	H	0.48	0/1861	0.71	1/2520 (0.0%)
All	All	0.46	0/14839	0.72	11/20098 (0.1%)

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	27	LEU	CA-CB-CG	6.18	129.51	115.30
1	B	42	LEU	CA-CB-CG	6.17	129.50	115.30
1	F	154	LEU	CA-CB-CG	5.92	128.93	115.30
1	E	203	VAL	N-CA-C	-5.85	95.22	111.00
1	F	203	VAL	N-CA-C	-5.82	95.29	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1815	0	1796	32	0
1	B	1796	0	1771	25	0
1	C	1785	0	1758	41	0
1	D	1815	0	1796	37	0
1	E	1815	0	1796	46	0
1	F	1815	0	1796	36	0
1	G	1815	0	1796	53	0
1	H	1815	0	1796	45	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
2	G	2	0	0	0	0
2	H	1	0	0	0	0
3	A	124	0	0	2	0
3	B	89	0	0	3	0
3	C	83	0	0	2	0
3	D	84	0	0	3	0
3	E	82	0	0	4	0
3	F	140	0	0	4	0
3	G	102	0	0	4	0
3	H	98	0	0	1	0
All	All	15287	0	14305	298	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 298 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:59:LEU:HD22	1:E:94:MSE:HE3	1.45	0.97
1:F:1:MSE:HB2	1:F:105:ARG:HH21	1.32	0.95
1:A:215:LEU:HD22	1:A:222:LEU:HD22	1.51	0.92
1:C:219:ALA:HB1	1:C:221:LYS:HE2	1.48	0.92
1:E:161:HIS:HD2	1:E:196:LEU:HD13	1.35	0.90

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/228 (99%)	217 (96%)	9 (4%)	0	100	100
1	B	224/228 (98%)	211 (94%)	11 (5%)	2 (1%)	20	14
1	C	223/228 (98%)	210 (94%)	12 (5%)	1 (0%)	38	35
1	D	226/228 (99%)	217 (96%)	9 (4%)	0	100	100
1	E	226/228 (99%)	212 (94%)	14 (6%)	0	100	100
1	F	226/228 (99%)	217 (96%)	9 (4%)	0	100	100
1	G	226/228 (99%)	212 (94%)	14 (6%)	0	100	100
1	H	226/228 (99%)	213 (94%)	11 (5%)	2 (1%)	20	14
All	All	1803/1824 (99%)	1709 (95%)	89 (5%)	5 (0%)	44	44

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	162	LYS
1	H	165	ASN
1	C	28	ALA
1	B	163	GLY
1	B	29	PRO

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/180 (103%)	180 (97%)	6 (3%)	44	46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	184/180 (102%)	177 (96%)	7 (4%)	38	38
1	C	183/180 (102%)	178 (97%)	5 (3%)	50	54
1	D	186/180 (103%)	182 (98%)	4 (2%)	57	62
1	E	186/180 (103%)	177 (95%)	9 (5%)	30	27
1	F	186/180 (103%)	179 (96%)	7 (4%)	38	38
1	G	186/180 (103%)	180 (97%)	6 (3%)	44	46
1	H	186/180 (103%)	182 (98%)	4 (2%)	57	62
All	All	1483/1440 (103%)	1435 (97%)	48 (3%)	44	46

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	166	GLU
1	E	96	ASN
1	H	21	LEU
1	E	42	LEU
1	E	68	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	180	ASN
1	E	161	HIS
1	G	86	ASN
1	D	165	ASN
1	F	167	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	222/228 (97%)	-0.11	6 (2%)	55	61	16, 26, 38, 52	0
1	B	221/228 (96%)	0.04	8 (3%)	43	50	18, 28, 47, 62	0
1	C	220/228 (96%)	0.21	12 (5%)	26	32	22, 35, 50, 60	0
1	D	222/228 (97%)	0.14	8 (3%)	43	50	18, 31, 44, 65	0
1	E	222/228 (97%)	0.22	15 (6%)	18	23	21, 32, 49, 61	0
1	F	222/228 (97%)	-0.22	4 (1%)	69	73	15, 23, 37, 54	0
1	G	222/228 (97%)	0.02	8 (3%)	43	50	17, 30, 45, 64	0
1	H	222/228 (97%)	0.02	11 (4%)	30	36	16, 27, 43, 59	0
All	All	1773/1824 (97%)	0.04	72 (4%)	38	45	15, 29, 46, 65	0

The worst 5 of 72 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	228	ARG	7.1
1	H	164	LEU	5.4
1	B	164	LEU	5.4
1	E	163	GLY	4.6
1	E	228	ARG	4.5

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	CA	E	911	1/1	0.98	0.10	-0.42	45,45,45,45	0
2	CA	F	906	1/1	0.99	0.06	-0.94	26,26,26,26	0
2	CA	C	910	1/1	0.97	0.07	-1.37	51,51,51,51	0
2	CA	A	909	1/1	0.93	0.06	-1.75	45,45,45,45	0
2	CA	H	908	1/1	0.99	0.03	-1.83	32,32,32,32	0
2	CA	A	901	1/1	0.96	0.04	-2.01	29,29,29,29	0
2	CA	B	902	1/1	0.99	0.04	-2.14	32,32,32,32	0
2	CA	D	904	1/1	0.98	0.04	-2.18	35,35,35,35	0
2	CA	C	903	1/1	0.99	0.04	-2.48	39,39,39,39	0
2	CA	E	905	1/1	0.98	0.03	-2.63	32,32,32,32	0
2	CA	G	912	1/1	0.94	0.04	-2.77	45,45,45,45	0
2	CA	G	907	1/1	0.99	0.03	-2.81	34,34,34,34	0
2	CA	D	913	1/1	0.99	0.03	-	30,30,30,30	0
2	CA	F	914	1/1	0.99	0.03	-	31,31,31,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.