



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2017 – 08:38 PM EDT

PDB ID : 3UFQ
Title : Structure of rat nitric oxide synthase heme domain in complex with 6-(((3S,4S)-4-(((E)-5-(3-fluorophenyl)pent-4-en-1-yl)oxy)pyrrolidin-3-yl)methyl)-4-methylpyridin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : unknown
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2028 (2.08-2.04)
Clashscore	112137	2143 (2.08-2.04)
Ramachandran outliers	110173	2126 (2.08-2.04)
Sidechain outliers	110143	2126 (2.08-2.04)
RSRZ outliers	101464	2035 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>13%</div> <div> <div></div> <div>84%</div> <div>12%</div> <div>••</div> </div> </div>
1	B	422	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>8%</div> <div>•</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	A	860	-	-	-	X
5	HW5	A	800	-	-	-	X
5	HW5	B	800	-	-	-	X

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 7114 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	0	0
			3313	2121	566	605	21			
1	B	411	Total	C	N	O	S	0	1	0
			3348	2142	574	610	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 6-{{[(3S,4S)-4-{{[(4E)-5-(3-fluorophenyl)pent-4-en-1-yl]oxy}pyrrolidin-3-yl]methyl}-4-methylpyridin-2-amine (three-letter code: HW5) (formula: C₂₂H₂₈FN₃O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	F	N	O	0	0
			27	22	1	3	1		
5	B	1	Total	C	F	N	O	0	0
			27	22	1	3	1		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	99	Total	O	0	0
			99	99		
7	B	171	Total	O	0	0
			171	171		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.14Å 110.97Å 164.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.00 – 2.06 38.00 – 2.06	Depositor EDS
% Data completeness (in resolution range)	97.8 (38.00-2.06) 97.8 (38.00-2.06)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.06Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.190 , 0.231 0.210 , 0.244	Depositor DCC
R_{free} test set	2918 reflections (4.96%)	DCC
Wilson B-factor (Å ²)	38.0	Xtriage
Anisotropy	0.284	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7114	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, HW5, H4B, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/3406	0.69	0/4621
1	B	0.78	0/3444	0.73	0/4669
All	All	0.73	0/6850	0.71	0/9290

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3313	0	3221	36	0
1	B	3348	0	3264	19	0
2	A	43	0	30	7	0
2	B	43	0	30	5	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
5	A	27	0	28	2	0
5	B	27	0	28	2	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	99	0	0	4	0
7	B	171	0	0	1	0
All	All	7114	0	6637	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:701:THR:HB	7:A:8:HOH:O	1.48	1.07
1:A:535:GLN:HG2	7:A:77:HOH:O	1.61	0.99
1:A:336:MET:CE	1:A:678:TRP:HZ2	1.85	0.89
1:B:391:THR:O	1:B:392:SER:HB2	1.76	0.84
1:B:706:TYR:OH	2:B:750:HEM:O1D	2.04	0.76
1:A:507:GLN:O	1:A:507:GLN:HG2	1.90	0.71
1:A:631:VAL:HG11	1:B:628:GLN:HG3	1.76	0.67
1:A:336:MET:HE1	1:A:678:TRP:HZ2	1.61	0.66
1:A:336:MET:CE	1:A:678:TRP:CZ2	2.75	0.65
1:B:595:VAL:O	1:B:599:CYS:HB2	1.99	0.63
1:A:336:MET:HE1	1:A:678:TRP:CZ2	2.34	0.61
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.83	0.59
2:A:750:HEM:HBA1	5:A:800:HW5:H23	1.87	0.55
1:A:336:MET:HE3	1:A:678:TRP:HZ2	1.70	0.54
1:A:525:GLN:HE21	1:A:528:GLY:HA2	1.72	0.54
1:B:504:ILE:O	1:B:508:GLN:HG2	2.08	0.53
1:B:584:PHE:CD1	2:B:750:HEM:HAC	2.44	0.53
1:A:328:GLU:HB3	1:B:323:GLU:HG2	1.92	0.52
1:A:302:LYS:HD3	1:A:311:VAL:HG11	1.92	0.52
1:B:584:PHE:CD1	2:B:750:HEM:CAC	2.94	0.51
1:A:323:GLU:HG2	1:B:328:GLU:HB3	1.93	0.51
1:A:701:THR:CB	7:A:8:HOH:O	2.28	0.50
1:A:609:GLU:HG3	7:A:22:HOH:O	2.12	0.50
1:A:330:ILE:HD11	1:B:696:LEU:HB3	1.95	0.48
1:B:593:ILE:HA	1:B:597:ASP:HB2	1.95	0.48
1:B:592:GLU:OE1	5:B:800:HW5:H17	2.14	0.48
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.96	0.48
1:B:322:LEU:HB2	1:B:699:ARG:HE	1.78	0.47
1:A:706:TYR:OH	2:A:750:HEM:O2D	2.32	0.46
1:A:706:TYR:OH	2:A:750:HEM:CGD	2.64	0.46
1:B:485:TYR:CE1	1:B:514:ARG:HA	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:569:ASN:HD22	1:A:569:ASN:H	1.64	0.45
2:A:750:HEM:CMC	2:A:750:HEM:HBC2	2.46	0.45
1:B:499:VAL:O	1:B:503:GLU:HG3	2.17	0.45
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.53	0.44
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.06	0.43
1:A:336:MET:HG3	1:B:306:TRP:NE1	2.33	0.43
1:A:686:SER:HA	1:A:691:PHE:CG	2.53	0.42
2:A:750:HEM:HBB2	2:A:750:HEM:HHC	2.01	0.42
1:A:517:PHE:HB2	1:A:560:LYS:HE3	2.02	0.42
1:A:571:LEU:C	1:A:571:LEU:HD23	2.40	0.42
1:A:360:GLU:OE1	1:A:701:THR:CG2	2.68	0.42
1:A:332:MET:CE	1:B:301:LEU:HD13	2.50	0.41
1:A:588:TYR:CD1	1:A:593:ILE:HD11	2.55	0.41
2:A:750:HEM:HMC1	2:A:750:HEM:HBC2	2.03	0.41
1:A:675:ASP:O	1:A:679:ILE:HG12	2.21	0.41
2:B:750:HEM:HHC	2:B:750:HEM:HBB2	2.03	0.41
1:A:592:GLU:OE1	5:A:800:HW5:N01	2.54	0.41
1:A:706:TYR:OH	2:A:750:HEM:O1D	2.38	0.41
2:B:750:HEM:HBA2	5:B:800:HW5:H23	2.02	0.41
1:A:299:ARG:O	1:A:317:HIS:CE1	2.74	0.41
1:A:306:TRP:CD2	1:B:336:MET:HE3	2.56	0.41
1:A:478:GLN:HA	1:A:566:ALA:O	2.21	0.40
1:A:307:GLU:HG3	7:B:192:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/422 (96%)	391 (97%)	11 (3%)	1 (0%)	51	42
1	B	408/422 (97%)	402 (98%)	6 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	811/844 (96%)	793 (98%)	17 (2%)	1 (0%)	55 48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	489	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	363/377 (96%)	348 (96%)	15 (4%)	35 27
1	B	367/377 (97%)	357 (97%)	10 (3%)	50 44
All	All	730/754 (97%)	705 (97%)	25 (3%)	42 35

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	337	LEU
1	A	350	THR
1	A	370	LYS
1	A	380	ARG
1	A	507	GLN
1	A	508	GLN
1	A	523	LEU
1	A	527	ASN
1	A	547	ARG
1	A	552	ASP
1	A	569	ASN
1	A	601	ASN
1	A	620	LYS
1	A	645	LYS
1	A	653	SER
1	B	332	MET
1	B	353	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	389	GLU
1	B	390	SER
1	B	392	SER
1	B	454	ASN
1	B	507	GLN
1	B	532	GLU
1	B	547	ARG
1	B	653	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	454	ASN
1	A	569	ASN
1	A	601	ASN
1	A	605	ASN
1	A	697	ASN
1	B	353	GLN
1	B	364	GLN
1	B	385	ASN
1	B	454	ASN
1	B	507	GLN
1	B	535	GLN
1	B	601	ASN
1	B	605	ASN
1	B	697	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	750	1	28,50,50	2.43	10 (35%)	17,82,82	1.86	5 (29%)
3	H4B	A	760	-	14,18,18	0.96	1 (7%)	12,26,26	2.42	5 (41%)
5	HW5	A	800	-	26,29,29	0.79	0	32,38,38	1.76	6 (18%)
4	ACT	A	860	-	1,3,3	2.06	1 (100%)	0,3,3	0.00	-
2	HEM	B	750	1	28,50,50	2.47	11 (39%)	17,82,82	1.58	3 (17%)
3	H4B	B	760	-	14,18,18	1.01	1 (7%)	12,26,26	2.66	6 (50%)
5	HW5	B	800	-	26,29,29	0.79	1 (3%)	32,38,38	1.86	7 (21%)
4	ACT	B	860	-	1,3,3	1.53	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/6/54/54	0/0/8/8
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
5	HW5	A	800	-	-	0/13/23/23	0/3/3/3
4	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/6/54/54	0/0/8/8
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
5	HW5	B	800	-	-	0/13/23/23	0/3/3/3
4	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	C3B-C2B	-5.48	1.33	1.40
2	B	750	HEM	C3C-C2C	-5.45	1.33	1.40
2	A	750	HEM	C3B-C2B	-5.21	1.33	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C3C-C2C	-4.27	1.34	1.40
3	B	760	H4B	C4-C4A	-2.31	1.38	1.41
3	A	760	H4B	C4-C4A	-2.31	1.38	1.41
2	A	750	HEM	CMC-C2C	2.02	1.55	1.51
5	B	800	HW5	C24-C23	2.04	1.41	1.37
4	A	860	ACT	CH3-C	2.06	1.51	1.48
2	A	750	HEM	CMD-C2D	2.09	1.55	1.51
2	B	750	HEM	CMC-C2C	2.24	1.56	1.51
2	A	750	HEM	CMB-C2B	2.38	1.56	1.51
2	B	750	HEM	C1B-NB	2.43	1.39	1.36
2	B	750	HEM	CMD-C2D	2.45	1.56	1.51
2	B	750	HEM	CMB-C2B	2.48	1.56	1.51
2	A	750	HEM	CAD-C3D	2.50	1.57	1.52
2	B	750	HEM	CAD-C3D	2.80	1.57	1.52
2	B	750	HEM	C4D-ND	3.11	1.40	1.36
2	B	750	HEM	C3B-CAB	3.26	1.54	1.47
2	A	750	HEM	C3C-CAC	3.43	1.54	1.47
2	A	750	HEM	C3B-CAB	3.73	1.55	1.47
2	B	750	HEM	C3C-CAC	3.79	1.55	1.47
2	B	750	HEM	C3D-C2D	4.66	1.51	1.37
2	A	750	HEM	C3D-C2D	4.79	1.51	1.37
2	A	750	HEM	C4D-ND	5.01	1.42	1.36

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	800	HW5	C12-C13-C14	-4.07	117.47	125.50
5	A	800	HW5	C05-C06-N01	-3.96	118.62	122.91
5	A	800	HW5	C12-C13-C14	-3.80	117.99	125.50
2	A	750	HEM	CBA-CAA-C2A	-3.64	105.52	112.48
5	B	800	HW5	C05-C06-N01	-3.43	119.19	122.91
2	A	750	HEM	CAD-CBD-CGD	-3.33	106.97	112.66
3	A	760	H4B	N3-C2-N1	-3.19	120.27	125.45
2	B	750	HEM	CBD-CAD-C3D	-3.14	106.47	112.47
2	B	750	HEM	C1D-C2D-C3D	-3.00	104.91	107.00
5	A	800	HW5	C24-C23-C22	-2.89	119.50	123.29
2	B	750	HEM	CBA-CAA-C2A	-2.79	107.15	112.48
5	B	800	HW5	C24-C23-C22	-2.69	119.76	123.29
3	B	760	H4B	N3-C2-N1	-2.54	121.34	125.45
2	A	750	HEM	CBD-CAD-C3D	-2.49	107.72	112.47
5	B	800	HW5	C21-C14-C13	-2.48	120.21	127.00
3	B	760	H4B	C4A-C4-N3	-2.08	117.81	123.91

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	HEM	C3C-C4C-NC	-2.05	107.06	110.94
3	B	760	H4B	C2-N1-C8A	2.24	119.56	114.51
5	B	800	HW5	C08-C06-N01	2.28	120.72	117.12
3	B	760	H4B	N2-C2-N3	2.37	121.04	117.24
3	A	760	H4B	C2-N1-C8A	2.56	120.27	114.51
5	A	800	HW5	C2'-N1'-C5'	3.10	112.62	105.40
5	A	800	HW5	C08-C06-N01	3.27	122.27	117.12
3	B	760	H4B	C4-N3-C2	3.44	121.01	116.06
3	A	760	H4B	N2-C2-N3	3.66	123.09	117.24
3	A	760	H4B	C4-N3-C2	3.71	121.39	116.06
5	B	800	HW5	C2'-N1'-C5'	3.71	114.03	105.40
2	A	750	HEM	C4C-C3C-C2C	3.80	109.55	106.90
3	A	760	H4B	C4-C4A-C8A	4.06	118.24	114.56
5	A	800	HW5	C02-N01-C06	4.28	121.20	118.17
5	B	800	HW5	C02-N01-C06	5.60	122.13	118.17
3	B	760	H4B	C4-C4A-C8A	6.88	120.79	114.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	7	0
5	A	800	HW5	2	0
2	B	750	HEM	5	0
5	B	800	HW5	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/422 (96%)	0.57	54 (13%) 4 3	28, 52, 89, 112	0
1	B	411/422 (97%)	0.13	20 (4%) 30 31	26, 41, 66, 90	0
All	All	818/844 (96%)	0.35	74 (9%) 10 10	26, 46, 84, 112	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	348	VAL	8.6
1	B	300	PHE	6.6
1	B	350	THR	6.4
1	B	718	GLY	6.2
1	A	488	PRO	5.7
1	A	716	TRP	5.6
1	A	351	LYS	4.9
1	A	713	THR	4.7
1	A	715	VAL	4.6
1	A	350	THR	4.6
1	A	352	ASP	4.5
1	A	299	ARG	4.5
1	A	506	ILE	4.4
1	B	619	ARG	4.4
1	B	352	ASP	3.9
1	B	338	PRO	3.8
1	A	489	ASP	3.8
1	A	388	ILE	3.7
1	A	385	ASN	3.7
1	A	486	LYS	3.6
1	A	503	GLU	3.4
1	A	712	ASN	3.4
1	A	389	GLU	3.2
1	A	392	SER	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	391	THR	3.1
1	A	507	GLN	3.1
1	A	300	PHE	3.0
1	B	351	LYS	3.0
1	A	552	ASP	3.0
1	A	355	PHE	2.9
1	B	349	ARG	2.9
1	A	470	HIS	2.9
1	A	390	SER	2.9
1	A	678	TRP	2.8
1	A	714	HIS	2.7
1	A	711	TRP	2.7
1	B	299	ARG	2.7
1	A	386	LYS	2.7
1	A	487	GLN	2.7
1	B	677	VAL	2.7
1	A	591	THR	2.6
1	A	415	CYS	2.6
1	A	619	ARG	2.6
1	B	591	THR	2.6
1	A	677	VAL	2.6
1	A	490	GLY	2.6
1	A	354	LEU	2.6
1	A	685	GLY	2.6
1	A	584	PHE	2.6
1	A	567	VAL	2.5
1	A	593	ILE	2.5
1	A	679	ILE	2.4
1	A	681	PRO	2.4
1	A	491	SER	2.4
1	B	681	PRO	2.4
1	A	480	ILE	2.3
1	B	667	ARG	2.3
1	A	479	LEU	2.3
1	B	479	LEU	2.3
1	B	321	THR	2.3
1	A	469	LYS	2.3
1	B	678	TRP	2.3
1	A	514	ARG	2.3
1	B	620	LYS	2.2
1	A	321	THR	2.2
1	A	393	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	328	GLU	2.2
1	A	466	THR	2.2
1	A	682	PRO	2.2
1	A	338	PRO	2.1
1	B	389	GLU	2.1
1	A	382	GLU	2.0
1	B	561	TRP	2.0
1	A	416	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ACT	A	860	4/4	0.91	0.23	9.25	63,64,65,65	0
5	HW5	A	800	27/27	0.84	0.37	3.80	30,50,96,96	0
5	HW5	B	800	27/27	0.88	0.27	2.31	31,50,90,90	0
4	ACT	B	860	4/4	0.93	0.14	1.75	53,54,55,55	0
3	H4B	A	760	17/17	0.96	0.20	0.75	31,33,39,42	0
2	HEM	A	750	43/43	0.97	0.21	0.74	29,34,45,49	0
2	HEM	B	750	43/43	0.98	0.17	0.40	20,32,43,48	0
3	H4B	B	760	17/17	0.97	0.17	0.18	28,32,38,41	0
6	ZN	A	900	1/1	1.00	0.08	-0.98	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.