



# Full wwPDB/EMDatabank EM Map/Model Validation Report ⓘ

Aug 17, 2017 – 04:25 PM EDT

PDB ID : 4UF8  
EMDB ID: : EMD-2936  
Title : Electron cryo-microscopy structure of PB1-p62 filaments  
Authors : Ciuffa, R.; Lamark, T.; Tarafder, A.; Guesdon, A.; Rybina, S.; Hagen, W.J.H.;  
Johansen, T.; Sachse, C.  
Deposited on : unknown  
Resolution : 10.90 Å(reported)  
Based on PDB ID : 2KKC

This is a Full wwPDB/EMDatabank EM Map/Model Validation Report  
for a publicly released PDB/EMDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

MolProbity : 4.02b-467  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 10.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	125131	1336
Ramachandran outliers	121729	1120
Sidechain outliers	121581	1026

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	100	
1	B	100	
1	C	100	
1	I	100	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3120 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called SEQUESTOSOME-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	100	Total	C	N	O	S	0	0
			780	492	132	151	5		
1	B	100	Total	C	N	O	S	0	0
			780	492	132	151	5		
1	C	100	Total	C	N	O	S	0	0
			780	492	132	151	5		
1	I	100	Total	C	N	O	S	0	0
			780	492	132	151	5		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SEQUESTOSOME-1




- Molecule 1: SEQUESTOSOME-1



- Molecule 1: SEQUESTOSOME-1



- Molecule 1: SEQUESTOSOME-1



## 4 Experimental information

Property	Value	Source
Reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=Not provided°, rise=Not provided Å, axial sym=Not provided	Depositor
Number of segments used	50620	Depositor
Resolution determination method	Not provided	Depositor
CTF correction method	CTFFIND, CONVOLUTION IMAGES WIENER FILTER RECONSTRUCTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	15	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	5000	Depositor
Magnification	75000	Depositor
Image detector	FEI FALCON II (4k x 4k)	Depositor

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 2$	RMSZ	$\# Z  > 2$
1	A	0.56	0/796	0.91	2/1071 (0.2%)
1	B	0.56	0/796	0.91	2/1071 (0.2%)
1	C	0.56	0/796	0.91	2/1071 (0.2%)
1	I	0.56	0/796	0.91	2/1071 (0.2%)
All	All	0.56	0/3184	0.91	8/4284 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	37	ALA	CB-CA-C	6.08	119.22	110.10
1	C	37	ALA	CB-CA-C	6.06	119.19	110.10
1	B	37	ALA	CB-CA-C	6.06	119.19	110.10
1	I	37	ALA	CB-CA-C	6.03	119.14	110.10
1	A	37	ALA	N-CA-CB	-5.09	102.98	110.10
1	I	37	ALA	N-CA-CB	-5.05	103.03	110.10
1	C	37	ALA	N-CA-CB	-5.04	103.05	110.10
1	B	37	ALA	N-CA-CB	-5.04	103.05	110.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	780	0	758	21	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	780	0	758	14	0
1	C	780	0	758	13	0
1	I	780	0	758	21	0
All	All	3120	0	3032	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:32:GLU:HG2	1:C:37:ALA:HB3	1.55	0.87
1:A:32:GLU:HG2	1:A:37:ALA:HB3	1.55	0.86
1:I:32:GLU:HG2	1:I:37:ALA:HB3	1.55	0.86
1:B:32:GLU:HG2	1:B:37:ALA:HB3	1.55	0.86
1:A:59:ARG:NH2	1:I:92:ASP:HB3	1.96	0.80
1:A:59:ARG:HH22	1:I:92:ASP:CB	2.02	0.73
1:A:59:ARG:NH2	1:I:92:ASP:OD1	2.27	0.66
1:A:59:ARG:HH22	1:I:92:ASP:HB3	1.57	0.66
1:A:90:VAL:HG21	1:A:94:ILE:O	1.98	0.64
1:C:90:VAL:HG21	1:C:94:ILE:O	1.98	0.64
1:B:90:VAL:HG21	1:B:94:ILE:O	1.98	0.64
1:I:90:VAL:HG21	1:I:94:ILE:O	1.98	0.64
1:A:59:ARG:HH22	1:I:92:ASP:CG	2.01	0.63
1:A:32:GLU:HG2	1:A:37:ALA:CB	2.29	0.62
1:I:32:GLU:HG2	1:I:37:ALA:CB	2.29	0.61
1:B:32:GLU:HG2	1:B:37:ALA:CB	2.29	0.60
1:C:32:GLU:HG2	1:C:37:ALA:CB	2.29	0.60
1:A:58:LEU:HD12	1:A:59:ARG:O	2.04	0.58
1:I:58:LEU:HD12	1:I:59:ARG:O	2.04	0.57
1:C:58:LEU:HD12	1:C:59:ARG:O	2.04	0.57
1:A:59:ARG:NH1	1:I:92:ASP:OD1	2.35	0.57
1:B:34:GLU:OE2	1:B:36:GLU:HB3	2.05	0.57
1:I:34:GLU:OE2	1:I:36:GLU:HB3	2.05	0.57
1:C:34:GLU:OE2	1:C:36:GLU:HB3	2.05	0.57
1:B:58:LEU:HD12	1:B:59:ARG:O	2.04	0.57
1:A:34:GLU:OE2	1:A:36:GLU:HB3	2.05	0.56
1:A:35:ALA:HA	1:A:38:ALA:O	2.06	0.56
1:A:32:GLU:HG3	1:A:33:ALA:N	2.21	0.55
1:A:32:GLU:HG3	1:A:33:ALA:H	1.71	0.55
1:C:32:GLU:HG3	1:C:33:ALA:N	2.22	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:35:ALA:HA	1:C:38:ALA:O	2.06	0.55
1:I:32:GLU:HG3	1:I:33:ALA:H	1.72	0.55
1:I:35:ALA:HA	1:I:38:ALA:O	2.06	0.55
1:I:32:GLU:HG3	1:I:33:ALA:N	2.22	0.55
1:B:32:GLU:HG3	1:B:33:ALA:H	1.72	0.55
1:B:35:ALA:HA	1:B:38:ALA:O	2.06	0.55
1:B:32:GLU:HG3	1:B:33:ALA:N	2.22	0.55
1:C:32:GLU:HG3	1:C:33:ALA:H	1.72	0.54
1:A:11:LEU:HD12	1:A:100:LYS:HG3	1.93	0.51
1:I:11:LEU:HD12	1:I:100:LYS:HG3	1.93	0.51
1:B:11:LEU:HD12	1:B:100:LYS:HG3	1.93	0.51
1:C:11:LEU:HD12	1:C:100:LYS:HG3	1.93	0.51
1:C:66:HIS:CD2	1:C:76:ALA:HB2	2.52	0.45
1:I:66:HIS:CD2	1:I:76:ALA:HB2	2.52	0.45
1:B:66:HIS:CD2	1:B:76:ALA:HB2	2.52	0.45
1:A:66:HIS:CD2	1:A:76:ALA:HB2	2.52	0.45
1:A:69:ASP:OD2	1:A:75:VAL:HG21	2.18	0.44
1:B:67:TYR:CE2	1:B:69:ASP:OD1	2.71	0.43
1:I:67:TYR:CE2	1:I:69:ASP:OD1	2.71	0.43
1:C:69:ASP:OD2	1:C:75:VAL:HG21	2.18	0.43
1:C:67:TYR:CE2	1:C:69:ASP:OD1	2.71	0.43
1:A:67:TYR:CE2	1:A:69:ASP:OD1	2.71	0.43
1:I:69:ASP:OD2	1:I:75:VAL:HG21	2.18	0.43
1:B:69:ASP:OD2	1:B:75:VAL:HG21	2.18	0.43
1:A:7:LYS:HG2	1:A:22:ARG:HG2	2.01	0.42
1:B:7:LYS:HG2	1:B:22:ARG:HG2	2.01	0.42
1:I:7:LYS:HG2	1:I:22:ARG:HG2	2.01	0.42
1:C:7:LYS:HG2	1:C:22:ARG:HG2	2.01	0.41
1:B:58:LEU:HD22	1:B:101:GLU:HB3	2.03	0.41
1:I:58:LEU:HD22	1:I:101:GLU:HB3	2.03	0.41
1:A:59:ARG:CZ	1:I:92:ASP:OD1	2.69	0.41
1:A:58:LEU:HD22	1:A:101:GLU:HB3	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM



entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/100 (98%)	87 (89%)	7 (7%)	4 (4%)	3	30
1	B	98/100 (98%)	87 (89%)	7 (7%)	4 (4%)	3	30
1	C	98/100 (98%)	87 (89%)	7 (7%)	4 (4%)	3	30
1	I	98/100 (98%)	87 (89%)	7 (7%)	4 (4%)	3	30
All	All	392/400 (98%)	348 (89%)	28 (7%)	16 (4%)	6	30

All (16) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	78	SER
1	B	78	SER
1	C	78	SER
1	I	78	SER
1	A	17	ALA
1	A	76	ALA
1	B	76	ALA
1	C	17	ALA
1	C	76	ALA
1	I	17	ALA
1	I	76	ALA
1	A	42	GLY
1	B	17	ALA
1	B	42	GLY
1	C	42	GLY
1	I	42	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/80 (100%)	76 (95%)	4 (5%)	28	60

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	80/80 (100%)	76 (95%)	4 (5%)	28	60
1	C	80/80 (100%)	76 (95%)	4 (5%)	28	60
1	I	80/80 (100%)	76 (95%)	4 (5%)	28	60
All	All	320/320 (100%)	304 (95%)	16 (5%)	33	60

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	10	LEU
1	A	85	MET
1	A	101	GLU
1	B	5	THR
1	B	10	LEU
1	B	85	MET
1	B	101	GLU
1	C	5	THR
1	C	10	LEU
1	C	85	MET
1	C	101	GLU
1	I	5	THR
1	I	10	LEU
1	I	85	MET
1	I	101	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.