



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 10:36 pm GMT

PDB ID : 3UJN
Title : Formyl Glycinamide Ribonucleotide Amidotransferase from Salmonella Typhimurium : Role of the ATP complexation and glutaminase domain in catalytic coupling
Authors : Anand, R.; Morar, M.; Tanwar, A.S.; Panjekar, S.
Deposited on : 2011-11-08
Resolution : 2.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2168 (3.00-2.96)
Clashscore	112137	2535 (3.00-2.96)
Ramachandran outliers	110173	2451 (3.00-2.96)
Sidechain outliers	110143	2454 (3.00-2.96)
RSRZ outliers	101464	2192 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1303	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="position: absolute; bottom: -10px; left: 0;">86%</div> <div style="position: absolute; bottom: -10px; right: 0;">12% ..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	2009	-	-	X	X
3	ADP	A	2005	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MG	A	2006	-	-	-	X
4	MG	A	2007	-	-	-	X
4	MG	A	2008	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 10205 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphoribosylformylglycinamide synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1286	9910	6217	1770	1875	48	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	EXPRESSION TAG	UNP P74881
A	-6	LEU	-	EXPRESSION TAG	UNP P74881
A	-5	VAL	-	EXPRESSION TAG	UNP P74881
A	-4	PRO	-	EXPRESSION TAG	UNP P74881
A	-3	ARG	-	EXPRESSION TAG	UNP P74881
A	-2	GLY	-	EXPRESSION TAG	UNP P74881
A	-1	SER	-	EXPRESSION TAG	UNP P74881
A	0	HIS	-	EXPRESSION TAG	UNP P74881

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Mg	0	0
			3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	255	Total	O	0	0
			255	255		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphoribosylformylglycinamide synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	146.96Å 146.96Å 141.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.98 19.80 – 2.98	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.98) 99.2 (19.80-2.98)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.98Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.208 , 0.264 0.219 , 0.218	Depositor DCC
R_{free} test set	1030 reflections (2.95%)	DCC
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 25.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.010 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	10205	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CYG, SO4, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/10107	0.78	7/13717 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	332	LEU	CA-CB-CG	9.00	136.00	115.30
1	A	409	LYS	N-CA-C	-6.04	94.69	111.00
1	A	788	GLY	N-CA-C	-5.37	99.68	113.10
1	A	1144	LEU	CA-CB-CG	5.36	127.63	115.30
1	A	385	LEU	N-CA-C	5.20	125.04	111.00
1	A	43	LEU	CA-CB-CG	5.05	126.92	115.30
1	A	1236	ASN	N-CA-C	-5.00	97.49	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1135	CYG	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9910	0	9693	84	0
2	A	10	0	0	6	0
3	A	27	0	11	1	0
4	A	3	0	0	0	0
5	A	255	0	0	6	0
All	All	10205	0	9704	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:PHE:HD2	2:A:2009:SO4:O3	1.57	0.86
1:A:13:ALA:HB3	2:A:2009:SO4:O1	1.77	0.85
1:A:13:ALA:HB3	2:A:2009:SO4:S	2.20	0.81
1:A:824:ASN:HD21	1:A:958:ALA:H	1.34	0.74
1:A:175:ASN:HD21	1:A:182:LEU:H	1.36	0.74
1:A:175:ASN:HD22	1:A:180:LEU:HB2	1.56	0.71
1:A:637:VAL:HG13	1:A:891:LEU:HD21	1.75	0.68
1:A:317:ARG:HH22	1:A:548:GLN:NE2	1.94	0.66
1:A:387:GLY:HA2	1:A:697:ILE:HD12	1.79	0.64
1:A:13:ALA:CB	2:A:2009:SO4:O1	2.46	0.62
1:A:1135:CYG:O	1:A:1137:GLY:N	2.35	0.59
1:A:14:PHE:CD2	2:A:2009:SO4:O3	2.49	0.59
1:A:723:ILE:HD12	1:A:728:ILE:HD11	1.85	0.57
1:A:396:VAL:HG13	1:A:850:LEU:HD22	1.87	0.57
1:A:329:LYS:NZ	1:A:419:ASN:HD21	2.03	0.57
1:A:786:GLN:HE22	1:A:791:GLN:HG2	1.72	0.55
1:A:-3:ARG:HH22	1:A:152:HIS:CE1	2.24	0.55
1:A:120:THR:H	1:A:123:GLN:HE21	1.53	0.55
1:A:1004:GLN:NE2	1:A:1233:TYR:H	2.03	0.55
1:A:725:ALA:HB2	1:A:882:TRP:CD1	2.41	0.55
1:A:1004:GLN:HE21	1:A:1233:TYR:H	1.55	0.54
1:A:820:SER:H	1:A:930:GLN:HE22	1.56	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:326:ALA:HB2	1:A:420:ILE:HD12	1.91	0.52
1:A:513:PRO:O	1:A:517:SER:HB2	2.09	0.52
1:A:628:ARG:HA	1:A:631:ILE:HD13	1.92	0.50
1:A:383:PRO:HD3	1:A:660:VAL:HG13	1.93	0.50
1:A:633:ILE:O	1:A:637:VAL:HG23	2.13	0.49
1:A:317:ARG:HH22	1:A:548:GLN:HE22	1.60	0.49
1:A:1241:PRO:O	1:A:1244:ILE:HG12	2.13	0.49
1:A:175:ASN:ND2	1:A:180:LEU:HB2	2.27	0.48
1:A:742:ALA:HB2	1:A:798:LEU:HD12	1.96	0.48
1:A:168:ARG:NH1	1:A:188:ASP:OD1	2.46	0.48
1:A:358:ARG:HD3	1:A:358:ARG:H	1.78	0.48
1:A:-3:ARG:NH1	1:A:0:HIS:HB2	2.29	0.47
1:A:96:HIS:HE1	1:A:103:VAL:O	1.98	0.47
1:A:824:ASN:HD21	1:A:958:ALA:N	2.08	0.47
1:A:1218:VAL:HG13	1:A:1244:ILE:HG23	1.97	0.47
1:A:1175:THR:HG21	1:A:1221:ARG:HE	1.80	0.47
1:A:305:TRP:HB3	1:A:306:PRO:HD3	1.97	0.46
1:A:882:TRP:CE3	1:A:929:ILE:HG23	2.50	0.46
1:A:1222:TYR:OH	1:A:1245:THR:HG21	2.15	0.46
1:A:379:GLU:HB3	1:A:475:PRO:HB2	1.97	0.46
1:A:81:PRO:HD3	1:A:138:THR:HG21	1.97	0.46
1:A:114:ILE:HD11	1:A:131:LEU:HD11	1.98	0.46
1:A:257:LYS:NZ	1:A:426:GLN:HE22	2.14	0.45
1:A:396:VAL:HG22	1:A:850:LEU:HB2	1.98	0.45
1:A:916:HIS:HD2	5:A:1359:HOH:O	1.98	0.45
1:A:696:SER:OG	1:A:804:ALA:HB3	2.17	0.45
1:A:123:GLN:O	1:A:127:VAL:HG23	2.17	0.45
1:A:727:GLN:HG3	5:A:1444:HOH:O	2.17	0.44
1:A:873:LEU:HD13	1:A:879:LEU:HD21	1.99	0.44
1:A:483:GLU:O	1:A:487:ARG:HG2	2.17	0.44
1:A:654:THR:HA	1:A:658:ARG:HH12	1.82	0.44
1:A:870:MET:O	1:A:874:VAL:HG23	2.17	0.44
1:A:329:LYS:O	1:A:383:PRO:HD2	2.18	0.43
1:A:371:LEU:HD21	1:A:1164:ARG:HD2	2.01	0.43
1:A:501:HIS:HD1	1:A:502:ASP:N	2.16	0.43
1:A:782:LYS:NZ	1:A:784:ARG:HH22	2.15	0.43
1:A:398:SER:HB3	1:A:399:HIS:H	1.64	0.42
1:A:411:ILE:HG12	1:A:741:MET:HG2	2.02	0.42
1:A:1006:HIS:HE1	5:A:1523:HOH:O	2.01	0.42
1:A:1220:LEU:H	1:A:1246:ALA:HB3	1.83	0.42
1:A:13:ALA:HB3	2:A:2009:SO4:O3	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:824:ASN:ND2	1:A:958:ALA:H	2.08	0.42
1:A:668:GLN:HG2	3:A:2005:ADP:H1'	2.02	0.41
1:A:1041:LYS:HD2	1:A:1068:ILE:HD11	2.03	0.41
1:A:5:LEU:HD22	1:A:59:LEU:HD12	2.03	0.41
1:A:776:LYS:HG2	5:A:1434:HOH:O	2.21	0.41
1:A:1192:ILE:HD13	1:A:1193:ALA:H	1.85	0.41
1:A:1:MET:HG3	5:A:1389:HOH:O	2.21	0.41
1:A:725:ALA:HB2	1:A:882:TRP:HD1	1.84	0.41
1:A:1196:HIS:NE2	1:A:1238:ASN:ND2	2.63	0.41
1:A:317:ARG:NH1	1:A:549:GLU:H	2.19	0.41
1:A:336:SER:HB2	1:A:411:ILE:HB	2.02	0.41
1:A:310:THR:HG21	1:A:471:GLN:HB3	2.03	0.41
1:A:1134:VAL:O	1:A:1135:CYG:O	2.38	0.41
1:A:1204:ARG:NH2	5:A:1327:HOH:O	2.52	0.41
1:A:1201:VAL:HG21	1:A:1244:ILE:HB	2.03	0.41
1:A:293:VAL:HB	1:A:739:ASN:HD21	1.86	0.40
1:A:329:LYS:HZ3	1:A:685:ALA:HA	1.87	0.40
1:A:828:LEU:HD13	1:A:890:LEU:HD13	2.03	0.40
1:A:253:LEU:HB3	1:A:425:VAL:HG11	2.03	0.40
1:A:625:ALA:HA	1:A:852:ASP:HB2	2.04	0.40
1:A:753:LEU:HD13	1:A:798:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1281/1303 (98%)	1208 (94%)	66 (5%)	7 (0%)	32 72

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	13	ALA
1	A	1136	ASN
1	A	117	SER
1	A	623	GLY
1	A	647	ALA
1	A	1275	GLU
1	A	-4	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1029/1040 (99%)	984 (96%)	45 (4%)	33 70

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-6	LEU
1	A	-5	VAL
1	A	-4	PRO
1	A	-3	ARG
1	A	6	ARG
1	A	43	LEU
1	A	104	ASP
1	A	141	SER
1	A	169	GLN
1	A	215	GLU
1	A	253	LEU
1	A	258	ASP
1	A	290	LEU
1	A	332	LEU
1	A	337	VAL
1	A	358	ARG
1	A	382	ARG
1	A	412	MET
1	A	413	LEU
1	A	508	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	510	ASN
1	A	515	LEU
1	A	533	ASP
1	A	571	ARG
1	A	582	GLU
1	A	588	LEU
1	A	596	GLN
1	A	610	PRO
1	A	622	LYS
1	A	723	ILE
1	A	750	ASP
1	A	774	VAL
1	A	783	THR
1	A	789	ASN
1	A	790	GLU
1	A	823	ASP
1	A	846	VAL
1	A	891	LEU
1	A	976	GLU
1	A	1180	LEU
1	A	1190	MET
1	A	1192	ILE
1	A	1195	SER
1	A	1205	ASP
1	A	1245	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	HIS
1	A	96	HIS
1	A	102	GLN
1	A	123	GLN
1	A	175	ASN
1	A	276	ASN
1	A	298	HIS
1	A	408	HIS
1	A	419	ASN
1	A	426	GLN
1	A	445	ASN
1	A	471	GLN
1	A	510	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	545	ASN
1	A	548	GLN
1	A	739	ASN
1	A	786	GLN
1	A	824	ASN
1	A	883	HIS
1	A	922	ASN
1	A	930	GLN
1	A	969	ASN
1	A	993	GLN
1	A	1004	GLN
1	A	1006	HIS
1	A	1061	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	CYG	A	1135	1	9,14,15	5.20	2 (22%)	6,17,19	6.33	4 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CYG	A	1135	1	-	0/10/16/18	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1135	CYG	CA-C	2.50	1.53	1.50
1	A	1135	CYG	OE2-CD1	15.24	1.44	1.21

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1135	CYG	OE2-CD1-CG1	-10.99	114.33	123.95
1	A	1135	CYG	O-C-CA	-5.84	108.88	125.02
1	A	1135	CYG	CB1-CG1-CD1	-3.93	107.31	113.12
1	A	1135	CYG	CG1-CD1-SG	8.26	121.60	113.28

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1135	CYG	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	2002	-	4,4,4	0.51	0	6,6,6	0.20	0
3	ADP	A	2005	4	25,29,29	1.39	3 (12%)	24,45,45	2.15	1 (4%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	2009	-	4,4,4	0.33	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	2002	-	-	0/0/0/0	0/0/0/0
3	ADP	A	2005	4	-	0/12/32/32	0/3/3/3
2	SO4	A	2009	-	-	0/0/0/0	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2005	ADP	O3'-C3'	-5.20	1.31	1.43
3	A	2005	ADP	C5-N7	-2.22	1.31	1.39
3	A	2005	ADP	O4'-C1'	2.29	1.44	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2005	ADP	N3-C2-N1	-9.82	120.31	128.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2005	ADP	1	0
2	A	2009	SO4	6	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	1285/1303 (98%)	-0.32	16 (1%)	79 59	10, 26, 46, 88	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-4	PRO	7.1
1	A	-7	GLY	6.6
1	A	-5	VAL	5.9
1	A	-6	LEU	4.9
1	A	-2	GLY	3.9
1	A	118	THR	3.5
1	A	-1	SER	3.2
1	A	610	PRO	3.0
1	A	-3	ARG	3.0
1	A	216	HIS	2.6
1	A	154	GLN	2.5
1	A	276	ASN	2.4
1	A	66	SER	2.4
1	A	152	HIS	2.3
1	A	790	GLU	2.2
1	A	31	HIS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	CYG	A	1135	15/16	0.94	0.17	-	25,27,30,30	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SO4	A	2009	5/5	0.89	0.45	8.44	23,25,27,27	0
4	MG	A	2007	1/1	0.98	0.25	8.38	7,7,7,7	0
3	ADP	A	2005	27/27	0.95	0.35	7.99	1,1,3,5	27
4	MG	A	2006	1/1	0.60	0.25	4.24	15,15,15,15	0
4	MG	A	2008	1/1	0.89	0.22	3.47	11,11,11,11	0
2	SO4	A	2002	5/5	0.94	0.19	-0.02	38,38,40,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.