



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 11, 2017 – 05:52 PM EDT

PDB ID : 5ULX  
Title : Structure of human DNA polymerase iota bound to template 1-methyl-deoxy adenosine crystallized in the presence of dCTP  
Authors : Jain, R.; Aggarwal, A.K.  
Deposited on : unknown  
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	7	<div> <div>29%</div> <div>71%</div> </div>
2	T	11	<div> <div>36%</div> <div>45%</div> <div>18%</div> </div>
3	A	420	<div> <div>9%</div> <div>65%</div> <div>23%</div> <div>• 11%</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3442 atoms, of which 26 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*AP\*GP\*GP\*AP\*CP\*CP\*(DOC))-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	P	7	Total	C	H	N	O	P	0	0	0
			151	67	12	29	37	6			

- Molecule 2 is a DNA chain called DNA (5'-D(P\*(MA7)P\*GP\*GP\*GP\*TP\*CP\*CP\*T)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	T	9	Total	C	H	N	O	P	0	0	1
			181	79	14	30	50	8			

- Molecule 3 is a protein called DNA polymerase iota.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	373	Total	C	N	O	S	0	3	0
			2818	1789	490	518	21			

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	P	13	Total	O	0	0
			13	13		
5	T	21	Total	O	0	0
			21	21		
5	A	257	Total	O	0	0
			257	257		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(\*AP\*GP\*GP\*AP\*CP\*CP\*(DOC))-3')

Chain P: 




- Molecule 2: DNA (5'-D(P\*(MA7)P\*GP\*GP\*GP\*TP\*CP\*CP\*T)-3')

Chain T: 



- Molecule 3: DNA polymerase iota

Chain A: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.92Å 97.92Å 202.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.96 – 1.96 48.96 – 1.96	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.96-1.96) 99.9 (48.96-1.96)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 1.97Å)	Xtriage
Refinement program	PHENIX 1.10.1 _2155	Depositor
R, $R_{free}$	0.216 , 0.239 0.215 , 0.238	Depositor DCC
$R_{free}$ test set	3151 reflections (7.53%)	DCC
Wilson B-factor (Å <sup>2</sup> )	33.3	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 63.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3442	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DOC, MA7, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	P	0.60	0/136	0.75	0/208
2	T	0.67	0/160	0.92	0/245
3	A	0.29	0/2866	0.49	0/3884
All	All	0.34	0/3162	0.54	0/4337

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	139	12	79	3	1
2	T	167	14	94	4	1
3	A	2818	0	2794	98	0
4	A	1	0	0	1	0
5	A	257	0	0	17	0
5	P	13	0	0	0	0
5	T	21	0	0	1	0
All	All	3416	26	2967	104	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:T:840:MA7:O4'	2:T:840:MA7:C4'	1.69	1.21
1:P:873:DOC:O4'	1:P:873:DOC:C1'	1.65	1.18
3:A:409:CYS:SG	5:A:792:HOH:O	2.18	1.00
3:A:290:ASN:ND2	4:A:501:CL:CL	2.37	0.94
3:A:365:PRO:HB2	3:A:368:VAL:HG23	1.57	0.83
3:A:189[B]:ASN:ND2	5:A:605:HOH:O	2.11	0.83
3:A:90:GLN:OE1	3:A:90:GLN:N	2.12	0.80
3:A:251:GLU:OE2	5:A:603:HOH:O	2.00	0.78
3:A:185:GLU:OE1	5:A:602:HOH:O	2.00	0.78
3:A:326:ALA:HA	3:A:380:MET:HE1	1.66	0.77
3:A:304:GLU:HG3	3:A:328:LEU:HG	1.67	0.77
3:A:247:ALA:O	3:A:251:GLU:HG3	1.84	0.76
3:A:314:GLU:HG2	3:A:391:PHE:HZ	1.49	0.76
3:A:380:MET:SD	5:A:807:HOH:O	2.46	0.73
3:A:302:PHE:HE1	3:A:413:LEU:HD21	1.55	0.70
3:A:380:MET:O	3:A:384:VAL:HG23	1.91	0.69
3:A:411:CYS:SG	5:A:792:HOH:O	2.51	0.67
3:A:365:PRO:HB2	3:A:368:VAL:CG2	2.24	0.66
3:A:335:ASP:OD2	3:A:337:ARG:HD3	1.94	0.66
3:A:302:PHE:CE1	3:A:413:LEU:HD21	2.30	0.66
3:A:101:ARG:NH2	5:A:610:HOH:O	2.23	0.65
3:A:27:SER:OG	5:A:606:HOH:O	2.14	0.65
3:A:314:GLU:HG2	3:A:391:PHE:CZ	2.31	0.65
3:A:306:ASP:O	3:A:405:LEU:HD12	1.96	0.65
3:A:391:PHE:CZ	3:A:395:VAL:HG11	2.32	0.64
3:A:117:SER:OG	5:A:601:HOH:O	1.88	0.64
3:A:413:LEU:O	3:A:414:LYS:HG3	1.98	0.63
3:A:236:ILE:CD1	3:A:261:LEU:HB2	2.30	0.62
3:A:343:ARG:HG2	3:A:345:ILE:HD11	1.82	0.62
3:A:347:ARG:HE	3:A:404:THR:HG23	1.65	0.61
3:A:236:ILE:HD12	3:A:255:ILE:HG22	1.83	0.61
3:A:236:ILE:HG22	3:A:242:ILE:HG21	1.84	0.59
3:A:265:SER:HB3	3:A:268:ILE:HD13	1.84	0.59
3:A:396:ASN:O	5:A:607:HOH:O	2.17	0.59
2:T:840:MA7:H8	3:A:59:GLN:OE1	2.04	0.58
3:A:343:ARG:HD2	3:A:359:SER:HB2	1.87	0.57
3:A:265:SER:CB	3:A:268:ILE:HD13	2.34	0.56
3:A:61:TYR:O	3:A:81:VAL:HG23	2.05	0.56
2:T:846:DC:OP1	5:T:901:HOH:O	2.18	0.55
3:A:35:LEU:HD23	5:A:709:HOH:O	2.07	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:384:VAL:O	3:A:388:MET:HG2	2.07	0.55
3:A:325:LEU:HD21	3:A:383:MET:HB2	1.90	0.54
3:A:339:PRO:HG3	3:A:410:PHE:HB3	1.88	0.54
3:A:236:ILE:HD11	3:A:261:LEU:HB2	1.91	0.53
3:A:105:MET:CG	3:A:193:LEU:HD11	2.39	0.53
3:A:280:ARG:NH2	5:A:609:HOH:O	2.20	0.53
3:A:61:TYR:C	3:A:81:VAL:HG23	2.30	0.53
3:A:391:PHE:O	3:A:395:VAL:HG12	2.09	0.52
3:A:55:LEU:HD12	3:A:56:GLY:H	1.76	0.51
3:A:236:ILE:HD13	3:A:261:LEU:HB2	1.93	0.51
3:A:27:SER:N	5:A:606:HOH:O	2.33	0.51
3:A:339:PRO:CG	3:A:410:PHE:HB3	2.41	0.50
3:A:55:LEU:HD12	3:A:56:GLY:N	2.26	0.50
3:A:339:PRO:CB	3:A:410:PHE:HB3	2.42	0.50
3:A:105:MET:HE1	5:A:748:HOH:O	2.11	0.50
3:A:386:ILE:O	3:A:390:LEU:HG	2.12	0.50
3:A:36:ASP:OD1	5:A:608:HOH:O	2.19	0.49
3:A:152:THR:HG22	5:A:656:HOH:O	2.13	0.49
3:A:347:ARG:HB3	3:A:404:THR:HG23	1.95	0.49
1:P:870:DA:H2'	1:P:871:DC:C6	2.48	0.48
3:A:249:CYS:SG	3:A:273:LEU:HD21	2.53	0.48
3:A:236:ILE:CG2	3:A:242:ILE:HD13	2.44	0.48
3:A:266:PRO:O	3:A:270:GLU:HG3	2.14	0.47
3:A:103:ARG:HD3	5:A:791:HOH:O	2.13	0.47
3:A:203:LYS:HD3	3:A:293:VAL:HG22	1.96	0.47
3:A:381:THR:OG1	3:A:382:PRO:HD3	2.14	0.47
3:A:329:LEU:O	3:A:332:VAL:HG12	2.15	0.47
3:A:59:GLN:OE1	3:A:64[B]:VAL:HG11	2.14	0.46
3:A:302:PHE:HE1	3:A:413:LEU:CD2	2.25	0.46
3:A:163:ILE:HG21	3:A:174:LEU:HD11	1.97	0.46
3:A:121:GLU:HB2	3:A:294:ILE:O	2.15	0.46
3:A:313:SER:C	3:A:315:VAL:H	2.19	0.46
3:A:310:LYS:HE3	3:A:349:TYR:CB	2.45	0.46
3:A:304:GLU:HG3	3:A:328:LEU:CG	2.42	0.46
3:A:345:ILE:HD12	3:A:345:ILE:N	2.30	0.46
3:A:267:LYS:H	3:A:267:LYS:CD	2.27	0.45
3:A:343:ARG:HG2	3:A:345:ILE:CD1	2.46	0.45
3:A:309:LYS:HD3	3:A:309:LYS:HA	1.52	0.45
3:A:105:MET:HG3	3:A:193:LEU:HD11	1.99	0.45
3:A:236:ILE:CD1	3:A:255:ILE:HG22	2.45	0.45
2:T:844:DT:H2''	2:T:845:DC:H5'	1.99	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:148:LEU:HD23	3:A:148:LEU:C	2.37	0.45
3:A:255:ILE:HD11	3:A:264:PHE:CG	2.51	0.44
3:A:60:LYS:HE2	3:A:307:SER:O	2.17	0.44
3:A:100:THR:O	3:A:104:GLU:HG3	2.18	0.44
3:A:313:SER:O	3:A:315:VAL:N	2.51	0.44
3:A:36:ASP:O	3:A:37:CYS:C	2.55	0.44
3:A:50:LEU:CD2	3:A:92:VAL:HG11	2.48	0.44
3:A:379:VAL:HG23	3:A:379:VAL:O	2.18	0.44
3:A:268:ILE:HD12	3:A:268:ILE:N	2.33	0.43
3:A:265:SER:OG	3:A:267:LYS:HG2	2.18	0.43
3:A:413:LEU:C	3:A:414:LYS:HG3	2.39	0.42
3:A:391:PHE:CE1	3:A:395:VAL:HG11	2.54	0.42
3:A:277:VAL:O	3:A:281:ILE:HG23	2.19	0.42
3:A:329:LEU:HA	3:A:332:VAL:HG12	2.00	0.42
3:A:325:LEU:O	3:A:329:LEU:HG	2.20	0.42
3:A:413:LEU:HD12	3:A:413:LEU:N	2.35	0.42
3:A:339:PRO:HB3	3:A:410:PHE:HB3	2.00	0.41
3:A:379:VAL:O	3:A:383:MET:HG2	2.19	0.41
3:A:31:VAL:CG1	3:A:130:VAL:HB	2.50	0.41
3:A:303:SER:O	3:A:304:GLU:HG2	2.21	0.41
3:A:153:VAL:HB	3:A:174:LEU:HD22	2.03	0.41
3:A:255:ILE:HD11	3:A:264:PHE:CD1	2.56	0.41
1:P:871:DC:H2''	1:P:872:DC:H5'	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:P:867:DA:O5'	2:T:847:DT:O3'[10_665]	1.81	0.39

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	368/420 (88%)	357 (97%)	8 (2%)	3 (1%)	22	10

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	37	CYS
3	A	310	LYS
3	A	314	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	302/376 (80%)	293 (97%)	9 (3%)	46	34

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	52	ASP
3	A	64[A]	VAL
3	A	64[B]	VAL
3	A	80	ASN
3	A	272	GLU
3	A	308	PHE
3	A	309	LYS
3	A	312	SER
3	A	394	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	DOC	P	873	1,2	13,19,20	5.43	10 (76%)	12,26,29	1.23	1 (8%)
2	MA7	T	840	2	17,24,25	3.83	10 (58%)	15,35,38	2.26	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DOC	P	873	1,2	-	0/3/18/19	0/2/2/2
2	MA7	T	840	2	-	0/3/21/22	0/3/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	P	873	DOC	C2'-C1'	-7.43	1.33	1.51
1	P	873	DOC	O4'-C4'	-6.82	1.29	1.44
2	T	840	MA7	C3'-C4'	-6.72	1.34	1.53
2	T	840	MA7	O4'-C1'	-4.60	1.31	1.42
2	T	840	MA7	O5'-C5'	-3.80	1.39	1.44
1	P	873	DOC	O5'-C5'	-3.37	1.40	1.44
2	T	840	MA7	C6-N1	-2.75	1.31	1.36
2	T	840	MA7	C5-C4	-2.32	1.35	1.40
2	T	840	MA7	C2'-C3'	2.02	1.58	1.52
2	T	840	MA7	C2'-C1'	2.91	1.60	1.52
1	P	873	DOC	C5-C4	3.00	1.48	1.41
2	T	840	MA7	C6-N6	3.12	1.48	1.35
1	P	873	DOC	C4-N4	3.14	1.45	1.35
2	T	840	MA7	C2-N3	3.49	1.36	1.30
1	P	873	DOC	C6-C5	5.13	1.49	1.38
1	P	873	DOC	C4-N3	5.58	1.45	1.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	P	873	DOC	C2-N3	5.63	1.49	1.38
1	P	873	DOC	C6-N1	7.43	1.45	1.35
1	P	873	DOC	O4'-C1'	10.18	1.65	1.42
2	T	840	MA7	O4'-C4'	10.62	1.69	1.45

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	840	MA7	C4'-O4'-C1'	-2.78	102.62	109.42
2	T	840	MA7	O4'-C1'-C2'	-2.35	101.73	106.25
2	T	840	MA7	C2'-C3'-C4'	2.26	107.55	102.73
1	P	873	DOC	C3'-C2'-C1'	2.50	105.47	102.69
2	T	840	MA7	O4'-C1'-N9	2.51	112.01	107.78
2	T	840	MA7	N6-C6-N1	6.74	124.97	120.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	P	873	DOC	1	0
2	T	840	MA7	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	P	6/7 (85%)	0.02	0 100 100	33, 47, 51, 56	0
2	T	8/11 (72%)	-0.07	0 100 100	31, 37, 46, 63	0
3	A	373/420 (88%)	0.63	38 (10%) 7 12	20, 44, 83, 105	0
All	All	387/438 (88%)	0.61	38 (9%) 8 13	20, 44, 82, 105	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	333	CYS	7.3
3	A	395	VAL	6.9
3	A	349	TYR	5.7
3	A	397	VAL	5.7
3	A	348	ARG	5.5
3	A	367	HIS	4.8
3	A	381	THR	4.6
3	A	311	CYS	4.6
3	A	312	SER	4.6
3	A	334	GLN	4.5
3	A	391	PHE	4.4
3	A	403	LEU	4.1
3	A	379	VAL	4.1
3	A	350	SER	4.0
3	A	332	VAL	3.6
3	A	368	VAL	3.5
3	A	390	LEU	3.2
3	A	396	ASN	3.2
3	A	335	ASP	3.1
3	A	244	TYR	2.9
3	A	366	SER	2.9
3	A	378	ASP	2.8
3	A	380	MET	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	A	405	LEU	2.7
3	A	273	LEU	2.7
3	A	329	LEU	2.7
3	A	310	LYS	2.5
3	A	143	LEU	2.4
3	A	365	PRO	2.4
3	A	81	VAL	2.3
3	A	269	LEU	2.2
3	A	383	MET	2.2
3	A	404	THR	2.1
3	A	394	MET	2.1
3	A	313	SER	2.1
3	A	336	GLY	2.1
3	A	148	LEU	2.1
3	A	386	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q < 0.9
2	MA7	T	840	22/23	0.94	0.13	-	41,58,72,79	0
1	DOC	P	873	18/19	0.96	0.12	-	27,34,44,44	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	CL	A	501	1/1	0.81	0.09	-	85,85,85,85	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.