



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 04:10 am GMT

PDB ID : 1UMO  
Title : THE CRYSTAL STRUCTURE OF CYTOGLOBIN: THE FOURTH GLOBIN TYPE DISCOVERED IN MAN  
Authors : De Sanctis, D.; Dewilde, S.; Pesce, A.; Moens, L.; Ascenzi, P.; Hankeln, T.; Burmester, T.; Bolognesi, M.  
Deposited on : 2003-08-26  
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	<div> <div>2%</div> <div> <div></div> <div>58%</div> <div>18%</div> <div>• •</div> <div>19%</div> </div> </div>
1	B	190	<div> <div>3%</div> <div> <div></div> <div>55%</div> <div>21%</div> <div>5%</div> <div>19%</div> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2713 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOGLOBIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	154	Total	C	N	O	S	0	16	0
			1365	876	242	242	5			
1	B	154	Total	C	N	O	S	0	0	0
			1232	795	213	220	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	38	SER	CYS	ENGINEERED MUTATION	UNP Q8WWM9
A	83	SER	CYS	ENGINEERED MUTATION	UNP Q8WWM9
B	38	SER	CYS	ENGINEERED MUTATION	UNP Q8WWM9
B	83	SER	CYS	ENGINEERED MUTATION	UNP Q8WWM9

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	17	Total 17	O 17	0	0
3	B	13	Total 13	O 13	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.62Å 72.00Å 98.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.88 – 2.59 29.83 – 2.59	Depositor EDS
% Data completeness (in resolution range)	98.9 (29.88-2.59) 98.5 (29.83-2.59)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	16.19 (at 2.57Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.208 , 0.284 0.216 , 0.289	Depositor DCC
$R_{free}$ test set	514 reflections (5.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.6	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 54.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2713	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	0/1399	0.78	3/1891 (0.2%)
1	B	0.67	0/1263	0.82	5/1710 (0.3%)
All	All	0.67	0/2662	0.80	8/3601 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	98	ASP	CB-CG-OD2	7.92	125.43	118.30
1	B	68	ASP	CB-CG-OD2	7.42	124.97	118.30
1	A	98	ASP	CB-CG-OD2	7.03	124.63	118.30
1	B	142	ASP	CB-CG-OD2	5.68	123.41	118.30
1	A	40	ASP	CB-CG-OD2	5.52	123.27	118.30
1	A	68	ASP	CB-CG-OD2	5.35	123.11	118.30
1	B	40	ASP	CB-CG-OD2	5.30	123.08	118.30
1	B	63	PHE	N-CA-C	-5.21	96.93	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1365	0	1372	38	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1232	0	1232	19	0
2	A	43	0	30	1	0
2	B	43	0	30	4	0
3	A	17	0	0	3	0
3	B	13	0	0	2	0
All	All	2713	0	2664	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81[A]:HIS:O	1:A:85[A]:VAL:HG23	1.61	0.99
1:A:74[B]:ARG:O	1:A:75[B]:SER:C	2.03	0.95
1:A:75[B]:SER:HB3	1:A:78[B]:LEU:HB2	1.48	0.93
1:A:75[B]:SER:O	1:A:78[B]:LEU:N	2.04	0.91
1:B:18:GLU:O	1:B:153:LYS:NZ	2.08	0.86
1:A:100:ASP:OD2	3:A:2009:HOH:O	1.96	0.81
1:A:74[B]:ARG:O	1:A:75[B]:SER:O	2.08	0.71
1:A:75[B]:SER:HB3	1:A:78[B]:LEU:CB	2.20	0.69
1:A:72[A]:MET:O	1:A:78[A]:LEU:HG	1.95	0.66
1:A:81[A]:HIS:O	1:A:85[A]:VAL:CG2	2.42	0.65
1:B:63:PHE:O	1:B:64:LYS:CB	2.45	0.64
1:A:74[B]:ARG:C	1:A:76[B]:PRO:HD3	2.18	0.64
1:A:77[A]:GLN:NE2	1:A:80[A]:LYS:NZ	2.47	0.62
1:A:39:GLU:O	1:A:43:VAL:HG13	1.99	0.61
1:B:63:PHE:O	1:B:64:LYS:HB3	2.02	0.58
1:B:19:LEU:HA	1:B:153:LYS:NZ	2.19	0.58
1:A:159:TYR:O	1:A:163:THR:HB	2.04	0.57
2:B:1172:HEM:HBB2	2:B:1172:HEM:HMB1	1.87	0.56
1:B:87:GLY:O	1:B:91:THR:HG23	2.05	0.56
1:B:81:HIS:O	1:B:85:VAL:HG23	2.05	0.56
1:A:74[B]:ARG:C	1:A:76[B]:PRO:CD	2.75	0.55
1:B:80:LYS:O	1:B:84:ARG:HD2	2.07	0.54
1:A:75[B]:SER:O	1:A:78[B]:LEU:CB	2.56	0.53
1:A:81[B]:HIS:O	1:A:84[B]:ARG:N	2.39	0.53
1:A:77[A]:GLN:NE2	1:A:80[A]:LYS:HZ2	2.07	0.51
1:A:75[B]:SER:O	1:A:78[B]:LEU:HB3	2.10	0.51
1:B:139:PHE:O	1:B:143:PHE:HB2	2.11	0.51
1:B:51:VAL:HG21	1:B:69:PRO:HG3	1.93	0.50

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:169:VAL:HG12	1:A:169:VAL:O	2.10	0.50
1:A:115:LEU:HD21	1:A:169:VAL:CG1	2.41	0.50
1:A:47:VAL:O	1:A:51:VAL:HG23	2.12	0.50
1:A:72[B]:MET:O	1:A:73[B]:GLU:C	2.50	0.50
1:A:75[B]:SER:C	1:A:77[B]:GLN:N	2.65	0.50
1:A:120:GLU:HB3	3:A:2011:HOH:O	2.12	0.49
1:A:43:VAL:HG11	1:A:79[B]:ARG:HD2	1.95	0.49
1:A:77[A]:GLN:NE2	1:A:80[A]:LYS:HZ1	2.11	0.48
1:A:73[A]:GLU:C	1:A:74[A]:ARG:HG2	2.33	0.48
1:B:27:VAL:HG11	1:B:154:LEU:HD22	1.94	0.48
1:B:20:SER:O	1:B:24:ARG:HG2	2.14	0.48
1:A:78[B]:LEU:O	1:A:78[B]:LEU:HD13	2.15	0.47
1:A:75[B]:SER:O	1:A:78[B]:LEU:CA	2.62	0.47
1:B:21:GLU:HA	1:B:24:ARG:HG3	1.96	0.46
1:A:128:SER:HB3	1:A:155:ARG:HG3	1.98	0.46
1:A:30:MET:HE1	1:A:142:ASP:HB2	1.96	0.46
1:B:144:PRO:HB2	3:B:2011:HOH:O	2.16	0.46
1:A:115:LEU:HD21	1:A:169:VAL:HG11	1.98	0.45
1:A:75[B]:SER:C	1:A:77[B]:GLN:H	2.20	0.45
1:B:84:ARG:HG2	2:B:1172:HEM:CAA	2.47	0.45
2:B:1172:HEM:HBB2	2:B:1172:HEM:CMB	2.47	0.45
1:B:155:ARG:HD2	3:B:2012:HOH:O	2.20	0.42
1:B:96:LEU:HD23	1:B:102:VAL:HG11	2.02	0.41
1:A:34:LEU:HD11	1:A:135:VAL:HG22	2.02	0.41
1:B:19:LEU:HA	1:B:153:LYS:HZ3	1.83	0.41
1:A:74[B]:ARG:O	1:A:79[B]:ARG:HG2	2.21	0.41
1:A:121:PRO:HD2	3:A:2011:HOH:O	2.20	0.41
1:A:75[B]:SER:O	1:A:77[B]:GLN:N	2.54	0.41
1:B:84:ARG:HG2	2:B:1172:HEM:C2A	2.56	0.41
1:A:119:VAL:CG1	2:A:1172:HEM:HAC	2.51	0.41
1:B:75:SER:HA	1:B:76:PRO:HD3	1.88	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/190 (88%)	155 (92%)	10 (6%)	3 (2%)	10	19
1	B	152/190 (80%)	148 (97%)	3 (2%)	1 (1%)	25	49
All	All	320/380 (84%)	303 (95%)	13 (4%)	4 (1%)	18	29

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	64	LYS
1	A	74[A]	ARG
1	A	74[B]	ARG
1	A	145	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/162 (90%)	121 (83%)	24 (17%)	2	4
1	B	130/162 (80%)	104 (80%)	26 (20%)	1	2
All	All	275/324 (85%)	225 (82%)	50 (18%)	2	3

All (50) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LEU
1	A	30	MET
1	A	57	LYS
1	A	58	GLN
1	A	71[A]	GLU
1	A	71[B]	GLU
1	A	74[A]	ARG
1	A	74[B]	ARG
1	A	77[A]	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	77[B]	GLN
1	A	78[A]	LEU
1	A	78[B]	LEU
1	A	79[A]	ARG
1	A	79[B]	ARG
1	A	84[A]	ARG
1	A	84[B]	ARG
1	A	103	SER
1	A	104	SER
1	A	108	LEU
1	A	117	HIS
1	A	121	PRO
1	A	142	ASP
1	A	149	ARG
1	A	163	THR
1	B	18	GLU
1	B	19	LEU
1	B	33	ARG
1	B	40	ASP
1	B	46	LEU
1	B	48	ARG
1	B	67	GLU
1	B	77	GLN
1	B	79	ARG
1	B	84	ARG
1	B	89	LEU
1	B	91	THR
1	B	100	ASP
1	B	103	SER
1	B	104	SER
1	B	108	LEU
1	B	116	LYS
1	B	118	LYS
1	B	132	LEU
1	B	149	ARG
1	B	153	LYS
1	B	155	ARG
1	B	161	HIS
1	B	163	THR
1	B	167	LYS
1	B	168	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN
1	A	148	GLN
1	A	161	HIS
1	B	58	GLN
1	B	117	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	1172	1	28,50,50	2.18	7 (25%)	17,82,82	1.61	4 (23%)
2	HEM	B	1172	1	28,50,50	2.15	6 (21%)	17,82,82	1.65	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	1172	1	-	0/6/54/54	0/0/8/8
2	HEM	B	1172	1	-	0/6/54/54	0/0/8/8

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1172	HEM	C3C-C2C	-5.08	1.33	1.40
2	B	1172	HEM	C3C-C2C	-4.68	1.34	1.40
2	A	1172	HEM	C3B-C2B	-4.14	1.34	1.40
2	B	1172	HEM	C3B-C2B	-3.89	1.35	1.40
2	B	1172	HEM	CMA-C3A	2.29	1.56	1.51
2	A	1172	HEM	CAA-C2A	2.55	1.56	1.52
2	A	1172	HEM	C4D-ND	3.23	1.40	1.36
2	A	1172	HEM	C3C-CAC	3.86	1.55	1.47
2	B	1172	HEM	C3B-CAB	3.91	1.55	1.47
2	A	1172	HEM	C3B-CAB	3.97	1.55	1.47
2	B	1172	HEM	C3C-CAC	3.97	1.55	1.47
2	A	1172	HEM	C3D-C2D	4.69	1.51	1.37
2	B	1172	HEM	C3D-C2D	5.64	1.54	1.37

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1172	HEM	CAA-CBA-CGA	-2.82	107.84	112.66
2	A	1172	HEM	CMA-C3A-C4A	-2.74	124.25	128.46
2	B	1172	HEM	C1D-C2D-C3D	-2.30	105.40	107.00
2	B	1172	HEM	CMA-C3A-C4A	-2.19	125.10	128.46
2	A	1172	HEM	CAA-CBA-CGA	-2.18	108.93	112.66
2	A	1172	HEM	CAD-CBD-CGD	-2.07	109.12	112.66
2	B	1172	HEM	CMD-C2D-C1D	-2.00	125.38	128.46
2	A	1172	HEM	CMB-C2B-C3B	2.15	128.89	124.89
2	B	1172	HEM	CMD-C2D-C3D	2.26	129.21	124.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1172	HEM	1	0
2	B	1172	HEM	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	154/190 (81%)	0.21	4 (2%) 56 49	24, 40, 52, 57	0
1	B	154/190 (81%)	0.22	6 (3%) 40 32	24, 42, 59, 67	0
All	All	308/380 (81%)	0.21	10 (3%) 48 40	24, 41, 55, 67	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	141	SER	3.9
1	A	65	HIS	3.2
1	B	142	ASP	3.1
1	B	140	ALA	3.1
1	B	143	PHE	3.0
1	A	77[A]	GLN	2.6
1	A	171	TRP	2.4
1	B	170	GLY	2.2
1	A	76[A]	PRO	2.1
1	B	36	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	HEM	B	1172	43/43	0.96	0.13	-0.90	34,36,40,43	0
2	HEM	A	1172	43/43	0.96	0.14	-1.05	26,32,41,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.