



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 3, 2017 – 09:41 PM EDT

PDB ID : 5UNY
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with (RS)-3-[(2-amino-4-methylquinolin-7-yl)methoxy]-5-(2-(methylamino)propyl) benzonitrile
Authors : Li, H.; Poulos, T.L.
Deposited on : unknown
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5868 (1.84-1.80)
Clashscore	112137	6856 (1.84-1.80)
Ramachandran outliers	110173	6780 (1.84-1.80)
Sidechain outliers	110143	6780 (1.84-1.80)
RSRZ outliers	101464	5947 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>19%</div> <div> <div></div> <div>86%</div> <div>10%</div> <div>••</div> </div> </div>
1	B	422	<div> <div>9%</div> <div> <div></div> <div>89%</div> <div>8%</div> <div>•</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	M4R	A	803[A]	-	-	-	X
4	M4R	B	803[A]	-	-	-	X
5	8FD	A	804[B]	-	-	-	X
5	8FD	B	804[B]	-	-	-	X
6	ACT	A	805	-	-	-	X
6	ACT	B	805	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7354 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	410	Total	C	N	O	S	0	4	0
			3354	2146	572	614	22			
1	B	411	Total	C	N	O	S	0	3	0
			3357	2148	574	614	21			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



-
- The chemical structure of M4R is a complex organic molecule. It features a central benzene ring (C21-C26) substituted with a nitrile group (C27-C28), a chiral amine group (C29-C33), and a phenoxy group (C12-C13) linked to a pyridine ring (C04-C11) with an amine substituent (C02-C03). The structure is labeled with atom identifiers and includes stereochemical information (R/S) for the chiral center.
- N#CC1=CC(=CC=C1C2=CC=CC=C2OCC3=CC=CC=C3N4C=CC(=CC=C4)N)C[C@H](N)C


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	1
			27	22	4	1		



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	1
			27	22	4	1		

- Molecule 5 is 3-[(2-amino-4-methylquinolin-7-yl)methoxy]-5-[(2S)-2-(methylamino)propyl]benzonitrile (three-letter code: 8FD) (formula: C₂₂H₂₄N₄O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	1
			27	22	4	1		
5	B	1	Total	C	N	O	0	1
			27	22	4	1		

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	175	Total	O	0	0
			175	175		
8	B	231	Total	O	0	0
			231	231		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.97Å 111.94Å 164.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.23 – 1.82 39.23 – 1.82	Depositor EDS
% Data completeness (in resolution range)	91.6 (39.23-1.82) 92.5 (39.23-1.82)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.82Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.193 , 0.238 0.194 , 0.236	Depositor DCC
R_{free} test set	3997 reflections (4.96%)	DCC
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.551	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 61.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7354	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, H4B, M4R, 8FD, ACT, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/3456	0.51	0/4688
1	B	0.40	0/3459	0.51	0/4689
All	All	0.39	0/6915	0.51	0/9377

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3354	0	3264	27	0
1	B	3357	0	3275	18	0
2	A	43	0	30	2	0
2	B	43	0	30	3	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	27	0	0	1	0
4	B	27	0	0	1	0
5	A	27	0	0	1	0
5	B	27	0	0	1	0
6	A	4	0	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	4	0	3	0	0
7	A	1	0	0	0	0
8	A	175	0	0	1	0
8	B	231	0	0	3	0
All	All	7354	0	6635	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:801:HEM:HHC	2:A:801:HEM:HBB2	1.69	0.74
2:B:801:HEM:HBB2	2:B:801:HEM:HHC	1.69	0.72
2:B:801:HEM:HBC2	2:B:801:HEM:HMC2	1.74	0.69
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.75	0.69
1:A:516:ARG:NH2	8:A:901:HOH:O	2.25	0.67
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.78	0.66
1:B:706:TYR:OH	2:B:801:HEM:O2D	2.11	0.65
1:A:467:ASP:OD1	1:A:467:ASP:N	2.28	0.64
1:A:706:TYR:OH	2:A:801:HEM:O2D	2.17	0.61
1:B:667:ARG:NH1	1:B:668:CYS:HB3	2.20	0.56
1:B:516:ARG:NH1	8:B:907:HOH:O	2.40	0.54
1:A:491:SER:OG	1:A:492:THR:N	2.39	0.54
1:A:660:LYS:O	1:A:663[B]:GLU:HG3	2.08	0.52
1:B:607:LEU:HD13	1:B:626:LYS:HG2	1.93	0.50
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.94	0.50
1:B:595:VAL:O	1:B:599:CYS:HB2	2.12	0.49
1:A:330:ILE:HD11	1:B:696:LEU:HD22	1.94	0.49
1:B:600:ASP:OD2	8:B:901:HOH:O	2.20	0.49
1:B:567:VAL:HG21	4:B:803[A]:M4R:C07	2.43	0.48
1:B:508:GLN:NE2	8:B:910:HOH:O	2.47	0.47
1:B:336:MET:HE2	1:B:336:MET:HB2	1.82	0.47
1:B:322:LEU:HD13	1:B:699:ARG:NH2	2.30	0.46
1:A:359:LYS:HE2	1:A:381:LEU:HD21	1.98	0.46
1:A:595:VAL:O	1:A:599:CYS:HB2	2.16	0.46
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.98	0.45
1:A:496:PRO:HA	1:A:499:VAL:HG23	1.97	0.45
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.99	0.45
1:B:567:VAL:HG21	5:B:804[B]:8FD:C07	2.46	0.45
1:B:323:GLU:O	1:B:699:ARG:HD3	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:553:TRP:HB2	1:A:613:LYS:NZ	2.33	0.44
1:A:546:ILE:HG22	1:A:554:PHE:HE2	1.83	0.44
1:A:567:VAL:HG21	4:A:803[A]:M4R:C07	2.48	0.44
1:A:567:VAL:HG21	5:A:804[B]:8FD:C07	2.48	0.43
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.53	0.43
1:A:360:GLU:O	1:A:364:GLN:HG2	2.18	0.43
1:A:475:TRP:CE2	1:A:710:PRO:HB2	2.53	0.43
1:A:684:SER:HB3	1:A:687:ILE:HD11	2.00	0.43
1:B:366:TYR:HA	1:B:369:ILE:HG12	2.01	0.42
1:A:475:TRP:CZ2	1:A:531:PRO:HG3	2.54	0.42
1:A:596:ARG:NH2	1:A:600:ASP:OD2	2.51	0.42
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.08	0.41
1:A:546:ILE:HG12	1:A:560:LYS:HA	2.03	0.41
1:A:709:ASP:HA	1:A:710:PRO:HD3	1.87	0.41
1:A:328:GLU:H	1:A:328:GLU:HG3	1.62	0.41
1:A:537:PRO:HA	1:A:538:PRO:HD3	1.93	0.41
1:B:701:THR:HA	1:B:702:PRO:C	2.41	0.40
1:A:686:SER:HA	1:A:691:PHE:CG	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/422 (97%)	391 (96%)	18 (4%)	0	100	100
1	B	410/422 (97%)	399 (97%)	11 (3%)	0	100	100
All	All	819/844 (97%)	790 (96%)	29 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/377 (98%)	359 (97%)	10 (3%)	50	34
1	B	369/377 (98%)	358 (97%)	11 (3%)	46	29
All	All	738/754 (98%)	717 (97%)	21 (3%)	48	33

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	348	VAL
1	A	467	ASP
1	A	486	LYS
1	A	503	GLU
1	A	516	ARG
1	A	523	LEU
1	A	527	ASN
1	A	547	ARG
1	A	613	LYS
1	A	715	VAL
1	B	321	THR
1	B	348	VAL
1	B	351	LYS
1	B	353	GLN
1	B	360	GLU
1	B	389	GLU
1	B	392	SER
1	B	486	LYS
1	B	516	ARG
1	B	547	ARG
1	B	667	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	801	1	28,50,50	2.33	9 (32%)	17,82,82	1.33	1 (5%)
3	H4B	A	802	-	14,18,18	0.84	0	12,26,26	2.48	5 (41%)
4	M4R	A	803[A]	-	29,29,29	1.51	2 (6%)	37,40,40	1.08	3 (8%)
5	8FD	A	804[B]	-	29,29,29	1.50	2 (6%)	37,40,40	1.20	4 (10%)
6	ACT	A	805	-	1,3,3	1.32	0	0,3,3	0.00	-
2	HEM	B	801	1	28,50,50	2.32	12 (42%)	17,82,82	1.73	3 (17%)
3	H4B	B	802	-	14,18,18	0.75	0	12,26,26	2.33	4 (33%)
4	M4R	B	803[A]	-	29,29,29	1.53	3 (10%)	37,40,40	1.01	2 (5%)
5	8FD	B	804[B]	-	29,29,29	1.53	2 (6%)	37,40,40	1.07	3 (8%)
6	ACT	B	805	-	1,3,3	1.16	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	801	1	-	0/6/54/54	0/0/8/8
3	H4B	A	802	-	-	0/8/17/17	0/2/2/2
4	M4R	A	803[A]	-	-	0/13/13/13	0/3/3/3
5	8FD	A	804[B]	-	-	0/13/13/13	0/3/3/3
6	ACT	A	805	-	-	0/0/0/0	0/0/0/0
2	HEM	B	801	1	-	0/6/54/54	0/0/8/8
3	H4B	B	802	-	-	0/8/17/17	0/2/2/2
4	M4R	B	803[A]	-	-	0/13/13/13	0/3/3/3
5	8FD	B	804[B]	-	-	0/13/13/13	0/3/3/3
6	ACT	B	805	-	-	0/0/0/0	0/0/0/0

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	804[B]	8FD	C23-C27	-6.50	1.28	1.44
4	B	803[A]	M4R	C23-C27	-6.49	1.28	1.44
5	A	804[B]	8FD	C23-C27	-6.46	1.29	1.44
4	A	803[A]	M4R	C23-C27	-6.38	1.29	1.44
2	A	801	HEM	C3B-C2B	-5.33	1.33	1.40
2	B	801	HEM	C3B-C2B	-4.98	1.33	1.40
2	B	801	HEM	C3C-C2C	-3.93	1.35	1.40
2	A	801	HEM	C3C-C2C	-3.87	1.35	1.40
5	B	804[B]	8FD	C05-C10	-2.12	1.39	1.42
4	B	803[A]	M4R	C05-C10	-2.11	1.39	1.42
4	B	803[A]	M4R	C09-C10	-2.07	1.38	1.41
2	A	801	HEM	CMC-C2C	2.00	1.55	1.51
2	A	801	HEM	CMD-C2D	2.06	1.55	1.51
2	B	801	HEM	C1D-ND	2.07	1.40	1.36
2	B	801	HEM	CMD-C2D	2.08	1.55	1.51
2	B	801	HEM	CMB-C2B	2.10	1.56	1.51
2	B	801	HEM	CMC-C2C	2.10	1.56	1.51
2	B	801	HEM	C4B-NB	2.10	1.40	1.36
5	A	804[B]	8FD	C02-N01	2.34	1.35	1.33
2	A	801	HEM	C1D-ND	2.45	1.41	1.36
4	A	803[A]	M4R	C02-N01	2.48	1.35	1.33
2	B	801	HEM	C4D-ND	2.67	1.39	1.36
2	B	801	HEM	C1B-NB	2.98	1.40	1.36
2	A	801	HEM	C4D-ND	3.71	1.41	1.36
2	B	801	HEM	C3B-CAB	3.71	1.55	1.47
2	A	801	HEM	C3C-CAC	3.72	1.55	1.47
2	B	801	HEM	C3C-CAC	3.79	1.55	1.47
2	A	801	HEM	C3B-CAB	4.01	1.55	1.47
2	A	801	HEM	C3D-C2D	5.31	1.53	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	HEM	C3D-C2D	5.37	1.53	1.37

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	H4B	N3-C2-N1	-3.03	120.54	125.45
2	B	801	HEM	C1D-C2D-C3D	-2.99	104.92	107.00
2	B	801	HEM	CBA-CAA-C2A	-2.91	106.92	112.48
3	B	802	H4B	N3-C2-N1	-2.84	120.84	125.45
5	A	804[B]	8FD	C05-C10-N01	-2.82	119.92	122.84
2	B	801	HEM	CAD-CBD-CGD	-2.74	107.97	112.66
2	A	801	HEM	CAD-CBD-CGD	-2.57	108.28	112.66
4	A	803[A]	M4R	C05-C10-N01	-2.43	120.33	122.84
5	B	804[B]	8FD	C05-C10-N01	-2.32	120.44	122.84
4	B	803[A]	M4R	C05-C10-N01	-2.18	120.59	122.84
3	A	802	H4B	N2-C2-N3	2.13	120.64	117.24
5	B	804[B]	8FD	N02-C02-N01	2.15	119.39	118.06
4	A	803[A]	M4R	N02-C02-N01	2.25	119.45	118.06
5	A	804[B]	8FD	C12-O13-C21	2.38	123.54	117.61
4	B	803[A]	M4R	C04-C05-C10	2.61	119.47	117.99
5	A	804[B]	8FD	N02-C02-N01	2.65	119.70	118.06
5	B	804[B]	8FD	C04-C05-C10	2.75	119.55	117.99
3	B	802	H4B	C4-N3-C2	2.86	120.17	116.06
3	B	802	H4B	C2-N1-C8A	2.87	120.97	114.51
3	A	802	H4B	C4-N3-C2	2.98	120.35	116.06
3	A	802	H4B	C2-N1-C8A	3.00	121.28	114.51
4	A	803[A]	M4R	C04-C05-C10	3.01	119.69	117.99
5	A	804[B]	8FD	C04-C05-C10	3.19	119.79	117.99
3	B	802	H4B	C4-C4A-C8A	5.33	119.39	114.56
3	A	802	H4B	C4-C4A-C8A	5.64	119.67	114.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	HEM	2	0
4	A	803[A]	M4R	1	0
5	A	804[B]	8FD	1	0
2	B	801	HEM	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	803[A]	M4R	1	0
5	B	804[B]	8FD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	410/422 (97%)	1.06	82 (20%) 1 1	30, 62, 115, 162	0
1	B	411/422 (97%)	0.44	37 (9%) 10 8	30, 51, 87, 117	0
All	All	821/844 (97%)	0.75	119 (14%) 3 2	30, 56, 108, 162	0

All (119) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	715	VAL	9.5
1	A	716	TRP	8.5
1	A	486	LYS	7.9
1	A	488	PRO	7.3
1	A	348	VAL	7.0
1	A	347	ASP	7.0
1	B	300	PHE	6.6
1	A	355	PHE	5.9
1	A	507	GLN	5.7
1	A	388	ILE	5.3
1	A	489	ASP	5.2
1	B	350	THR	4.9
1	A	352	ASP	4.9
1	A	551	PHE	4.8
1	A	351	LYS	4.8
1	A	506	ILE	4.8
1	A	713	THR	4.7
1	A	711	TRP	4.7
1	A	299	ARG	4.6
1	A	349	ARG	4.5
1	A	391	THR	4.5
1	A	712	ASN	4.4
1	B	677	VAL	4.3
1	A	508	GLN	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	718	GLY	4.2
1	A	678	TRP	4.2
1	B	619	ARG	4.1
1	A	503	GLU	4.1
1	A	389	GLU	4.1
1	A	714	HIS	4.0
1	A	677	VAL	3.9
1	B	348	VAL	3.9
1	B	351	LYS	3.9
1	A	679	ILE	3.8
1	A	643	SER	3.8
1	A	386	LYS	3.8
1	B	680	VAL	3.7
1	B	678	TRP	3.6
1	A	470	HIS	3.6
1	A	467	ASP	3.5
1	A	567	VAL	3.5
1	A	390	SER	3.5
1	A	322	LEU	3.4
1	A	681	PRO	3.4
1	A	511	LYS	3.4
1	A	393	THR	3.4
1	A	676	TRP	3.3
1	A	499	VAL	3.3
1	A	553	TRP	3.3
1	B	301	LEU	3.3
1	B	299	ARG	3.3
1	B	352	ASP	3.2
1	B	617	ASP	3.2
1	A	392	SER	3.2
1	A	504	ILE	3.1
1	A	480	ILE	3.1
1	A	667	ARG	3.0
1	A	373	GLY	3.0
1	A	619	ARG	2.9
1	A	505	CYS	2.9
1	B	329	HIS	2.9
1	A	321	THR	2.8
1	A	350	THR	2.8
1	A	516	ARG	2.8
1	B	676	TRP	2.8
1	A	487	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	682	PRO	2.8
1	B	593	ILE	2.8
1	A	300	PHE	2.8
1	A	369	ILE	2.8
1	A	588	TYR	2.8
1	A	685	GLY	2.7
1	A	566	ALA	2.7
1	A	691	PHE	2.7
1	B	682	PRO	2.7
1	A	492	THR	2.7
1	B	611	ALA	2.6
1	B	311	VAL	2.6
1	B	561	TRP	2.6
1	A	584	PHE	2.6
1	B	616	LEU	2.6
1	B	479	LEU	2.6
1	A	479	LEU	2.5
1	A	311	VAL	2.5
1	B	490	GLY	2.5
1	B	567	VAL	2.5
1	A	481	ARG	2.4
1	A	490	GLY	2.4
1	A	385	ASN	2.4
1	B	620	LYS	2.4
1	A	609	GLU	2.4
1	B	302	LYS	2.4
1	B	679	ILE	2.4
1	A	611	ALA	2.3
1	B	309	ASP	2.3
1	B	551	PHE	2.3
1	A	686	SER	2.3
1	B	691	PHE	2.3
1	B	310	VAL	2.3
1	A	565	PRO	2.3
1	A	469	LYS	2.2
1	B	375	LYS	2.2
1	B	318	LEU	2.2
1	A	415	CYS	2.2
1	B	612	LYS	2.2
1	B	313	THR	2.2
1	A	617	ASP	2.1
1	B	566	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	561	TRP	2.1
1	A	466	THR	2.1
1	A	591	THR	2.1
1	A	552	ASP	2.1
1	A	515	GLY	2.1
1	B	562	TYR	2.0
1	A	680	VAL	2.0
1	A	394	TYR	2.0
1	A	328	GLU	2.0
1	A	593	ILE	2.0
1	A	443	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	ACT	A	805	4/4	0.80	0.22	6.13	56,62,63,63	0
6	ACT	B	805	4/4	0.95	0.19	5.26	49,56,57,58	0
4	M4R	B	803[A]	27/27	0.91	0.33	3.85	32,52,83,85	27
5	8FD	B	804[B]	27/27	0.90	0.33	3.76	32,52,82,84	27
4	M4R	A	803[A]	27/27	0.88	0.35	3.50	32,56,84,88	27
5	8FD	A	804[B]	27/27	0.89	0.34	3.29	33,54,84,87	27
2	HEM	A	801	43/43	0.97	0.22	1.13	26,40,71,74	0
2	HEM	B	801	43/43	0.97	0.18	0.82	28,40,62,65	0
3	H4B	B	802	17/17	0.89	0.19	0.19	36,45,52,53	0
3	H4B	A	802	17/17	0.92	0.18	0.16	32,47,59,65	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	ZN	A	806	1/1	0.99	0.08	-1.22	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.