



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 01:39 am GMT

PDB ID : 4UP8  
Title : Crystal structure of Entamoeba histolytica lysyl-tRNA synthetase apo form  
Authors : Bonnefond, L.; Nureki, O.  
Deposited on : 2014-06-14  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949

**i**

## X-RAY DIFFRACTION

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R <sub>free</sub>	100719	1586 (2.90-2.90)
Clashscore	112137	1807 (2.90-2.90)
Ramachandran outliers	110173	1768 (2.90-2.90)
Sidechain outliers	110143	1770 (2.90-2.90)
RSRZ outliers	101464	1596 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	777	<div> <div></div> <div>63%</div> <div>10%</div> <div>25%</div> </div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4448 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LYSINE-TRNA LIGASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	580	Total	C	N	O	S	0	0	0
			4448	2827	766	830	25			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	EXPRESSION TAG	UNP C4M7X2
A	-6	HIS	-	EXPRESSION TAG	UNP C4M7X2
A	-5	HIS	-	EXPRESSION TAG	UNP C4M7X2
A	-4	HIS	-	EXPRESSION TAG	UNP C4M7X2
A	-3	HIS	-	EXPRESSION TAG	UNP C4M7X2
A	-2	HIS	-	EXPRESSION TAG	UNP C4M7X2
A	-1	HIS	-	EXPRESSION TAG	UNP C4M7X2
A	0	MET	-	EXPRESSION TAG	UNP C4M7X2
A	1	LEU	-	EXPRESSION TAG	UNP C4M7X2
A	582	LYS	LEU	CONFLICT	UNP C4M7X2
A	764	ILE	VAL	ENGINEERED MUTATION	UNP C4M7X2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	156.48Å 156.48Å 93.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.54 – 2.90 34.54 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (34.54-2.90) 99.8 (34.54-2.90)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.11 (at 2.90Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.216 , 0.251 0.216 , 0.251	Depositor DCC
$R_{free}$ test set	1405 reflections (4.83%)	DCC
Wilson B-factor (Å <sup>2</sup> )	83.0	Xtriage
Anisotropy	0.130	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 52.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.033 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4448	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/4534	0.55	5/6137 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	327	PRO	N-CA-CB	6.20	110.74	103.30
1	A	329	PRO	N-CA-CB	5.94	110.43	103.30
1	A	325	PRO	N-CA-CB	5.77	110.22	103.30
1	A	432	LYS	CB-CG-CD	5.13	124.95	111.60
1	A	431	ASP	C-N-CA	-5.09	108.97	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	398	ARG	Peptide
1	A	432	LYS	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4448	0	4285	59	0
All	All	4448	0	4285	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:560:ARG:NH1	1:A:561:PRO:O	2.10	0.84
1:A:422:GLU:HG2	1:A:424:GLN:HG3	1.65	0.79
1:A:142:PRO:HG2	1:A:164:VAL:HG11	1.67	0.77
1:A:9:ARG:HG2	1:A:138:LEU:HA	1.67	0.76
1:A:574:GLU:O	1:A:577:MET:HB2	1.93	0.69
1:A:431:ASP:O	1:A:434:PHE:HB3	1.92	0.69
1:A:432:LYS:HA	1:A:435:GLY:H	1.59	0.67
1:A:258:THR:HG23	1:A:259:HIS:HD2	1.62	0.65
1:A:570:ARG:O	1:A:574:GLU:HG3	1.97	0.64
1:A:420:CYS:SG	1:A:421:ALA:N	2.70	0.64
1:A:433:LEU:O	1:A:437:LEU:HB2	2.00	0.62
1:A:392:PHE:O	1:A:415:ARG:NH2	2.33	0.62
1:A:391:LYS:HZ3	1:A:437:LEU:HD11	1.66	0.61
1:A:301:GLU:OE1	1:A:302:ILE:N	2.26	0.60
1:A:577:MET:O	1:A:580:GLN:HB3	2.01	0.60
1:A:11:LYS:O	1:A:15:GLU:HG3	2.01	0.60
1:A:2:SER:OG	1:A:575:ASP:OD1	2.23	0.57
1:A:258:THR:HG23	1:A:259:HIS:CD2	2.40	0.56
1:A:398:ARG:HH21	1:A:466:PRO:HG2	1.70	0.56
1:A:560:ARG:HD2	1:A:561:PRO:HD2	1.87	0.56
1:A:156:ARG:NH2	1:A:560:ARG:O	2.36	0.55
1:A:435:GLY:HA2	1:A:439:GLU:HB3	1.89	0.55
1:A:406:LEU:O	1:A:410:LYS:HD2	2.08	0.54
1:A:294:THR:OG1	1:A:295:GLY:N	2.41	0.54
1:A:406:LEU:HG	1:A:410:LYS:NZ	2.24	0.53
1:A:563:ASP:OD1	1:A:566:GLU:HG3	2.09	0.52
1:A:421:ALA:O	1:A:424:GLN:HG2	2.10	0.52
1:A:383:SER:HB3	1:A:386:HIS:CD2	2.45	0.51
1:A:417:ASN:OD1	1:A:419:ILE:N	2.23	0.51
1:A:60:VAL:HG13	1:A:73:ILE:HG23	1.92	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:406:LEU:HD13	1:A:426:THR:HA	1.95	0.49
1:A:62:SER:HB3	1:A:74:ASP:HB2	1.93	0.49
1:A:391:LYS:NZ	1:A:437:LEU:HD11	2.26	0.49
1:A:383:SER:HB3	1:A:386:HIS:HD2	1.78	0.49
1:A:406:LEU:HG	1:A:410:LYS:HZ2	1.77	0.48
1:A:411:LYS:O	1:A:414:ILE:HG13	2.14	0.47
1:A:472:PHE:CE2	1:A:484:ALA:HB3	2.49	0.47
1:A:437:LEU:HD12	1:A:437:LEU:HA	1.74	0.47
1:A:412:GLN:HA	1:A:415:ARG:HG2	1.97	0.46
1:A:435:GLY:HA2	1:A:439:GLU:CB	2.45	0.46
1:A:432:LYS:CG	1:A:435:GLY:HA3	2.45	0.46
1:A:301:GLU:CD	1:A:302:ILE:H	2.15	0.46
1:A:417:ASN:OD1	1:A:418:ALA:N	2.49	0.45
1:A:432:LYS:HG3	1:A:435:GLY:HA3	1.98	0.45
1:A:514:LEU:HD12	1:A:515:VAL:H	1.82	0.45
1:A:279:MET:HG3	1:A:472:PHE:CE2	2.52	0.44
1:A:456:SER:HB2	1:A:469:THR:HG21	1.99	0.44
1:A:560:ARG:HH11	1:A:560:ARG:HG3	1.83	0.44
1:A:411:LYS:O	1:A:415:ARG:HB3	2.17	0.44
1:A:451:GLN:OE1	1:A:451:GLN:N	2.49	0.44
1:A:175:ARG:O	1:A:178:VAL:HB	2.19	0.43
1:A:-1:HIS:O	1:A:3:LYS:HG3	2.20	0.41
1:A:495:ARG:O	1:A:499:GLU:HG3	2.20	0.41
1:A:47:SER:N	1:A:118:ARG:O	2.53	0.41
1:A:347:ASN:HA	1:A:348:TYR:HA	1.94	0.41
1:A:301:GLU:CD	1:A:302:ILE:N	2.73	0.40
1:A:307:MET:O	1:A:309:ILE:N	2.51	0.40
1:A:144:MET:HE2	1:A:164:VAL:HA	2.04	0.40
1:A:425:THR:O	1:A:429:VAL:HG12	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	576/777 (74%)	535 (93%)	31 (5%)	10 (2%)	11	36

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	325	PRO
1	A	327	PRO
1	A	399	PRO
1	A	309	ILE
1	A	323	LYS
1	A	329	PRO
1	A	328	ARG
1	A	330	PHE
1	A	331	ASN
1	A	398	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	454/679 (67%)	449 (99%)	5 (1%)	78	94

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	258	THR
1	A	279	MET
1	A	301	GLU
1	A	438	ILE
1	A	564	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	259	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	386	HIS
1	A	483	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	580/777 (74%)	0.01	11 (1%) 67 64	44, 76, 139, 164	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	583	ALA	6.0
1	A	414	ILE	3.6
1	A	367	THR	3.1
1	A	350	TYR	2.9
1	A	329	PRO	2.8
1	A	364	ASP	2.7
1	A	322	PHE	2.7
1	A	419	ILE	2.5
1	A	306	LEU	2.4
1	A	346	LEU	2.4
1	A	349	TYR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.