



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 07:31 am GMT

PDB ID : 3UQX
Title : Crystal structure of BACE1 with its inhibitor
Authors : Chen, T.T.; Chen, W.Y.; Xu, Y.C.
Deposited on : 2011-11-21
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3453 (1.70-1.70)
Clashscore	112137	3876 (1.70-1.70)
Ramachandran outliers	110173	3815 (1.70-1.70)
Sidechain outliers	110143	3815 (1.70-1.70)
RSRZ outliers	101464	3491 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	<div> <div>3%</div> <div> <div></div> <div>74%</div> <div>11%</div> <div>13%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	404	-	-	-	X
4	URE	A	407	-	X	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	URE	A	408	-	-	-	X
4	URE	A	409	-	-	-	X
4	URE	A	410	-	X	-	-
5	CL	A	412	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3357 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-secretase 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	375	2944	1893	481	556	14	0	8	0

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-39	MET	-	EXPRESSION TAG	UNP P56817
A	-38	GLY	-	EXPRESSION TAG	UNP P56817
A	-37	SER	-	EXPRESSION TAG	UNP P56817
A	-36	SER	-	EXPRESSION TAG	UNP P56817
A	-35	HIS	-	EXPRESSION TAG	UNP P56817
A	-34	HIS	-	EXPRESSION TAG	UNP P56817
A	-33	HIS	-	EXPRESSION TAG	UNP P56817
A	-32	HIS	-	EXPRESSION TAG	UNP P56817
A	-31	HIS	-	EXPRESSION TAG	UNP P56817
A	-30	HIS	-	EXPRESSION TAG	UNP P56817
A	-29	SER	-	EXPRESSION TAG	UNP P56817
A	-28	ALA	-	EXPRESSION TAG	UNP P56817
A	-27	GLY	-	EXPRESSION TAG	UNP P56817
A	-26	GLU	-	EXPRESSION TAG	UNP P56817
A	-25	ASN	-	EXPRESSION TAG	UNP P56817
A	-24	LEU	-	EXPRESSION TAG	UNP P56817
A	-23	TYR	-	EXPRESSION TAG	UNP P56817
A	-22	PHE	-	EXPRESSION TAG	UNP P56817
A	-21	GLN	-	EXPRESSION TAG	UNP P56817
A	-20	GLY	-	EXPRESSION TAG	UNP P56817
A	-19	THR	-	EXPRESSION TAG	UNP P56817
A	75	ALA	LYS	ENGINEERED MUTATION	UNP P56817
A	77	ALA	GLU	ENGINEERED MUTATION	UNP P56817


- Molecule 2 is N-[(1R)-1-(4-FLUOROPHENYL)ETHYL]-N'-[(2S,3S)-3-HYDROXY-4-{4-[(1S)-1-HYDROXYETHYL]-1H-1,2,3-TRIAZOL-1-YL}-1-PHENYLBUTAN-2-YL]-5-[METHYL(METHYLSULFONYL)AMINO]BENZENE-1,3-DICARBOXAMIDE (three-letter code:

ZPZ) (formula: $\text{C}_{32}\text{H}_{37}\text{FN}_6\text{O}_6\text{S}$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	0	0
			46	32	1	6	6	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is UREA (three-letter code: URE) (formula: $\text{CH}_4\text{N}_2\text{O}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			4	1	2	1		
4	A	1	Total	C	N	O	0	0
			4	1	2	1		
4	A	1	Total	C	N	O	0	0
			4	1	2	1		
4	A	1	Total	C	N	O	0	0
			4	1	2	1		
4	A	1	Total	C	N	O	0	0
			4	1	2	1		
4	A	1	Total	C	N	O	0	0
			4	1	2	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Cl	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	322	Total 322	O 322	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-secretase 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	104.55Å 128.28Å 76.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.80 – 1.70 35.80 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.9 (35.80-1.70) 98.9 (35.80-1.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.29 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.159 , 0.186 0.159 , 0.186	Depositor DCC
R_{free} test set	2842 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.548	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3357	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: URE, ZPZ, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.57	17/3042 (0.6%)	1.50	28/4140 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	380	GLU	CD-OE1	7.97	1.34	1.25
1	A	255	GLU	CD-OE1	6.90	1.33	1.25
1	A	79	GLU	CD-OE1	6.19	1.32	1.25
1	A	262	TRP	CG-CD1	6.14	1.45	1.36
1	A	104	GLU	CD-OE1	6.08	1.32	1.25
1	A	120	GLY	N-CA	6.08	1.55	1.46
1	A	196	GLU	CD-OE1	5.71	1.31	1.25
1	A	100	ALA	C-O	5.68	1.34	1.23
1	A	290	GLU	CD-OE2	5.59	1.31	1.25
1	A	76	TRP	CD2-CE2	5.53	1.48	1.41
1	A	116	GLU	CD-OE2	5.48	1.31	1.25
1	A	49	HIS	CB-CG	5.37	1.59	1.50
1	A	204	VAL	C-O	5.37	1.33	1.23
1	A	189	TRP	CD2-CE3	5.22	1.48	1.40
1	A	-3	GLY	N-CA	5.06	1.53	1.46
1	A	295	SER	CA-CB	5.05	1.60	1.52
1	A	76	TRP	CZ3-CH2	5.05	1.48	1.40

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	50	ARG	NE-CZ-NH2	-12.68	113.96	120.30
1	A	50	ARG	NE-CZ-NH1	10.03	125.31	120.30
1	A	130	ASP	CB-CG-OD2	9.54	126.89	118.30
1	A	350	LYS	CD-CE-NZ	9.47	133.48	111.70
1	A	349	ARG	NE-CZ-NH1	9.09	124.84	120.30
1	A	349	ARG	NE-CZ-NH2	-8.66	115.97	120.30
1	A	128	ARG	NE-CZ-NH1	8.44	124.52	120.30
1	A	212	ASP	CB-CG-OD1	8.33	125.80	118.30
1	A	15	TYR	CB-CG-CD1	-6.92	116.85	121.00
1	A	60	TYR	CG-CD2-CE2	-6.79	115.87	121.30
1	A	296	PHE	CB-CG-CD1	-6.71	116.10	120.80
1	A	349	ARG	CG-CD-NE	-6.69	97.75	111.80
1	A	27	LEU	CB-CG-CD1	-5.99	100.82	111.00
1	A	61	ARG	NE-CZ-NH2	-5.99	117.31	120.30
1	A	4[A]	ASP	CB-CG-OD1	5.97	123.67	118.30
1	A	4[B]	ASP	CB-CG-OD1	5.97	123.67	118.30
1	A	194	ARG	NE-CZ-NH2	-5.87	117.37	120.30
1	A	14	TYR	CB-CG-CD1	-5.85	117.49	121.00
1	A	96	ARG	NE-CZ-NH2	-5.65	117.48	120.30
1	A	255	GLU	CG-CD-OE1	5.65	129.59	118.30
1	A	62	ASP	CB-CG-OD2	-5.57	113.29	118.30
1	A	297	ARG	NE-CZ-NH2	-5.51	117.54	120.30
1	A	306	LEU	CB-CG-CD1	-5.48	101.69	111.00
1	A	50	ARG	CD-NE-CZ	5.45	131.23	123.60
1	A	198	TYR	CB-CG-CD1	-5.33	117.80	121.00
1	A	128	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	A	333	MET	CG-SD-CE	-5.16	91.94	100.20
1	A	54	ARG	NE-CZ-NH1	-5.07	117.77	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	197	TRP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2944	0	2840	16	0
2	A	46	0	36	1	0
3	A	15	0	0	0	0
4	A	28	0	28	3	0
5	A	2	0	0	1	0
6	A	322	0	0	4	0
All	All	3357	0	2904	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:364:GLU:HG2	6:A:750:HOH:O	1.64	0.98
1:A:128:ARG:HH11	4:A:407:URE:HN22	1.10	0.92
1:A:8:GLY:HA3	1:A:170:VAL:HG12	1.60	0.84
1:A:8:GLY:HA3	1:A:170:VAL:CG1	2.17	0.74
1:A:8:GLY:CA	1:A:170:VAL:HG12	2.26	0.65
1:A:301:LEU:H	1:A:304:GLN:HE21	1.47	0.60
1:A:128:ARG:NH1	4:A:407:URE:HN22	1.92	0.58
1:A:301:LEU:H	1:A:304:GLN:NE2	2.03	0.56
1:A:290:GLU:OE1	6:A:669:HOH:O	2.19	0.54
1:A:371:GLU:HG2	6:A:788:HOH:O	2.10	0.51
1:A:218:LYS:HE3	1:A:381:ASP:O	2.12	0.50
4:A:410:URE:N1	6:A:775:HOH:O	2.35	0.50
1:A:34:GLY:O	2:A:401:ZPZ:H32	2.15	0.45
1:A:222:TYR:HA	1:A:223:ASP:HA	1.85	0.45
1:A:350:LYS:HB3	1:A:350:LYS:HE3	1.73	0.44
1:A:194:ARG:HD2	1:A:202:ILE:HD11	2.00	0.42
1:A:23:PRO:HA	5:A:412:CL:CL	2.58	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	377/433 (87%)	370 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/368 (86%)	311 (99%)	4 (1%)	73	60

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	SER
1	A	111	ASN
1	A	197	TRP
1	A	315	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	28	ASN
1	A	111	ASN
1	A	278	ASN
1	A	304	GLN
1	A	362	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ZPZ	A	401	-	46,49,49	2.33	16 (34%)	59,70,70	2.84	24 (40%)
3	SO4	A	402	-	4,4,4	0.69	0	6,6,6	0.72	0
3	SO4	A	403	-	4,4,4	0.34	0	6,6,6	0.51	0
3	SO4	A	404	-	4,4,4	0.71	0	6,6,6	0.64	0
4	URE	A	405	-	3,3,3	0.11	0	3,3,3	1.03	0
4	URE	A	406	-	3,3,3	0.61	0	3,3,3	2.75	2 (66%)
4	URE	A	407	-	3,3,3	1.95	1 (33%)	3,3,3	2.21	2 (66%)
4	URE	A	408	-	3,3,3	0.71	0	3,3,3	2.13	1 (33%)
4	URE	A	409	-	3,3,3	0.97	0	3,3,3	0.27	0
4	URE	A	410	-	3,3,3	1.64	0	3,3,3	3.55	3 (100%)
4	URE	A	411	-	3,3,3	0.54	0	3,3,3	1.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ZPZ	A	401	-	-	0/42/46/46	0/4/4/4
3	SO4	A	402	-	-	0/0/0/0	0/0/0/0
3	SO4	A	403	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	404	-	-	0/0/0/0	0/0/0/0
4	URE	A	405	-	-	0/0/0/0	0/0/0/0
4	URE	A	406	-	-	0/0/0/0	0/0/0/0
4	URE	A	407	-	-	0/0/0/0	0/0/0/0
4	URE	A	408	-	-	0/0/0/0	0/0/0/0
4	URE	A	409	-	-	0/0/0/0	0/0/0/0
4	URE	A	410	-	-	0/0/0/0	0/0/0/0
4	URE	A	411	-	-	0/0/0/0	0/0/0/0

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	ZPZ	CBJ-CBF	-3.37	1.43	1.50
2	A	401	ZPZ	CBQ-NBD	-2.61	1.41	1.46
4	A	407	URE	O-C	-2.38	1.16	1.26
2	A	401	ZPZ	CAS-CBK	2.33	1.42	1.39
2	A	401	ZPZ	CAV-CBI	2.40	1.43	1.39
2	A	401	ZPZ	CBE-NBC	2.45	1.39	1.34
2	A	401	ZPZ	CAQ-CAS	2.48	1.43	1.38
2	A	401	ZPZ	CAC-NBR	2.60	1.51	1.47
2	A	401	ZPZ	CAU-CBJ	2.63	1.43	1.39
2	A	401	ZPZ	CAW-CBJ	3.87	1.45	1.39
2	A	401	ZPZ	CBL-NBR	4.03	1.51	1.44
2	A	401	ZPZ	OAG-SBT	4.07	1.50	1.43
2	A	401	ZPZ	CAV-CBL	4.09	1.47	1.39
2	A	401	ZPZ	CBF-NBD	4.33	1.44	1.34
2	A	401	ZPZ	CAP-CBH	4.51	1.48	1.38
2	A	401	ZPZ	CAT-CBK	4.53	1.46	1.39
2	A	401	ZPZ	CAM-CAO	5.53	1.49	1.38

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	ZPZ	CBI-CAV-CBL	-9.43	109.83	119.70
2	A	401	ZPZ	CAR-CAT-CBK	-8.13	112.95	121.20
2	A	401	ZPZ	CAQ-CAS-CBK	-5.64	115.48	121.20
2	A	401	ZPZ	CAY-CBQ-NBD	-5.15	104.13	110.17
4	A	406	URE	O-C-N1	-3.51	112.52	121.01
4	A	407	URE	O-C-N1	-3.00	113.75	121.01
2	A	401	ZPZ	CAM-CAO-CBH	-2.84	116.31	120.64
2	A	401	ZPZ	CBJ-CAW-CBL	-2.80	116.78	119.70
2	A	401	ZPZ	CAT-CBK-CBO	-2.76	115.39	120.79

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	ZPZ	OAG-SBT-NBR	-2.65	103.64	107.09
2	A	401	ZPZ	CAX-CBM-CBN	-2.62	125.27	129.30
2	A	401	ZPZ	CAU-CBJ-CAW	-2.59	116.47	119.63
2	A	401	ZPZ	CAS-CAQ-CBG	-2.50	115.72	118.35
4	A	410	URE	O-C-N2	-2.30	115.45	121.01
2	A	401	ZPZ	CAN-CAP-CBH	-2.19	117.30	120.64
2	A	401	ZPZ	CAY-CBH-CAO	-2.16	116.54	120.91
2	A	401	ZPZ	FAK-CBG-CAQ	-2.05	114.95	118.53
4	A	410	URE	O-C-N1	-2.02	116.12	121.01
4	A	407	URE	N2-C-N1	2.08	121.90	117.83
2	A	401	ZPZ	OAH-SBT-NBR	2.53	110.38	107.09
2	A	401	ZPZ	CBI-CAU-CBJ	2.65	123.54	120.46
2	A	401	ZPZ	CBH-CAY-CBQ	2.74	118.33	113.36
2	A	401	ZPZ	CAO-CBH-CAP	2.85	122.69	118.16
4	A	406	URE	N2-C-N1	3.00	123.69	117.83
4	A	408	URE	N2-C-N1	3.15	123.98	117.83
2	A	401	ZPZ	OAI-CBN-CAA	3.36	124.53	108.67
2	A	401	ZPZ	CAR-CBG-CAQ	3.44	127.65	122.86
2	A	401	ZPZ	CAU-CBI-CAV	3.54	123.97	119.63
2	A	401	ZPZ	CAV-CBL-CAW	4.52	129.16	120.02
2	A	401	ZPZ	NBA-NBB-NBS	4.89	111.00	107.31
4	A	410	URE	N2-C-N1	5.34	128.26	117.83
2	A	401	ZPZ	CAT-CBK-CAS	6.99	127.07	118.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	ZPZ	1	0
4	A	407	URE	2	0
4	A	410	URE	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	375/433 (86%)	-0.34	12 (3%)	48 54	6, 12, 34, 56	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	10	SER	3.9
1	A	365	PHE	3.4
1	A	254	THR	3.4
1	A	170	VAL	3.0
1	A	386	ILE	2.8
1	A	273	GLY	2.6
1	A	272	ALA	2.5
1	A	49	HIS	2.4
1	A	253	SER	2.3
1	A	251	ALA	2.3
1	A	279	ILE	2.1
1	A	364	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	URE	A	409	4/4	0.86	0.17	11.41	21,25,31,35	0
5	CL	A	412	1/1	1.00	0.10	8.42	7,7,7,7	1
4	URE	A	408	4/4	0.88	0.13	7.82	28,30,40,41	0
3	SO4	A	404	5/5	0.85	0.14	5.08	26,33,36,36	5
5	CL	A	413	1/1	0.99	0.09	1.66	15,15,15,15	1
4	URE	A	405	4/4	0.96	0.10	1.42	16,18,23,23	0
4	URE	A	407	4/4	0.98	0.08	1.21	13,16,19,19	0
2	ZPZ	A	401	46/46	0.98	0.07	-0.33	7,9,14,27	0
4	URE	A	410	4/4	0.78	0.19	-	29,30,36,42	0
4	URE	A	411	4/4	0.94	0.09	-	31,35,38,43	0
4	URE	A	406	4/4	0.95	0.11	-	13,16,21,28	0
3	SO4	A	403	5/5	0.95	0.28	-	37,40,41,43	5
3	SO4	A	402	5/5	0.99	0.12	-	28,29,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.