



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2017 – 09:47 PM EST

PDB ID : 5UQX  
Title : USP7 in complex with GNE6776 (6'-amino-4'-ethyl-5'-(4-hydroxyphenyl)-N-methyl-[3,3'-bipyridine]-6-carboxamide)  
Authors : Murray, J.M.; Rouge, L.  
Deposited on : unknown  
Resolution : 2.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1804 (2.26-2.22)
Clashscore	112137	1957 (2.26-2.22)
Ramachandran outliers	110173	1916 (2.26-2.22)
Sidechain outliers	110143	1917 (2.26-2.22)
RSRZ outliers	101464	1809 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	<div> <div>8%</div> <div> <div></div> <div>75%</div> <div>14%</div> <div>5%</div> <div>6%</div> </div> </div>
1	B	369	<div> <div>7%</div> <div> <div></div> <div>76%</div> <div>12%</div> <div>•</div> <div>10%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	346	Total	C	N	O	S	0	0	0
			2811	1775	480	540	16			
1	B	332	Total	C	N	O	S	0	0	0
			2697	1709	462	510	16			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	187	MET	-	initiating methionine	UNP Q93009
A	188	GLY	-	expression tag	UNP Q93009
A	189	SER	-	expression tag	UNP Q93009
A	190	SER	-	expression tag	UNP Q93009
A	191	HIS	-	expression tag	UNP Q93009
A	192	HIS	-	expression tag	UNP Q93009
A	193	HIS	-	expression tag	UNP Q93009
A	194	HIS	-	expression tag	UNP Q93009
A	195	HIS	-	expression tag	UNP Q93009
A	196	HIS	-	expression tag	UNP Q93009
A	197	SER	-	expression tag	UNP Q93009
A	198	SER	-	expression tag	UNP Q93009
A	199	GLY	-	expression tag	UNP Q93009
A	200	LEU	-	expression tag	UNP Q93009
A	201	VAL	-	expression tag	UNP Q93009
A	202	PRO	-	expression tag	UNP Q93009
A	203	ARG	-	expression tag	UNP Q93009
A	204	GLY	-	expression tag	UNP Q93009
A	205	SER	-	expression tag	UNP Q93009
A	206	HIS	-	expression tag	UNP Q93009
A	207	MET	-	expression tag	UNP Q93009
B	187	MET	-	initiating methionine	UNP Q93009
B	188	GLY	-	expression tag	UNP Q93009
B	189	SER	-	expression tag	UNP Q93009
B	190	SER	-	expression tag	UNP Q93009

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	191	HIS	-	expression tag	UNP Q93009
B	192	HIS	-	expression tag	UNP Q93009
B	193	HIS	-	expression tag	UNP Q93009
B	194	HIS	-	expression tag	UNP Q93009
B	195	HIS	-	expression tag	UNP Q93009
B	196	HIS	-	expression tag	UNP Q93009
B	197	SER	-	expression tag	UNP Q93009
B	198	SER	-	expression tag	UNP Q93009
B	199	GLY	-	expression tag	UNP Q93009
B	200	LEU	-	expression tag	UNP Q93009
B	201	VAL	-	expression tag	UNP Q93009
B	202	PRO	-	expression tag	UNP Q93009
B	203	ARG	-	expression tag	UNP Q93009
B	204	GLY	-	expression tag	UNP Q93009
B	205	SER	-	expression tag	UNP Q93009
B	206	HIS	-	expression tag	UNP Q93009
B	207	MET	-	expression tag	UNP Q93009

- Molecule 2 is 6'-amino-4'-ethyl-5'-(4-hydroxyphenyl)-N-methyl[3,3'-bipyridine]-6-carboxamide (three-letter code: 8JP) (formula: C<sub>20</sub>H<sub>20</sub>N<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			26	20	4	2		
2	B	1	Total	C	N	O	0	0
			26	20	4	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	133	Total 133	O 133	0	0
3	B	102	Total 102	O 102	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7



- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.36Å 67.53Å 76.86Å 90.00° 96.99° 90.00°	Depositor
Resolution (Å)	38.15 – 2.23 38.14 – 2.10	Depositor EDS
% Data completeness (in resolution range)	82.5 (38.15-2.23) 82.5 (38.14-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.61 (at 2.10Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.188 , 0.223 0.184 , 0.220	Depositor DCC
$R_{free}$ test set	1584 reflections (5.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	37.7	Xtriage
Anisotropy	0.307	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 53.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.035 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5795	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.63 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.6943e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 8JP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/2871	0.63	0/3874
1	B	0.44	1/2752 (0.0%)	0.58	0/3705
All	All	0.45	1/5623 (0.0%)	0.61	0/7579

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	223	CYS	CB-SG	5.27	1.91	1.82

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2811	0	2741	52	0
1	B	2697	0	2655	25	0
2	A	26	0	0	0	0
2	B	26	0	0	1	0
3	A	133	0	0	10	1
3	B	102	0	0	4	0
All	All	5795	0	5396	76	1



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:300:CYS:SG	3:A:816:HOH:O	2.16	1.03
1:A:503:ASP:HB2	1:A:507:VAL:HG13	1.56	0.85
1:A:308:GLU:OE1	3:A:701:HOH:O	1.95	0.84
1:A:509:HIS:HA	1:A:510:CYS:HB2	1.59	0.83
1:A:314:THR:HG23	1:A:316:VAL:H	1.43	0.82
1:A:232:LEU:HD11	3:A:816:HOH:O	1.81	0.80
1:B:492:GLU:OE2	3:B:701:HOH:O	1.99	0.79
1:A:422:ASN:HD22	1:A:508:ARG:HD2	1.47	0.78
1:A:549:ARG:O	3:A:702:HOH:O	2.04	0.75
1:A:419:ILE:HA	1:A:460:ASN:HD22	1.51	0.74
1:B:551:GLU:HA	1:B:554:LYS:HD3	1.67	0.74
1:B:314:THR:HG22	1:B:316:VAL:H	1.52	0.73
1:B:380:ASP:OD1	3:B:702:HOH:O	2.04	0.73
1:A:311:MET:O	1:A:314:THR:HG22	1.91	0.70
1:A:416:ASP:N	1:A:416:ASP:OD1	2.27	0.68
1:A:510:CYS:N	3:A:706:HOH:O	2.27	0.67
1:A:553:GLN:N	3:A:702:HOH:O	2.28	0.67
1:B:355:LYS:HB2	1:B:421:ILE:HD13	1.78	0.66
1:A:329:VAL:HG23	1:A:396:LEU:HD11	1.79	0.64
1:A:502:ASP:HB2	1:A:507:VAL:HG21	1.78	0.64
1:A:509:HIS:CA	1:A:510:CYS:HB2	2.30	0.62
1:A:228:LEU:HD12	1:A:299:LEU:HD13	1.82	0.61
1:A:448:TYR:HB3	1:A:518:TYR:HB3	1.81	0.61
1:B:448:TYR:HB3	1:B:518:TYR:HB3	1.81	0.60
1:B:328:MET:HG3	1:B:347:TYR:CD1	2.38	0.59
1:A:503:ASP:H	1:A:507:VAL:HG22	1.68	0.58
1:B:470:ASN:HD22	1:B:470:ASN:H	1.49	0.58
1:A:279:LEU:HD22	1:A:283:PHE:CE1	2.39	0.56
1:B:242:VAL:O	1:B:245:MET:HE2	2.05	0.56
1:A:506:SER:HB3	1:A:508:ARG:HH11	1.69	0.56
1:A:216:LEU:HD23	1:A:274:VAL:HB	1.88	0.55
1:A:225:MET:HG2	1:A:229:LEU:HD22	1.88	0.55
1:B:452:ALA:HB3	1:B:517:VAL:HB	1.89	0.54
1:A:520:ARG:NH2	3:A:714:HOH:O	2.40	0.53
1:B:308:GLU:HB2	1:B:320:ILE:HG13	1.90	0.52
1:A:509:HIS:C	3:A:706:HOH:O	2.47	0.52
1:A:456:HIS:NE2	1:A:463:GLY:HA3	2.24	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:507:VAL:H	1:A:508:ARG:HA	1.75	0.52
1:A:410:MET:HB2	1:A:419:ILE:HD11	1.90	0.51
1:A:219:GLN:O	1:B:289:ASP:HB3	2.12	0.50
1:B:307:VAL:O	1:B:311:MET:HG3	2.11	0.50
1:A:279:LEU:HD22	1:A:283:PHE:HE1	1.77	0.49
1:B:305:ASP:OD1	2:B:601:8JP:N25	2.45	0.48
1:A:470:ASN:HD22	1:A:470:ASN:H	1.61	0.48
1:A:415:THR:HG22	1:A:417:GLN:HG2	1.95	0.48
1:A:416:ASP:N	1:A:417:GLN:HG3	2.28	0.48
1:B:269:HIS:NE2	1:B:533:ASP:OD2	2.47	0.48
1:B:405:GLN:NE2	1:B:515:MET:SD	2.87	0.47
1:B:470:ASN:HD22	1:B:470:ASN:N	2.11	0.47
1:A:405:GLN:NE2	1:A:515:MET:SD	2.87	0.47
1:A:503:ASP:H	1:A:507:VAL:CG2	2.27	0.46
1:A:506:SER:HB3	1:A:508:ARG:HG3	1.97	0.46
1:A:414:GLN:HA	1:A:415:THR:HA	1.56	0.46
1:A:502:ASP:H	1:A:507:VAL:HG21	1.80	0.46
1:A:384:HIS:O	1:A:387:GLN:HG2	2.16	0.45
1:A:234:PHE:CD2	1:A:471:PRO:HB3	2.52	0.45
1:A:549:ARG:NH2	3:A:705:HOH:O	2.26	0.45
1:B:387:GLN:HB3	3:B:709:HOH:O	2.17	0.45
1:A:467:VAL:HG13	1:A:480:PHE:HB2	1.98	0.45
1:A:546:GLU:HG2	3:A:745:HOH:O	2.17	0.43
1:A:378:LYS:HE3	1:A:378:LYS:HB2	1.75	0.43
1:A:378:LYS:HG3	1:A:379:TYR:H	1.83	0.43
1:A:415:THR:C	1:A:417:GLN:HG3	2.40	0.43
1:A:457:SER:HA	1:A:510:CYS:O	2.20	0.42
1:B:279:LEU:HD22	1:B:283:PHE:HE1	1.85	0.42
1:B:420:LYS:HB2	1:B:460:ASN:HA	2.01	0.42
1:B:248:GLU:CD	1:B:248:GLU:H	2.22	0.42
1:A:383:GLU:N	1:A:383:GLU:OE1	2.27	0.42
1:B:314:THR:CG2	1:B:316:VAL:H	2.28	0.42
1:A:412:ASP:HA	1:A:413:PRO:HD3	1.76	0.42
1:B:344:ARG:HD2	1:B:345:GLU:N	2.35	0.41
1:A:416:ASP:HB2	1:A:417:GLN:HA	2.02	0.41
1:A:303:LEU:HD23	1:A:303:LEU:HA	1.82	0.41
1:B:210:HIS:CE1	3:B:727:HOH:O	2.73	0.40
1:B:304:LEU:HA	1:B:307:VAL:HG22	2.03	0.40
1:A:413:PRO:HB2	1:A:414:GLN:CB	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:709:HOH:O	3:A:718:HOH:O[2_655]	1.61	0.59

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	344/369 (93%)	326 (95%)	16 (5%)	2 (1%)	28	27
1	B	326/369 (88%)	315 (97%)	11 (3%)	0	100	100
All	All	670/738 (91%)	641 (96%)	27 (4%)	2 (0%)	44	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	385	GLY
1	A	510	CYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/333 (94%)	284 (91%)	29 (9%)	10	6
1	B	299/333 (90%)	281 (94%)	18 (6%)	22	20
All	All	612/666 (92%)	565 (92%)	47 (8%)	15	11

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	228	LEU
1	A	229	LEU
1	A	252	SER
1	A	258	LEU
1	A	260	LEU
1	A	279	LEU
1	A	299	LEU
1	A	316	VAL
1	A	343	ARG
1	A	345	GLU
1	A	353	SER
1	A	360	ILE
1	A	376	ASP
1	A	378	LYS
1	A	384	HIS
1	A	393	VAL
1	A	402	LEU
1	A	410	MET
1	A	414	GLN
1	A	415	THR
1	A	416	ASP
1	A	417	GLN
1	A	470	ASN
1	A	503	ASP
1	A	507	VAL
1	A	508	ARG
1	A	510	CYS
1	A	551	GLU
1	A	553	GLN
1	B	247	THR
1	B	258	LEU
1	B	260	LEU
1	B	279	LEU
1	B	314	THR
1	B	316	VAL
1	B	337	VAL
1	B	344	ARG
1	B	351	GLN
1	B	386	LEU
1	B	402	LEU
1	B	461	HIS
1	B	470	ASN
1	B	487	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	509	HIS
1	B	524	LEU
1	B	551	GLU
1	B	553	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	460	ASN
1	B	405	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	8JP	A	601	-	28,28,28	0.88	1 (3%)	35,39,39	1.35	6 (17%)
2	8JP	B	601	-	28,28,28	0.81	1 (3%)	35,39,39	1.55	7 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8JP	A	601	-	-	0/16/16/16	0/3/3/3
2	8JP	B	601	-	-	0/16/16/16	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	8JP	C7-N16	2.00	1.39	1.34
2	B	601	8JP	C7-N16	2.14	1.39	1.34

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	8JP	C26-N25-C23	-3.10	118.26	121.84
2	B	601	8JP	C4-C5-N6	-2.57	120.55	124.49
2	B	601	8JP	C17-C22-N21	-2.32	120.40	124.24
2	B	601	8JP	O24-C23-N25	-2.31	118.85	122.49
2	A	601	8JP	C9-C8-C7	-2.14	118.39	120.69
2	A	601	8JP	C19-C20-N21	-2.11	119.78	122.67
2	A	601	8JP	C5-N6-C7	2.08	121.16	118.84
2	B	601	8JP	C22-N21-C20	2.46	121.77	117.42
2	A	601	8JP	C23-C20-N21	2.48	120.48	117.61
2	A	601	8JP	C26-N25-C23	2.62	124.88	121.84
2	A	601	8JP	C22-N21-C20	2.91	122.57	117.42
2	B	601	8JP	C5-N6-C7	3.51	122.75	118.84
2	B	601	8JP	C23-C20-N21	3.52	121.69	117.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	8JP	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	346/369 (93%)	0.49	28 (8%)	13 12	26, 43, 115, 161	0
1	B	332/369 (89%)	0.24	24 (7%)	16 16	29, 50, 97, 141	0
All	All	678/738 (91%)	0.37	52 (7%)	14 13	26, 47, 103, 161	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	503	ASP	17.9
1	A	505	LEU	14.3
1	A	415	THR	13.0
1	A	461	HIS	9.7
1	A	504	ASP	9.5
1	A	506	SER	9.4
1	A	414	GLN	8.6
1	A	502	ASP	7.8
1	B	554	LYS	7.7
1	B	555	ARG	7.6
1	A	413	PRO	7.4
1	A	554	LYS	7.3
1	B	509	HIS	7.0
1	A	507	VAL	6.8
1	A	508	ARG	6.6
1	B	461	HIS	6.3
1	A	417	GLN	6.3
1	B	208	LYS	6.1
1	A	418	ASN	5.9
1	A	411	TYR	5.5
1	B	419	ILE	5.5
1	A	509	HIS	5.3
1	B	210	HIS	4.9
1	A	460	ASN	4.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	377	ASN	4.4
1	A	376	ASP	4.4
1	B	459	ASP	4.0
1	A	416	ASP	3.9
1	B	443	LYS	3.8
1	B	374	ASP	3.4
1	B	510	CYS	3.1
1	A	385	GLY	3.0
1	B	376	ASP	3.0
1	A	209	LYS	2.9
1	B	377	ASN	2.9
1	B	410	MET	2.8
1	A	550	ILE	2.8
1	A	412	ASP	2.8
1	B	550	ILE	2.7
1	B	336	GLU	2.6
1	A	210	HIS	2.5
1	A	388	GLU	2.5
1	B	553	GLN	2.5
1	B	462	GLY	2.5
1	B	209	LYS	2.5
1	B	539	GLN	2.4
1	B	551	GLU	2.4
1	B	500	GLY	2.4
1	A	462	GLY	2.2
1	B	331	TYR	2.2
1	A	553	GLN	2.1
1	B	335	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	8JP	A	601	26/26	0.95	0.14	-0.28	24,32,36,37	0
2	8JP	B	601	26/26	0.93	0.12	-0.58	38,48,53,54	0

## 6.5 Other polymers

There are no such residues in this entry.