



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 20, 2018 – 08:18 PM EST

PDB ID : 5UW4
Title : Activated yeast Glycogen Synthase in complex with UDP glucosamine
Authors : Mahalingan, K.K.; Hurley, T.D.
Deposited on : 2017-02-20
Resolution : 2.98 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2168 (3.00-2.96)
Clashscore	112137	2535 (3.00-2.96)
Ramachandran outliers	110173	2451 (3.00-2.96)
Sidechain outliers	110143	2454 (3.00-2.96)
RSRZ outliers	101464	2192 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	720	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>11%</div> <div>•</div> <div>11%</div> </div> </div>
1	B	720	<div> <div>3%</div> <div> <div></div> <div>79%</div> <div>9%</div> <div>•</div> <div>12%</div> </div> </div>
1	C	720	<div> <div>6%</div> <div> <div></div> <div>77%</div> <div>11%</div> <div>•</div> <div>11%</div> </div> </div>
1	D	720	<div> <div>6%</div> <div> <div></div> <div>75%</div> <div>13%</div> <div>•</div> <div>12%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycogen [starch] synthase isoform 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	638	Total	C	N	O	S	0	0	0
			5127	3272	894	942	19			
1	B	636	Total	C	N	O	S	0	0	0
			5084	3246	886	933	19			
1	A	638	Total	C	N	O	S	0	0	0
			4980	3185	853	924	18			
1	D	636	Total	C	N	O	S	0	0	0
			5005	3192	868	927	18			

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-19	MET	-	initiating methionine	UNP P27472
C	-18	GLY	-	expression tag	UNP P27472
C	-17	SER	-	expression tag	UNP P27472
C	-16	SER	-	expression tag	UNP P27472
C	-15	HIS	-	expression tag	UNP P27472
C	-14	HIS	-	expression tag	UNP P27472
C	-13	HIS	-	expression tag	UNP P27472
C	-12	HIS	-	expression tag	UNP P27472
C	-11	HIS	-	expression tag	UNP P27472
C	-10	HIS	-	expression tag	UNP P27472
C	-9	SER	-	expression tag	UNP P27472
C	-8	SER	-	expression tag	UNP P27472
C	-7	GLY	-	expression tag	UNP P27472
C	-6	LEU	-	expression tag	UNP P27472
C	-5	VAL	-	expression tag	UNP P27472
C	-4	PRO	-	expression tag	UNP P27472
C	-3	ARG	-	expression tag	UNP P27472
C	-2	GLY	-	expression tag	UNP P27472
C	-1	SER	-	expression tag	UNP P27472
C	0	HIS	-	expression tag	UNP P27472
C	169	GLN	GLU	engineered mutation	UNP P27472

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	535	SER	ALA	conflict	UNP P27472
B	-19	MET	-	initiating methionine	UNP P27472
B	-18	GLY	-	expression tag	UNP P27472
B	-17	SER	-	expression tag	UNP P27472
B	-16	SER	-	expression tag	UNP P27472
B	-15	HIS	-	expression tag	UNP P27472
B	-14	HIS	-	expression tag	UNP P27472
B	-13	HIS	-	expression tag	UNP P27472
B	-12	HIS	-	expression tag	UNP P27472
B	-11	HIS	-	expression tag	UNP P27472
B	-10	HIS	-	expression tag	UNP P27472
B	-9	SER	-	expression tag	UNP P27472
B	-8	SER	-	expression tag	UNP P27472
B	-7	GLY	-	expression tag	UNP P27472
B	-6	LEU	-	expression tag	UNP P27472
B	-5	VAL	-	expression tag	UNP P27472
B	-4	PRO	-	expression tag	UNP P27472
B	-3	ARG	-	expression tag	UNP P27472
B	-2	GLY	-	expression tag	UNP P27472
B	-1	SER	-	expression tag	UNP P27472
B	0	HIS	-	expression tag	UNP P27472
B	169	GLN	GLU	engineered mutation	UNP P27472
B	535	SER	ALA	conflict	UNP P27472
A	-19	MET	-	initiating methionine	UNP P27472
A	-18	GLY	-	expression tag	UNP P27472
A	-17	SER	-	expression tag	UNP P27472
A	-16	SER	-	expression tag	UNP P27472
A	-15	HIS	-	expression tag	UNP P27472
A	-14	HIS	-	expression tag	UNP P27472
A	-13	HIS	-	expression tag	UNP P27472
A	-12	HIS	-	expression tag	UNP P27472
A	-11	HIS	-	expression tag	UNP P27472
A	-10	HIS	-	expression tag	UNP P27472
A	-9	SER	-	expression tag	UNP P27472
A	-8	SER	-	expression tag	UNP P27472
A	-7	GLY	-	expression tag	UNP P27472
A	-6	LEU	-	expression tag	UNP P27472
A	-5	VAL	-	expression tag	UNP P27472
A	-4	PRO	-	expression tag	UNP P27472
A	-3	ARG	-	expression tag	UNP P27472
A	-2	GLY	-	expression tag	UNP P27472
A	-1	SER	-	expression tag	UNP P27472

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	expression tag	UNP P27472
A	169	GLN	GLU	engineered mutation	UNP P27472
A	535	SER	ALA	conflict	UNP P27472
D	-19	MET	-	initiating methionine	UNP P27472
D	-18	GLY	-	expression tag	UNP P27472
D	-17	SER	-	expression tag	UNP P27472
D	-16	SER	-	expression tag	UNP P27472
D	-15	HIS	-	expression tag	UNP P27472
D	-14	HIS	-	expression tag	UNP P27472
D	-13	HIS	-	expression tag	UNP P27472
D	-12	HIS	-	expression tag	UNP P27472
D	-11	HIS	-	expression tag	UNP P27472
D	-10	HIS	-	expression tag	UNP P27472
D	-9	SER	-	expression tag	UNP P27472
D	-8	SER	-	expression tag	UNP P27472
D	-7	GLY	-	expression tag	UNP P27472
D	-6	LEU	-	expression tag	UNP P27472
D	-5	VAL	-	expression tag	UNP P27472
D	-4	PRO	-	expression tag	UNP P27472
D	-3	ARG	-	expression tag	UNP P27472
D	-2	GLY	-	expression tag	UNP P27472
D	-1	SER	-	expression tag	UNP P27472
D	0	HIS	-	expression tag	UNP P27472
D	169	GLN	GLU	engineered mutation	UNP P27472
D	535	SER	ALA	conflict	UNP P27472

- Molecule 2 is D-GLUCOSAMINE (three-letter code: GCS) (formula: $C_6H_{13}NO_5$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	0	0
			12	6	1	5		

- Molecule 3 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	D	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

- Molecule 4 is ALPHA-D-GLUCOSE-6-PHOSPHATE (three-letter code: G6P) (formula: $C_6H_{13}O_9P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	O	P	0	0
			16	6	9	1		
4	B	1	Total	C	O	P	0	0
			16	6	9	1		
4	A	1	Total	C	O	P	0	0
			16	6	9	1		
4	D	1	Total	C	O	P	0	0
			16	6	9	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	O	0	0
			1	1		
5	B	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Glycogen [starch] synthase isoform 2





- Molecule 1: Glycogen [starch] synthase isoform 2



- Molecule 1: Glycogen [starch] synthase isoform 2



• Molecule 1: Glycogen [starch] synthase isoform 2



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	193.04Å 204.70Å 205.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	145.16 – 2.98 29.95 – 2.98	Depositor EDS
% Data completeness (in resolution range)	99.6 (145.16-2.98) 99.7 (29.95-2.98)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 3.00Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.188 , 0.257 0.197 , 0.263	Depositor DCC
R_{free} test set	4474 reflections (5.73%)	DCC
Wilson B-factor (Å ²)	74.6	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 59.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.003 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	20374	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, GCS, G6P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/5101	0.76	3/6943 (0.0%)
1	B	0.59	0/5206	0.83	1/7061 (0.0%)
1	C	0.57	0/5251	0.80	6/7119 (0.1%)
1	D	0.56	0/5123	0.78	3/6962 (0.0%)
All	All	0.57	0/20681	0.80	13/28085 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	199	ARG	NE-CZ-NH2	-6.70	116.95	120.30
1	B	34	ILE	N-CA-CB	6.51	125.78	110.80
1	C	320	ARG	NE-CZ-NH1	6.48	123.54	120.30
1	A	131	PRO	N-CA-C	-6.29	95.73	112.10
1	C	199	ARG	NE-CZ-NH1	6.12	123.36	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	18	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4980	0	4753	61	0
1	B	5084	0	4955	49	0
1	C	5127	0	5015	64	0
1	D	5005	0	4827	70	0
2	C	12	0	13	1	0
3	A	25	0	11	1	0
3	B	25	0	11	0	0
3	C	25	0	11	1	0
3	D	25	0	11	2	0
4	A	16	0	11	1	0
4	B	16	0	11	1	0
4	C	16	0	11	1	0
4	D	16	0	11	2	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
All	All	20374	0	19651	239	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 239 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:217:ASP:CB	1:C:220:HIS:HD2	1.36	1.39
1:C:217:ASP:CB	1:C:220:HIS:CD2	2.28	1.16
1:C:217:ASP:HB3	1:C:220:HIS:CD2	1.80	1.16
1:C:217:ASP:CG	1:C:220:HIS:CD2	2.20	1.14
1:C:217:ASP:HB3	1:C:220:HIS:HD2	0.97	1.10

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	636/720 (88%)	575 (90%)	58 (9%)	3 (0%)	32	72
1	B	632/720 (88%)	585 (93%)	46 (7%)	1 (0%)	51	85
1	C	636/720 (88%)	586 (92%)	50 (8%)	0	100	100
1	D	632/720 (88%)	579 (92%)	52 (8%)	1 (0%)	51	85
All	All	2536/2880 (88%)	2325 (92%)	206 (8%)	5 (0%)	51	85

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	401	PRO
1	A	544	ASN
1	A	183	ARG
1	A	363	SER
1	D	320	ARG

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	516/621 (83%)	504 (98%)	12 (2%)	56	84
1	B	539/621 (87%)	537 (100%)	2 (0%)	93	98
1	C	547/621 (88%)	543 (99%)	4 (1%)	87	95
1	D	523/621 (84%)	517 (99%)	6 (1%)	78	93
All	All	2125/2484 (86%)	2101 (99%)	24 (1%)	78	93

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	95	TRP
1	A	326	LYS
1	D	184	ILE
1	A	124	SER
1	A	125	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	403	ASN
1	D	372	GLN
1	B	634	ASN
1	B	19	ASN
1	D	283	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	UDP	A	801	-	21,26,26	0.83	0	22,40,40	1.83	2 (9%)
4	G6P	A	802	-	16,16,16	0.68	0	24,24,24	1.24	0
3	UDP	B	801	-	21,26,26	1.06	1 (4%)	22,40,40	1.79	1 (4%)
4	G6P	B	802	-	16,16,16	0.59	0	24,24,24	1.31	2 (8%)
2	GCS	C	801	-	12,12,12	0.73	0	15,17,17	1.63	3 (20%)
3	UDP	C	802	-	21,26,26	0.93	1 (4%)	22,40,40	2.21	4 (18%)
4	G6P	C	803	-	16,16,16	0.75	0	24,24,24	1.34	3 (12%)
3	UDP	D	801	-	21,26,26	0.93	1 (4%)	22,40,40	1.88	1 (4%)
4	G6P	D	802	-	16,16,16	0.66	0	24,24,24	1.45	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UDP	A	801	-	-	0/12/32/32	0/2/2/2
4	G6P	A	802	-	-	0/6/26/26	0/1/1/1
3	UDP	B	801	-	-	0/12/32/32	0/2/2/2
4	G6P	B	802	-	-	0/6/26/26	0/1/1/1
2	GCS	C	801	-	-	0/2/22/22	0/1/1/1
3	UDP	C	802	-	-	0/12/32/32	0/2/2/2
4	G6P	C	803	-	-	0/6/26/26	0/1/1/1
3	UDP	D	801	-	-	0/12/32/32	0/2/2/2
4	G6P	D	802	-	-	0/6/26/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	801	UDP	PB-O3A	2.45	1.64	1.60
3	C	802	UDP	O4'-C1'	2.62	1.44	1.41
3	B	801	UDP	PB-O3A	3.49	1.65	1.60

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	802	G6P	O3-C3-C2	-3.77	102.15	110.36
4	D	802	G6P	O3-C3-C4	-2.73	104.41	110.36
4	B	802	G6P	O3-C3-C4	-2.47	104.98	110.36
2	C	801	GCS	C3-C2-N2	-2.41	106.10	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	803	G6P	O3-C3-C2	-2.40	105.12	110.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	801	UDP	1	0
4	A	802	G6P	1	0
4	B	802	G6P	1	0
2	C	801	GCS	1	0
3	C	802	UDP	1	0
4	C	803	G6P	1	0
3	D	801	UDP	2	0
4	D	802	G6P	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	638/720 (88%)	0.19	29 (4%) 34 19	50, 86, 141, 178	0
1	B	636/720 (88%)	-0.02	19 (2%) 51 30	39, 71, 115, 166	0
1	C	638/720 (88%)	0.18	42 (6%) 19 10	45, 80, 129, 169	0
1	D	636/720 (88%)	0.25	44 (6%) 18 9	42, 87, 146, 164	0
All	All	2548/2880 (88%)	0.15	134 (5%) 27 15	39, 80, 138, 178	0

The worst 5 of 134 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	124	SER	5.7
1	D	630	LEU	5.4
1	C	125	LEU	5.1
1	D	2	SER	5.0
1	A	52	ALA	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	UDP	B	801	25/25	0.88	0.20	0.32	81,116,172,177	0
2	GCS	C	801	12/12	0.97	0.18	-0.10	60,67,78,78	0
3	UDP	C	802	25/25	0.96	0.16	-0.60	61,77,83,86	0
4	G6P	C	803	16/16	0.98	0.16	-0.62	56,66,75,76	0
3	UDP	D	801	25/25	0.92	0.14	-0.83	80,99,127,130	0
4	G6P	B	802	16/16	0.96	0.14	-0.85	39,49,55,58	0
4	G6P	D	802	16/16	0.98	0.13	-0.97	43,51,58,63	0
3	UDP	A	801	25/25	0.91	0.15	-1.11	79,94,109,114	0
4	G6P	A	802	16/16	0.97	0.12	-1.27	50,58,65,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.