



Full wwPDB X-ray Structure Validation Report i

Feb 13, 2017 – 07:02 pm GMT

PDB ID : 4V04
Title : FGFR1 in complex with ponatinib.
Authors : Tucker, J.; Klein, T.; Breed, J.; Breeze, A.; Overman, R.; Phillips, C.; Norman, R.A.
Deposited on : 2014-09-10
Resolution : 2.12 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4988 (2.14-2.10)
Clashscore	112137	5557 (2.14-2.10)
Ramachandran outliers	110173	5504 (2.14-2.10)
Sidechain outliers	110143	5505 (2.14-2.10)
RSRZ outliers	101464	5021 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	A	1767	-	-	-	X
2	EDO	A	1768	-	-	-	X
2	EDO	B	1769	-	-	-	X
2	EDO	B	1770	-	-	-	X
3	OLI	B	1771[A]	-	-	-	X
4	SO4	A	1776	-	-	-	X

2 Entry composition [\(i\)](#)

There are 5 unique types of molecules in this entry. The entry contains 5161 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FIBROBLAST GROWTH FACTOR RECEPTOR 1 (FMS-RELATED TYROSINE KINASE 2, PFEIFFER SYNDROME), ISOFORM CRA_B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	8	1
			2318	1476	395	428	19			
1	B	289	Total	C	N	O	S	0	6	0
			2276	1451	386	421	18			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	457	GLY	-	EXPRESSION TAG	UNP D3DSX2
A	488	ALA	CYS	ENGINEERED MUTATION	UNP D3DSX2
A	584	SER	CYS	ENGINEERED MUTATION	UNP D3DSX2
B	457	GLY	-	EXPRESSION TAG	UNP D3DSX2
B	488	ALA	CYS	ENGINEERED MUTATION	UNP D3DSX2
B	584	SER	CYS	ENGINEERED MUTATION	UNP D3DSX2

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is 3-(IMIDAZO[1,2-B]PYRIDAZIN-3-YLETHYNYL)-4-METHYL-N-{4-[(4-METHYLPYRROLIDIN-1-YL)METHYL]-3-(TRIFLUOROMETHYL)PHENYL}BENZAMIDE (three-letter code: 0LI) (formula: C₂₉H₂₇F₃N₆O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C 39	F 29	N 3	O 6	1	0
3	B	1	Total	C 39	F 29	N 3	O 6	1	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O 5	S 4	1	0
4	A	1	Total	O 5	S 4	1	0
4	A	1	Total	O 5	S 4	1	0
4	A	1	Total	O 5	S 4	1	0
4	B	1	Total	O 5	S 4	1	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	215	Total	O 215	O 215	0	0
5	B	205	Total	O 205	O 205	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FIBROBLAST GROWTH FACTOR RECEPTOR 1 (FMS-RELATED TYROSINE KINASE 2, PFEIFFER SYNDROME), ISOFORM CRA_B



- Molecule 1: FIBROBLAST GROWTH FACTOR RECEPTOR 1 (FMS-RELATED TYROSINE KINASE 2, PFEIFFER SYNDROME), ISOFORM CRA_B



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	207.25Å 58.09Å 65.23Å 90.00° 107.38° 90.00°	Depositor
Resolution (Å)	21.37 – 2.12 21.38 – 2.11	Depositor EDS
% Data completeness (in resolution range)	97.6 (21.37-2.12) 95.5 (21.38-2.11)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	3.32 (at 2.11Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R , R_{free}	0.186 , 0.228 0.244 , 0.270	Depositor DCC
R_{free} test set	2068 reflections (5.06%)	DCC
Wilson B-factor (Å ²)	37.2	Xtriage
Anisotropy	0.618	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 64.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.017 for -h-2*l,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5161	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 0LI, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/2364	0.64	0/3197
1	B	0.49	0/2321	0.62	0/3139
All	All	0.50	0/4685	0.63	0/6336

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2318	0	2274	11	0
1	B	2276	0	2221	8	1
2	A	24	0	36	0	0
2	B	20	0	30	0	0
3	A	39	0	27	7	0
3	B	39	0	27	6	0
4	A	20	0	0	1	0
4	B	5	0	0	0	0
5	A	215	0	0	1	3
5	B	205	0	0	0	0
All	All	5161	0	4615	20	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:514:LYS:HB2	3:A:1772[A]:0LI:H11	1.62	0.80
4:A:1773:SO4:O2	5:A:2209:HOH:O	2.00	0.79
1:A:622:ARG:HH11	3:A:1772[A]:0LI:H25	1.52	0.74
1:B:514:LYS:HB2	3:B:1771[A]:0LI:H11	1.76	0.68
1:B:622:ARG:HH11	3:B:1771[A]:0LI:H25	1.62	0.64
1:A:619[A]:CYS:SG	3:A:1772[A]:0LI:H24A	2.46	0.56
1:A:562:GLU:O	3:A:1772[A]:0LI:H1	2.11	0.50
1:A:607:VAL:HG13	1:A:639[A]:ILE:HD11	1.96	0.48
1:A:531:GLU:HG3	3:A:1772[A]:0LI:H14	1.97	0.46
1:B:514:LYS:HB2	3:B:1771[A]:0LI:C11	2.45	0.46
1:B:562:GLU:O	3:B:1771[A]:0LI:H1	2.16	0.46
1:A:610:GLY:HA3	1:A:639[B]:ILE:HD12	1.97	0.46
1:A:622:ARG:NH1	3:A:1772[A]:0LI:H25	2.27	0.45
1:A:504:LYS:HG2	1:A:506:ASN:H	1.82	0.45
1:A:477:ARG:NE	1:A:477:ARG:HA	2.33	0.44
1:A:619[A]:CYS:SG	3:A:1772[A]:0LI:C24	3.07	0.43
1:B:492:VAL:HG13	3:B:1771[A]:0LI:H11A	2.02	0.42
1:B:615:ALA:HA	1:B:679:HIS:CE1	2.55	0.42
1:B:461:SER:HA	1:B:464:GLU:O	2.20	0.41
1:B:531:GLU:HG3	3:B:1771[A]:0LI:H14	2.03	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2036:HOH:O	5:A:2036:HOH:O[2_656]	1.98	0.22
1:B:554:ASP:O	5:A:2190:HOH:O[4_556]	2.18	0.02
5:A:2034:HOH:O	5:A:2034:HOH:O[2_656]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	291/309 (94%)	286 (98%)	4 (1%)	1 (0%)	44 43
1	B	287/309 (93%)	283 (99%)	3 (1%)	1 (0%)	44 43
All	All	578/618 (94%)	569 (98%)	7 (1%)	2 (0%)	38 43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	460	VAL
1	A	587	PRO

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	247/269 (92%)	241 (98%)	6 (2%)	54 58
1	B	237/269 (88%)	231 (98%)	6 (2%)	53 56
All	All	484/538 (90%)	472 (98%)	12 (2%)	56 56

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	534	MET
1	A	609	ARG
1	A	642[A]	PHE
1	A	642[B]	PHE
1	A	720	ASP
1	A	731	MET
1	B	534	MET
1	B	594	GLN
1	B	644[B]	LEU
1	B	711	LYS
1	B	720	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	724	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	1766	-	3,3,3	0.42	0	2,2,2	0.29	0
2	EDO	A	1767	-	3,3,3	0.46	0	2,2,2	0.91	0
2	EDO	A	1768	-	3,3,3	0.81	0	2,2,2	0.08	0
2	EDO	A	1769	-	3,3,3	0.68	0	2,2,2	0.14	0
2	EDO	A	1770	-	3,3,3	0.84	0	2,2,2	0.06	0
2	EDO	A	1771	-	3,3,3	0.63	0	2,2,2	0.36	0
3	OLI	A	1772[A]	-	40,43,43	0.79	0	51,62,62	0.75	1 (1%)
4	SO4	A	1773	-	4,4,4	0.52	0	6,6,6	0.27	0
4	SO4	A	1774	-	4,4,4	0.20	0	6,6,6	0.46	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	1775	-	4,4,4	0.19	0	6,6,6	0.19	0
4	SO4	A	1776	-	4,4,4	0.35	0	6,6,6	0.16	0
2	EDO	B	1766	-	3,3,3	0.31	0	2,2,2	0.61	0
2	EDO	B	1767	-	3,3,3	0.54	0	2,2,2	0.42	0
2	EDO	B	1768	-	3,3,3	0.53	0	2,2,2	0.49	0
2	EDO	B	1769	-	3,3,3	0.58	0	2,2,2	0.41	0
2	EDO	B	1770	-	3,3,3	0.68	0	2,2,2	0.32	0
3	OLI	B	1771[A]	-	40,43,43	0.89	1 (2%)	51,62,62	0.95	3 (5%)
4	SO4	B	1772	-	4,4,4	0.29	0	6,6,6	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1766	-	-	0/1/1/1	0/0/0/0
2	EDO	A	1767	-	-	0/1/1/1	0/0/0/0
2	EDO	A	1768	-	-	0/1/1/1	0/0/0/0
2	EDO	A	1769	-	-	0/1/1/1	0/0/0/0
2	EDO	A	1770	-	-	0/1/1/1	0/0/0/0
2	EDO	A	1771	-	-	0/1/1/1	0/0/0/0
3	OLI	A	1772[A]	-	-	0/20/33/33	0/5/5/5
4	SO4	A	1773	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1774	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1775	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1776	-	-	0/0/0/0	0/0/0/0
2	EDO	B	1766	-	-	0/1/1/1	0/0/0/0
2	EDO	B	1767	-	-	0/1/1/1	0/0/0/0
2	EDO	B	1768	-	-	0/1/1/1	0/0/0/0
2	EDO	B	1769	-	-	0/1/1/1	0/0/0/0
2	EDO	B	1770	-	-	0/1/1/1	0/0/0/0
3	OLI	B	1771[A]	-	-	0/20/33/33	0/5/5/5
4	SO4	B	1772	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1771[A]	OLI	C82-C83	2.12	1.44	1.37

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1771[A]	OLI	C82-C81-C84	-2.17	117.97	120.22
3	A	1772[A]	OLI	C82-C81-C84	-2.12	118.02	120.22
3	B	1771[A]	OLI	C8-C9-C12	2.03	127.14	120.61
3	B	1771[A]	OLI	C19-C17-C16	2.85	124.11	121.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1772[A]	OLI	7	0
4	A	1773	SO4	1	0
3	B	1771[A]	OLI	6	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/309 (94%)	1.20	50 (17%) 2 2	26, 45, 87, 110	2 (0%)
1	B	289/309 (93%)	0.83	38 (13%) 4 5	33, 48, 82, 105	2 (0%)
All	All	582/618 (94%)	1.02	88 (15%) 3 3	26, 47, 86, 110	4 (0%)

All (88) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	644[B]	LEU	7.0
1	A	587	PRO	6.2
1	B	458	ALA	5.8
1	B	503	ASP	5.6
1	B	764	GLN	5.4
1	A	485	GLY	5.3
1	A	588	SER	5.3
1	B	579	PRO	5.3
1	A	520	ALA	5.2
1	A	486	GLU	5.1
1	B	651	ILE	5.0
1	A	658	THR	5.0
1	A	583	TYR	5.0
1	B	505	PRO	4.9
1	A	521	THR	4.8
1	A	503	ASP	4.7
1	B	489	PHE	4.6
1	B	658	THR	4.6
1	B	652	ASP	4.6
1	B	644[B]	LEU	4.6
1	A	581	LEU	4.5
1	A	654	TYR	4.4
1	A	463	TYR	4.2
1	B	459	GLY	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	590	ASN	3.9
1	B	502	LYS	3.9
1	A	490	GLY	3.7
1	A	560	ILE	3.7
1	B	654	TYR	3.7
1	A	653	TYR	3.5
1	B	659	ASN	3.5
1	A	651	ILE	3.4
1	A	660	GLY	3.3
1	A	589	HIS	3.3
1	B	559	VAL	3.3
1	A	582	GLU	3.3
1	B	487	GLY	3.2
1	A	659	ASN	3.2
1	A	516	LEU	3.2
1	A	592	GLU	3.1
1	A	519	ASP	3.1
1	B	765	GLU	3.0
1	B	508	VAL	3.0
1	A	591	PRO	3.0
1	A	471	TRP	2.9
1	A	652	ASP	2.9
1	A	502	LYS	2.9
1	A	642[A]	PHE	2.9
1	B	607	VAL	2.8
1	B	653	TYR	2.8
1	A	501	ASP	2.7
1	A	725	CYS	2.5
1	B	547	LEU	2.5
1	A	559	VAL	2.5
1	A	657	THR	2.5
1	B	657	THR	2.5
1	B	558	TYR	2.5
1	A	760	LEU	2.5
1	B	560	ILE	2.5
1	B	506	ASN	2.5
1	B	594	GLN	2.3
1	B	498	ILE	2.3
1	A	518	SER	2.3
1	B	604	ALA	2.3
1	A	580	GLY	2.3
1	B	488	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	639[A]	ILE	2.3
1	B	656	LYS	2.2
1	B	693	ILE	2.2
1	A	522	GLU	2.2
1	A	686	PHE	2.2
1	A	747	PHE	2.2
1	B	688	VAL	2.2
1	A	504	LYS	2.2
1	A	517	LYS	2.2
1	B	463	TYR	2.2
1	A	693	ILE	2.1
1	A	464	GLU	2.1
1	A	527	ASP	2.1
1	A	629	VAL	2.1
1	A	630	LEU	2.0
1	B	639	ILE	2.0
1	A	640	ALA	2.0
1	B	603	CYS	2.0
1	B	760	LEU	2.0
1	B	461	SER	2.0
1	B	537[A]	MET	2.0
1	A	465	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	EDO	A	1767	4/4	0.83	0.29	5.47	39,41,41,43	0
4	SO4	A	1776	5/5	0.79	0.24	5.30	69,70,73,75	5
2	EDO	A	1768	4/4	0.74	0.33	4.61	59,61,62,65	0
2	EDO	B	1769	4/4	0.66	0.27	3.29	62,65,69,70	0
2	EDO	B	1770	4/4	0.80	0.20	2.87	65,68,69,69	0
3	OLI	B	1771[A]	39/39	0.79	0.28	2.19	30,38,45,49	39
2	EDO	A	1770	4/4	0.55	0.24	1.87	57,57,59,60	0
2	EDO	B	1766	4/4	0.86	0.22	1.60	50,51,51,53	0
3	OLI	A	1772[A]	39/39	0.78	0.28	1.21	27,37,42,43	39
2	EDO	B	1767	4/4	0.86	0.17	0.66	61,61,62,62	0
4	SO4	B	1772	5/5	0.93	0.22	0.25	48,58,61,64	0
4	SO4	A	1774	5/5	0.93	0.18	-0.38	65,72,76,78	0
2	EDO	A	1766	4/4	0.91	0.15	-1.65	30,31,33,34	0
2	EDO	A	1771	4/4	0.78	0.24	-	58,61,62,64	0
2	EDO	B	1768	4/4	0.73	0.31	-	66,69,70,70	0
4	SO4	A	1775	5/5	0.83	0.29	-	89,89,92,93	0
2	EDO	A	1769	4/4	0.59	0.25	-	70,70,71,71	0
4	SO4	A	1773	5/5	0.88	0.22	-	33,41,42,44	5

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.