



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 07:16 am GMT

PDB ID : 2V17  
Title : STRUCTURE OF THE COMPLEX OF ANTIBODY MN423 WITH A  
FRAGMENT OF TAU PROTEIN  
Authors : Sevcik, J.; Skrabana, R.; Csokova, N.; Dvorsky, R.; Novak, M.  
Deposited on : 2007-05-22  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1368 (1.66-1.66)
Clashscore	112137	1468 (1.66-1.66)
Ramachandran outliers	110173	1438 (1.66-1.66)
Sidechain outliers	110143	1438 (1.66-1.66)
RSRZ outliers	101464	1371 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	6	<div> <div>17%</div> <div> <div></div> <div>83%</div> <div>17%</div> </div> </div>
2	H	222	<div> <div>9%</div> <div> <div></div> <div>78%</div> <div>18%</div> <div>• •</div> </div> </div>
3	L	214	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>19%</div> <div>•</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4069 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEPTIDE FRAGMENT.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	6	Total	C	N	O	0	0	0
			44	24	8	12			

- Molecule 2 is a protein called MONOCLONAL ANTIBODY FAB FRAGMENT MN423.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	222	Total	C	N	O	S	0	1	0
			1651	1038	273	331	9			

- Molecule 3 is a protein called MONOCLONAL ANTIBODY FAB FRAGMENT MN423.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	214	Total	C	N	O	S	0	1	0
			1672	1042	283	339	8			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total	O	0	0
			7	7		
4	H	343	Total	O	0	0
			343	343		
4	L	352	Total	O	0	0
			352	352		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



#### • Molecule 1: PEPTIDE FRAGMENT



#### • Molecule 2: MONOCLONAL ANTIBODY FAB FRAGMENT MN423



#### • Molecule 3: MONOCLONAL ANTIBODY FAB FRAGMENT MN423



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.54Å 36.81Å 85.54Å 90.00° 113.93° 90.00°	Depositor
Resolution (Å)	79.06 – 1.65 19.88 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.7 (79.06-1.65) 99.8 (19.88-1.65)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.93 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.156 , 0.218 0.170 , 0.228	Depositor DCC
$R_{free}$ test set	2504 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.0	Xtriage
Anisotropy	0.158	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 57.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.027 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4069	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.71	0/44	1.12	0/57
2	H	0.76	2/1696 (0.1%)	0.89	4/2316 (0.2%)
3	L	0.70	0/1715	0.93	4/2324 (0.2%)
All	All	0.73	2/3455 (0.1%)	0.91	8/4697 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	1	0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	74	ARG	CZ-NH1	-5.53	1.25	1.33
2	H	185	THR	CB-CG2	-5.07	1.35	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	70	ASP	CB-CG-OD1	9.14	126.52	118.30
3	L	93	ASP	CB-CG-OD2	6.71	124.34	118.30
3	L	143	ASP	CB-CG-OD1	6.28	123.95	118.30
2	H	185	THR	CA-CB-CG2	6.16	121.02	112.40
2	H	106	ARG	NE-CZ-NH2	-6.00	117.30	120.30
2	H	75	ASP	CB-CG-OD1	5.68	123.41	118.30
2	H	139	ASP	CB-CG-OD2	5.63	123.37	118.30
3	L	108	ARG	NE-CZ-NH1	5.08	122.84	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	H	185	THR	CB

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	44	0	31	0	0
2	H	1651	0	1604	60	0
3	L	1672	0	1610	49	0
4	A	7	0	0	0	0
4	H	343	0	0	29	0
4	L	352	0	0	21	1
All	All	4069	0	3245	103	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:173:HIS:CG	4:H:2272:HOH:O	1.83	1.27
3:L:180:THR:HB	4:L:2286:HOH:O	1.25	1.27
3:L:194:CYS:SG	4:L:2223:HOH:O	1.98	1.21
3:L:134:CYS:SG	4:L:2223:HOH:O	1.97	1.18
3:L:81:GLU:HB2	4:L:2166:HOH:O	1.47	1.13
2:H:60:THR:HB	4:H:2115:HOH:O	0.80	0.98
2:H:173:HIS:ND1	4:H:2272:HOH:O	1.89	0.94
3:L:134:CYS:CB	4:L:2223:HOH:O	2.13	0.87
2:H:222:SER:HA	4:H:2332:HOH:O	1.75	0.84
2:H:180:GLN:HE22	3:L:180:THR:HG21	1.42	0.83
3:L:39:LYS:NZ	3:L:81:GLU:O	2.14	0.79
2:H:193:PRO:O	2:H:196:THR:HG22	1.83	0.78
3:L:30:ARG:HE	3:L:92:ASN:HD21	1.32	0.77
2:H:185:THR:HG21	4:H:2236:HOH:O	1.86	0.75
3:L:134:CYS:HB2	4:L:2223:HOH:O	1.82	0.75

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:114:GLN:HG2	4:H:2197:HOH:O	1.90	0.70
3:L:180:THR:CG2	4:L:2222:HOH:O	2.39	0.70
2:H:173:HIS:HD1	3:L:174:SER:CB	2.03	0.69
2:H:30:THR:CG2	2:H:54:LYS:HB2	2.22	0.69
3:L:180:THR:HG22	4:L:2222:HOH:O	1.93	0.69
3:L:90:GLN:HE22	3:L:93:ASP:H	1.42	0.68
2:H:173:HIS:CE1	3:L:174:SER:OG	2.46	0.68
3:L:61:ARG:HD3	4:L:2124:HOH:O	1.94	0.67
3:L:210:ASN:HD22	3:L:212:ASN:H	1.40	0.67
2:H:173:HIS:CE1	4:H:2272:HOH:O	2.33	0.67
3:L:37:ARG:NE	3:L:39:LYS:HZ2	1.94	0.65
2:H:173:HIS:ND1	3:L:174:SER:OG	2.31	0.64
2:H:214:THR:HG23	4:H:2315:HOH:O	1.97	0.64
2:H:196:THR:HB	4:H:2295:HOH:O	1.97	0.64
2:H:43:LYS:N	4:H:2092:HOH:O	2.31	0.63
2:H:164:ASN:HB2	4:H:2256:HOH:O	1.97	0.63
3:L:149:LYS:HB2	3:L:193:THR:CG2	2.29	0.62
2:H:30:THR:HG23	2:H:58:TYR:OH	2.00	0.61
3:L:81:GLU:HG3	4:L:2168:HOH:O	1.99	0.61
2:H:16:GLY:HA3	4:H:2034:HOH:O	2.00	0.61
3:L:45:LYS:NZ	4:L:2097:HOH:O	2.32	0.61
2:H:60:THR:HG21	2:H:62:TYR:CE2	2.36	0.59
2:H:173:HIS:CD2	4:H:2272:HOH:O	2.29	0.59
2:H:16:GLY:HA2	4:H:2046:HOH:O	2.01	0.59
2:H:119:THR:HG22	4:H:2203:HOH:O	2.02	0.59
2:H:171:SER:HB2	4:L:2226:HOH:O	2.06	0.56
2:H:60:THR:HG21	2:H:62:TYR:CZ	2.41	0.56
2:H:180:GLN:NE2	4:H:2236:HOH:O	2.39	0.55
3:L:115:VAL:HG11	4:L:2328:HOH:O	2.07	0.55
3:L:198:HIS:HD2	3:L:200:THR:OG1	1.89	0.54
3:L:93:ASP:HB3	4:L:2176:HOH:O	2.07	0.54
2:H:164:ASN:HA	2:H:165:SER:OG	2.07	0.53
2:H:136:GLY:HA3	2:H:222:SER:HB2	1.90	0.53
2:H:119:THR:HG21	4:H:2239:HOH:O	2.08	0.53
3:L:39:LYS:HE2	3:L:42:LYS:NZ	2.24	0.53
3:L:4:ILE:HD11	3:L:97:THR:O	2.09	0.53
2:H:119:THR:HG23	4:H:2204:HOH:O	2.08	0.53
2:H:222:SER:C	4:H:2333:HOH:O	2.48	0.52
3:L:149:LYS:HB2	3:L:193:THR:HG22	1.92	0.52
3:L:115:VAL:HG22	3:L:136:LEU:CD2	2.40	0.51
2:H:74:ARG:HG3	2:H:74:ARG:HH11	1.74	0.51

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:197:TRP:CG	2:H:198:PRO:HA	2.46	0.51
3:L:107:LYS:HD2	4:L:2190:HOH:O	2.11	0.51
2:H:51:ILE:CD1	2:H:74:ARG:HD2	2.42	0.50
2:H:101:ASP:HA	2:H:108:THR:O	2.11	0.50
2:H:74:ARG:NH1	2:H:76:ASN:OD1	2.45	0.50
3:L:180:THR:HG23	4:L:2284:HOH:O	2.11	0.50
2:H:180:GLN:NE2	3:L:180:THR:HG21	2.18	0.49
3:L:30:ARG:HE	3:L:92:ASN:ND2	2.07	0.48
3:L:54:LEU:HG	4:L:2121:HOH:O	2.13	0.48
3:L:54:LEU:HD11	4:L:2128:HOH:O	2.14	0.47
2:H:60:THR:CG2	2:H:62:TYR:CE2	2.97	0.47
2:H:30:THR:HG23	2:H:54:LYS:HB2	1.93	0.47
3:L:210:ASN:ND2	3:L:212:ASN:H	2.11	0.47
3:L:90:GLN:NE2	3:L:92:ASN:H	2.13	0.47
2:H:114:GLN:HB2	4:L:2025:HOH:O	2.15	0.46
2:H:152:LYS:NZ	4:H:2235:HOH:O	2.48	0.46
2:H:51:ILE:HD13	2:H:74:ARG:HD2	1.98	0.46
2:H:141:THR:HA	4:H:2227:HOH:O	2.14	0.45
3:L:193:THR:HG23	3:L:193:THR:O	2.16	0.45
3:L:85:MET:HE1	3:L:103:LYS:HB2	2.00	0.44
3:L:214:CYS:HB3	4:L:2338:HOH:O	2.17	0.43
2:H:60:THR:HG22	2:H:61:GLU:N	2.33	0.43
2:H:74:ARG:CG	2:H:74:ARG:HH11	2.31	0.43
2:H:164:ASN:C	2:H:164:ASN:ND2	2.72	0.43
2:H:138:GLY:N	4:H:2225:HOH:O	2.51	0.43
2:H:169:SER:OG	4:H:2261:HOH:O	2.22	0.42
3:L:115:VAL:HG22	3:L:136:LEU:HD23	2.01	0.42
2:H:139:ASP:OD1	2:H:141:THR:HG23	2.19	0.42
2:H:16:GLY:CA	4:H:2034:HOH:O	2.65	0.42
2:H:53:ASN:C	2:H:53:ASN:HD22	2.23	0.42
2:H:137:CYS:N	4:H:2223:HOH:O	2.53	0.42
3:L:136:LEU:HD21	3:L:196:ALA:HB2	2.02	0.41
3:L:185:GLU:HG3	4:L:2292:HOH:O	2.20	0.41
2:H:200:GLN:HG3	4:H:2299:HOH:O	2.19	0.41
3:L:175:MET:HB3	3:L:175:MET:HE2	1.44	0.41
3:L:151:ASP:OD2	3:L:189:HIS:HD2	2.03	0.41
2:H:174:THR:HG23	4:H:2274:HOH:O	2.20	0.41
3:L:37:ARG:HE	3:L:39:LYS:HZ2	1.66	0.41
3:L:141:PRO:O	3:L:198:HIS:HE1	2.04	0.41
2:H:180:GLN:HE22	3:L:180:THR:CG2	2.24	0.41
2:H:42:GLY:C	4:H:2092:HOH:O	2.59	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:196:THR:HA	4:H:2299:HOH:O	2.20	0.41
2:H:30:THR:HG22	2:H:54:LYS:HB2	2.00	0.40
2:H:11:LEU:HA	2:H:119:THR:OG1	2.21	0.40
3:L:39:LYS:HG3	3:L:84:ALA:HB1	2.03	0.40
2:H:60:THR:CG2	2:H:61:GLU:N	2.84	0.40
3:L:24:ARG:HA	3:L:69:THR:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:L:2039:HOH:O	4:L:2051:HOH:O[2_555]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	4/6 (67%)	4 (100%)	0	0	100	100
2	H	221/222 (100%)	215 (97%)	4 (2%)	2 (1%)	20	4
3	L	213/214 (100%)	207 (97%)	6 (3%)	0	100	100
All	All	438/442 (99%)	426 (97%)	10 (2%)	2 (0%)	32	12

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	136	GLY
2	H	167	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	4/4 (100%)	3 (75%)	1 (25%)	1	0
2	H	184/183 (100%)	172 (94%)	12 (6%)	20	3
3	L	191/190 (100%)	179 (94%)	12 (6%)	21	4
All	All	379/377 (100%)	354 (93%)	25 (7%)	20	3

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	386	THR
2	H	1	GLU
2	H	53	ASN
2	H	67	LYS
2	H	74	ARG
2	H	140	THR
2	H	164	ASN
2	H	167	SER
2	H	170	SER
2	H	181	SER
2	H	185	THR
2	H	197	TRP
2	H	220	GLU
3	L	4	ILE
3	L	11	LEU
3	L	23[A]	CYS
3	L	23[B]	CYS
3	L	90	GLN
3	L	105	GLU
3	L	145	ASN
3	L	160	LEU
3	L	175	MET
3	L	190	ASN
3	L	210	ASN
3	L	214	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	53	ASN
2	H	164	ASN
2	H	180	GLN
3	L	22	ASN
3	L	90	GLN
3	L	92	ASN
3	L	161	ASN
3	L	189	HIS
3	L	198	HIS
3	L	210	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	6/6 (100%)	0.85	1 (16%) <b>2</b> <b>2</b>	12, 19, 25, 33	0
2	H	222/222 (100%)	0.40	19 (8%) <b>11</b> <b>10</b>	7, 13, 37, 54	0
3	L	214/214 (100%)	-0.10	3 (1%) <b>75</b> <b>79</b>	7, 13, 24, 43	0
All	All	442/442 (100%)	0.16	23 (5%) <b>28</b> <b>26</b>	7, 13, 29, 54	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	137	CYS	21.2
2	H	168	LEU	13.0
2	H	166	GLY	10.9
2	H	138	GLY	9.2
2	H	140	THR	8.9
2	H	141	THR	8.5
3	L	214	CYS	7.8
2	H	169	SER	7.7
2	H	139	ASP	7.6
2	H	136	GLY	6.9
2	H	165	SER	4.9
2	H	142	GLY	4.1
1	A	386	THR	3.9
2	H	167	SER	3.8
2	H	41	PRO	3.7
2	H	171	SER	3.4
3	L	202	THR	3.3
2	H	42	GLY	3.1
2	H	1	GLU	2.8
2	H	143	SER	2.6
3	L	213	GLU	2.3
2	H	170	SER	2.2
2	H	43	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.