



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 04:54 pm GMT

PDB ID : 1V5E
Title : Crystal Structure of Pyruvate oxidase containing FAD, from *Aerococcus viridans*
Authors : Hossain, M.T.; Suzuki, K.; Yamamoto, T.; Imamura, S.; Sekiguchi, T.; Takenaka, A.
Deposited on : 2003-11-22
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)
RSRZ outliers	101464	2714 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	590	<div> <div>8%</div> <div>79%</div> <div>20%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5390 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	590	Total	C	N	O	S	181	0	0
			4598	2912	775	891	20			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	719	Total	O	0	0
			719	719		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyruvate oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.66Å 106.12Å 155.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.60 42.17 – 1.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.60) 87.4 (42.17-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 1.39Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.231 , 0.250 0.240 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	14.9	Xtriage
Anisotropy	0.375	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 60.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.37$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5390	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/4694	0.58	1/6373 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	411	SER	N-CA-C	-5.44	96.32	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4598	0	4493	105	0
2	A	20	0	0	0	0
3	A	53	0	31	4	0
4	A	719	0	0	28	0
All	All	5390	0	4524	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:HIS:CE1	1:A:322:ALA:HB2	1.99	0.98
1:A:492:THR:HG22	1:A:493:ASP:H	1.33	0.94
1:A:258:TYR:H	1:A:266:ASN:HD21	1.22	0.85
1:A:221:HIS:NE2	1:A:322:ALA:HB2	1.98	0.76
1:A:60:ALA:HB1	4:A:2119:HOH:O	1.84	0.76
1:A:4:ASN:ND2	1:A:5:LYS:HG3	2.01	0.75
1:A:205:ASN:HD21	1:A:333:ASP:H	1.35	0.72
1:A:259:ARG:HG2	1:A:548:GLU:OE2	1.91	0.71
1:A:320:ASP:OD2	1:A:322:ALA:HB3	1.91	0.71
1:A:82:ALA:HB1	1:A:103:LEU:HG	1.74	0.70
1:A:467:PHE:HA	4:A:1830:HOH:O	1.93	0.69
1:A:258:TYR:H	1:A:266:ASN:ND2	1.90	0.69
1:A:291:PHE:O	1:A:294:VAL:HG22	1.94	0.68
1:A:322:ALA:HB1	4:A:1987:HOH:O	1.94	0.68
1:A:166:ASP:HA	1:A:169:LYS:HE2	1.77	0.67
1:A:368:PHE:CZ	1:A:544:PRO:HG3	2.33	0.63
1:A:137:TYR:HA	4:A:2120:HOH:O	1.99	0.62
1:A:151:MET:SD	1:A:155:LYS:HE2	2.40	0.61
1:A:258:TYR:N	1:A:266:ASN:HD21	1.95	0.61
1:A:8:ILE:HD12	1:A:170:VAL:HB	1.81	0.61
1:A:548:GLU:HG2	1:A:549:THR:HG23	1.83	0.60
1:A:29:SER:HB3	4:A:2047:HOH:O	2.02	0.59
1:A:56:VAL:HG13	1:A:422:PRO:HG3	1.84	0.59
1:A:533:VAL:HG22	4:A:2194:HOH:O	2.03	0.59
1:A:492:THR:HG22	1:A:493:ASP:N	2.12	0.57
1:A:541:GLN:N	1:A:541:GLN:HE21	2.01	0.57
1:A:468:SER:OG	1:A:538:LYS:HD3	2.04	0.57
1:A:136:ALA:H	1:A:140:GLN:NE2	2.04	0.56
1:A:541:GLN:NE2	1:A:541:GLN:N	2.53	0.56
1:A:378:HIS:HE1	1:A:517:ASP:OD2	1.89	0.55
1:A:221:HIS:HE1	4:A:1984:HOH:O	1.89	0.55
1:A:136:ALA:H	1:A:140:GLN:HE21	1.53	0.55
1:A:559:GLU:O	1:A:563:LYS:HG2	2.06	0.54
1:A:571:ALA:HB1	1:A:574:LEU:HD12	1.90	0.54
1:A:263:LYS:HB3	1:A:264:PRO:HD3	1.89	0.53
1:A:535:ILE:HD13	4:A:1818:HOH:O	2.07	0.53
1:A:46:VAL:HB	4:A:2146:HOH:O	2.09	0.53
1:A:459:ASN:C	1:A:461:PRO:HD3	2.29	0.52
1:A:221:HIS:CD2	1:A:322:ALA:HB2	2.44	0.52
1:A:563:LYS:HD3	4:A:1979:HOH:O	2.10	0.52
1:A:566:LYS:HD3	1:A:572:ALA:HA	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:539:ILE:HG23	4:A:1830:HOH:O	2.11	0.50
1:A:541:GLN:HE21	1:A:541:GLN:H	1.56	0.50
1:A:108:GLN:NE2	4:A:2120:HOH:O	2.45	0.49
1:A:39:MET:HE3	1:A:46:VAL:HG12	1.95	0.49
1:A:141:LEU:HB3	1:A:142:PRO:HD3	1.94	0.49
1:A:591:ILE:O	1:A:592:LYS:HB3	2.12	0.49
1:A:470:THR:HG22	1:A:470:THR:O	2.13	0.49
1:A:506:LYS:HG3	1:A:508:PHE:CE1	2.48	0.48
1:A:232:ILE:HG21	1:A:332:VAL:HG21	1.95	0.48
1:A:592:LYS:HG2	4:A:1887:HOH:O	2.12	0.48
1:A:555:LYS:HD3	1:A:592:LYS:HE2	1.94	0.48
1:A:426:GLY:CA	4:A:2119:HOH:O	2.62	0.48
1:A:463:ILE:HA	1:A:533:VAL:HG22	1.95	0.48
1:A:260:VAL:HB	3:A:1601:FAD:C4X	2.44	0.47
1:A:355:ILE:O	1:A:359:GLU:HG3	2.15	0.47
1:A:411:SER:O	1:A:415:ALA:HA	2.15	0.47
1:A:518:ARG:O	1:A:522:GLU:HG3	2.13	0.47
1:A:568:ARG:HG3	1:A:568:ARG:HH11	1.78	0.47
1:A:393:THR:HG22	1:A:439:ILE:HG22	1.96	0.47
1:A:354:TYR:O	1:A:357:MET:HG3	2.14	0.47
1:A:10:LEU:CD2	1:A:42:GLU:HG3	2.45	0.47
1:A:519:VAL:HG12	4:A:1818:HOH:O	2.13	0.47
1:A:82:ALA:CB	1:A:103:LEU:HG	2.43	0.47
1:A:256:SER:OG	1:A:261:GLY:HA2	2.15	0.46
1:A:357:MET:HG3	1:A:358:LEU:N	2.29	0.46
1:A:259:ARG:HG2	1:A:548:GLU:CD	2.36	0.46
1:A:551:LYS:HE3	4:A:1936:HOH:O	2.15	0.46
1:A:527:ASN:ND2	4:A:2194:HOH:O	2.48	0.46
1:A:205:ASN:ND2	1:A:332:VAL:HG23	2.31	0.46
1:A:566:LYS:HD2	4:A:1979:HOH:O	2.15	0.46
1:A:47:LYS:HE2	4:A:2210:HOH:O	2.16	0.45
1:A:34:SER:HB3	1:A:168:ALA:HB3	1.97	0.45
1:A:5:LYS:O	1:A:5:LYS:HD2	2.17	0.45
1:A:415:ALA:HB2	3:A:1601:FAD:HM82	1.99	0.44
1:A:63:MET:SD	1:A:66:LYS:HD2	2.57	0.44
1:A:41:GLU:HB2	4:A:2164:HOH:O	2.17	0.44
1:A:5:LYS:HD2	1:A:5:LYS:C	2.38	0.44
1:A:426:GLY:HA2	4:A:2119:HOH:O	2.17	0.44
1:A:543:ARG:NH2	1:A:546:PRO:HB3	2.32	0.44
1:A:10:LEU:HD13	4:A:2164:HOH:O	2.16	0.43
1:A:221:HIS:CD2	1:A:322:ALA:CB	3.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:HIS:NE2	1:A:322:ALA:CB	2.75	0.43
1:A:533:VAL:HG13	4:A:2194:HOH:O	2.18	0.43
1:A:322:ALA:HB3	4:A:1718:HOH:O	2.18	0.43
1:A:173:ASP:HB3	1:A:175:ASP:OD1	2.19	0.43
1:A:387:ILE:HG13	1:A:408:TRP:CE2	2.53	0.43
1:A:44:ASN:ND2	4:A:2146:HOH:O	2.48	0.43
1:A:563:LYS:HE2	4:A:2142:HOH:O	2.18	0.43
1:A:421:ILE:HB	1:A:422:PRO:HD3	2.00	0.43
1:A:262:TRP:HB3	1:A:264:PRO:HD2	2.01	0.43
1:A:10:LEU:HD22	1:A:42:GLU:HG3	2.01	0.42
1:A:260:VAL:N	3:A:1601:FAD:O4	2.49	0.42
1:A:32:LEU:HB2	1:A:36:MET:HE1	2.01	0.42
1:A:387:ILE:HG13	1:A:408:TRP:NE1	2.35	0.42
1:A:539:ILE:HG12	4:A:1830:HOH:O	2.18	0.42
1:A:173:ASP:HB2	1:A:176:GLN:HG2	2.01	0.41
1:A:176:GLN:NE2	1:A:176:GLN:HA	2.34	0.41
1:A:442:ASP:OD2	1:A:468:SER:HA	2.21	0.41
1:A:514:GLU:HG3	4:A:2257:HOH:O	2.21	0.41
1:A:86:ILE:HG23	1:A:87:ASN:N	2.36	0.41
1:A:412:PRO:HB2	3:A:1601:FAD:O3'	2.21	0.41
1:A:230:ARG:HE	1:A:250:PHE:HE1	1.69	0.41
1:A:25:TYR:CE2	1:A:49:LEU:HD13	2.56	0.40
1:A:332:VAL:HG22	1:A:333:ASP:N	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	588/590 (100%)	557 (95%)	29 (5%)	2 (0%)	44 22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	487	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	483/483 (100%)	473 (98%)	10 (2%)	59	32

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	45	ASN
1	A	84	HIS
1	A	213	TYR
1	A	330	ASN
1	A	357	MET
1	A	388	ASP
1	A	391	ASN
1	A	451	ASP
1	A	541	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	44	ASN
1	A	45	ASN
1	A	64	GLN
1	A	70	ASN
1	A	96	ASN
1	A	140	GLN
1	A	176	GLN
1	A	195	GLN
1	A	205	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	206	ASN
1	A	221	HIS
1	A	266	ASN
1	A	296	ASN
1	A	330	ASN
1	A	378	HIS
1	A	391	ASN
1	A	429	ASN
1	A	541	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FAD	A	1601	-	51,58,58	1.10	4 (7%)	54,89,89	1.96	5 (9%)
2	SO4	A	1603	-	4,4,4	0.30	0	6,6,6	0.09	0
2	SO4	A	1604	-	4,4,4	0.31	0	6,6,6	0.09	0
2	SO4	A	1605	-	4,4,4	0.27	0	6,6,6	0.11	0
2	SO4	A	1606	-	4,4,4	0.30	0	6,6,6	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	1601	-	-	0/28/50/50	0/6/6/6
2	SO4	A	1603	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1604	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1605	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1606	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1601	FAD	C9A-N10	2.30	1.41	1.38
3	A	1601	FAD	C4-N3	2.47	1.37	1.33
3	A	1601	FAD	C10-N1	2.56	1.36	1.33
3	A	1601	FAD	C4X-N5	3.81	1.38	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1601	FAD	C4X-C4-N3	-4.62	116.90	123.48
3	A	1601	FAD	C5X-C9A-N10	2.29	119.36	117.66
3	A	1601	FAD	C4X-N5-C5X	3.37	120.32	116.76
3	A	1601	FAD	C1'-N10-C9A	4.53	122.50	118.35
3	A	1601	FAD	C4-N3-C2	10.39	124.24	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1601	FAD	4	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	569/590 (96%)	1.08	46 (8%) 13 12	7, 12, 25, 48	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	470	THR	10.3
1	A	492	THR	9.7
1	A	3	ASP	7.7
1	A	469	ASN	6.1
1	A	4	ASN	5.1
1	A	41	GLU	4.9
1	A	541	GLN	4.2
1	A	36	MET	3.9
1	A	493	ASP	3.7
1	A	45	ASN	3.4
1	A	40	GLY	3.4
1	A	330	ASN	3.4
1	A	560	ASP	3.3
1	A	592	LYS	3.1
1	A	37	ASP	3.0
1	A	557	TYR	3.0
1	A	260	VAL	3.0
1	A	336	GLU	2.8
1	A	563	LYS	2.7
1	A	48	PHE	2.6
1	A	237	ILE	2.4
1	A	39	MET	2.4
1	A	568	ARG	2.2
1	A	202	GLU	2.2
1	A	176	GLN	2.1
1	A	282	PHE	2.1
1	A	5	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	195	GLN	2.1
1	A	518	ARG	2.1
1	A	335	VAL	2.1
1	A	413	LEU	2.1
1	A	174	ASN	2.1
1	A	512	ARG	2.1
1	A	397	ILE	2.1
1	A	391	ASN	2.1
1	A	363	GLU	2.1
1	A	11	ALA	2.1
1	A	322	ALA	2.1
1	A	524	VAL	2.1
1	A	549	THR	2.0
1	A	324	ALA	2.0
1	A	170	VAL	2.0
1	A	387	ILE	2.0
1	A	457	ARG	2.0
1	A	103	LEU	2.0
1	A	410	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SO4	A	1606	5/5	0.89	0.19	1.20	18,21,22,23	0
3	FAD	A	1601	53/53	0.88	0.16	0.40	7,10,16,22	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SO4	A	1603	5/5	0.93	0.10	-2.71	15,17,17,18	0
2	SO4	A	1604	5/5	0.81	0.27	-	36,36,36,36	0
2	SO4	A	1605	5/5	0.87	0.24	-	26,27,27,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.