



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 07:48 am GMT

PDB ID : 2V58  
Title : CRYSTAL STRUCTURE OF BIOTIN CARBOXYLASE FROM E.COLI IN  
COMPLEX WITH POTENT INHIBITOR 1  
Authors : Mochalkin, I.; Miller, J.R.  
Deposited on : 2008-10-02  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	449	<div> <div>%</div> <div>97%</div> <div>.</div> </div>
1	B	449	<div> <div>8%</div> <div>94%</div> <div>5% .</div> </div>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 7485 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BIOTIN CARBOXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	0	5	1
			3449	2173	616	638	22			
1	B	446	Total	C	N	O	S	0	6	1
			3447	2168	618	639	22			

- Molecule 2 is 6-(2,6-DIBROMOPHENYL)PYRIDO[2,3-D]PYRIMIDINE-2,7-DIAMINE (three-letter code: LZJ) (formula: C<sub>13</sub>H<sub>9</sub>Br<sub>2</sub>N<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	Br	C	N	0	0
			20	2	13	5		
2	B	1	Total	Br	C	N	0	0
			20	2	13	5		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Cl 1	0	0
3	A	1	Total 1	Cl 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	316	Total 316	O 316	0	0
4	B	231	Total 231	O 231	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: BIOTIN CARBOXYLASE



#### • Molecule 1: BIOTIN CARBOXYLASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.21Å 106.61Å 122.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	80.32 – 2.10 48.87 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.5 (80.32-2.10) 94.2 (48.87-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.46 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.193 , 0.219 0.202 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	25.1	Xtriage
Anisotropy	0.249	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7485	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LZJ, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/3537	0.47	0/4772
1	B	0.33	0/3540	0.46	0/4777
All	All	0.33	0/7077	0.47	0/9549

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3449	0	3485	6	0
1	B	3447	0	3473	15	0
2	A	20	0	9	0	0
2	B	20	0	9	1	0
3	A	1	0	0	0	0
3	B	1	0	0	1	0
4	A	316	0	0	0	0
4	B	231	0	0	0	0
All	All	7485	0	6976	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:292:ARG:NH1	3:B:1448:CL:CL	2.70	0.62
1:A:5:ILE:HG22	1:A:77:ALA:HB3	1.89	0.53
1:A:340:ASN:HD22	1:A:384:MET:HA	1.73	0.53
1:B:105:PRO:HG2	1:B:291:THR:HB	1.93	0.50
1:B:232:MET:CE	1:B:440:LEU:HB2	2.44	0.48
1:A:79:HIS:HE2	1:A:312:GLN:NE2	2.11	0.48
1:A:71:GLU:HG2	1:A:100:PHE:HZ	1.77	0.48
1:A:298:PRO:O	1:A:302:MET:HG2	2.15	0.47
1:B:5:ILE:HG22	1:B:77:ALA:HB3	1.96	0.47
1:B:252:LEU:HG	1:B:284:PHE:HE1	1.80	0.46
1:B:298:PRO:O	1:B:302:MET:HG2	2.17	0.45
1:B:232:MET:HE2	1:B:440:LEU:HB2	1.99	0.44
1:A:297:HIS:N	1:A:298:PRO:CD	2.81	0.44
1:B:232:MET:HE2	1:B:232:MET:HA	1.99	0.44
1:B:133:GLY:HA2	1:B:152:ILE:HD11	2.00	0.43
1:B:274:THR:HG21	1:B:294:GLN:HE22	1.85	0.42
1:B:297:HIS:N	1:B:298:PRO:CD	2.82	0.42
1:B:79:HIS:HE2	1:B:312:GLN:NE2	2.17	0.42
1:B:152:ILE:HG23	1:B:202:LYS:HB2	2.01	0.42
1:B:290:ASN:ND2	1:B:294:GLN:OE1	2.53	0.41
1:B:233:GLN:CD	2:B:1447:LZJ:H16	2.41	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	450/449 (100%)	442 (98%)	8 (2%)	0	100	100
1	B	450/449 (100%)	441 (98%)	9 (2%)	0	100	100
All	All	900/898 (100%)	883 (98%)	17 (2%)	0	100	100



There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/361 (101%)	361 (99%)	2 (1%)	89	92
1	B	363/361 (101%)	361 (99%)	2 (1%)	89	92
All	All	726/722 (101%)	722 (99%)	4 (1%)	89	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	237	GLN
1	A	323	ILE
1	B	441	GLU
1	B	442	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	180	GLN
1	A	206	ASN
1	A	312	GLN
1	A	319	GLN
1	A	340	ASN
1	A	404	ASN
1	B	290	ASN
1	B	294	GLN
1	B	312	GLN
1	B	319	GLN
1	B	404	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	LZJ	A	1448	-	22,22,22	0.84	0	28,32,32	2.03	9 (32%)
2	LZJ	B	1447	-	22,22,22	0.87	0	28,32,32	2.05	9 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LZJ	A	1448	-	-	0/4/4/4	0/3/3/3
2	LZJ	B	1447	-	-	0/4/4/4	0/3/3/3

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1448	LZJ	N1-C2-N3	-5.26	120.33	125.82
2	B	1447	LZJ	N1-C2-N3	-5.00	120.59	125.82
2	B	1447	LZJ	C5-C4-N3	-3.03	119.68	122.48
2	A	1448	LZJ	C5-C4-N3	-2.80	119.89	122.48
2	A	1448	LZJ	C5-C4-N7	-2.40	120.26	122.48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1447	LZJ	C5-C4-N7	-2.26	120.38	122.48
2	B	1447	LZJ	C5-C6-N1	-2.03	119.68	123.86
2	A	1448	LZJ	C9-C10-C5	-2.01	118.98	121.92
2	B	1447	LZJ	C6-N1-C2	2.55	120.70	116.26
2	A	1448	LZJ	C6-N1-C2	2.70	120.96	116.26
2	A	1448	LZJ	C14-C12-C18	2.91	118.70	115.53
2	B	1447	LZJ	N11-C2-N1	3.11	120.38	117.39
2	A	1448	LZJ	N11-C2-N1	3.15	120.41	117.39
2	A	1448	LZJ	C8-N7-C4	3.36	121.04	117.61
2	B	1447	LZJ	C14-C12-C18	3.46	119.30	115.53
2	B	1447	LZJ	C8-N7-C4	3.54	121.23	117.61
2	B	1447	LZJ	C2-N3-C4	4.59	120.51	115.16
2	A	1448	LZJ	C2-N3-C4	4.63	120.57	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1447	LZJ	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	447/449 (99%)	-0.25	3 (0%) 87 89	18, 22, 30, 41	0
1	B	446/449 (99%)	0.24	36 (8%) 13 16	18, 24, 49, 52	0
All	All	893/898 (99%)	-0.00	39 (4%) 35 41	18, 23, 41, 52	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	180	GLN	5.6
1	B	142	MET	5.1
1	B	154	TYR	4.8
1	A	446	LEU	4.4
1	B	172	VAL	4.3
1	B	183	SER	4.3
1	B	445	GLY	4.2
1	B	146	ARG	4.2
1	B	179	ALA	4.1
1	B	143	ASP	4.1
1	A	165	GLY	4.0
1	B	181	SER	3.9
1	B	141	ASP	3.7
1	B	176	ALA	3.7
1	B	152	ILE	3.7
1	B	177	GLU	3.7
1	B	150	LYS	3.6
1	B	153	GLY	3.5
1	B	148	ILE	3.3
1	B	174	GLY	3.3
1	B	184	MET	3.3
1	B	140	ASP	3.1
1	A	447	GLN	3.1
1	B	167	ARG	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	175	ASP	2.9
1	B	281	ASN	2.9
1	B	144	LYS	2.9
1	B	178	LEU	2.9
1	B	168	GLY	2.8
1	B	192	ALA	2.6
1	B	139	GLY	2.6
1	B	187	ALA	2.6
1	B	138	LEU	2.5
1	B	185	THR	2.5
1	B	151	ARG	2.4
1	B	173	ARG	2.3
1	B	137	PRO	2.2
1	B	282	GLY	2.2
1	B	254	ARG	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	LZJ	B	1447	20/20	0.97	0.11	-0.40	28,28,29,30	0
2	LZJ	A	1448	20/20	0.97	0.08	-0.46	25,26,28,30	0
3	CL	A	1449	1/1	0.99	0.06	-5.65	19,19,19,19	0
3	CL	B	1448	1/1	0.99	0.02	-6.15	25,25,25,25	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.