



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 03:58 am GMT

PDB ID : 3V8G
Title : Crystal structure of an asymmetric trimer of a glutamate transporter homologue (GltPh)
Authors : Verdon, G.; Boudker, O.
Deposited on : 2011-12-22
Resolution : 4.66 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1008 (5.66-3.66)
Clashscore	112137	1055 (5.60-3.70)
Ramachandran outliers	110173	1000 (5.58-3.68)
Sidechain outliers	110143	1024 (5.66-3.66)
RSRZ outliers	101464	1017 (5.66-3.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>14%</div> <div>68% 27% . .</div> </div>
1	B	422	<div> <div>11%</div> <div>67% 27% . .</div> </div>
1	C	422	<div> <div>7%</div> <div>52% 37% 6% 5%</div> </div>
1	D	422	<div> <div>12%</div> <div>68% 27% . .</div> </div>
1	E	422	<div> <div>10%</div> <div>68% 26% . 5%</div> </div>
1	F	422	<div> <div>6%</div> <div>52% 34% 7% 7%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 18038 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called sodium-coupled L-aspartate transporter.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	0	0	0
			3050	2007	492	533	18			
1	B	404	Total	C	N	O	S	0	0	0
			3001	1976	483	524	18			
1	C	400	Total	C	N	O	S	0	0	0
			2965	1955	475	518	17			
1	D	411	Total	C	N	O	S	0	0	0
			3050	2007	492	533	18			
1	E	403	Total	C	N	O	S	0	0	0
			2992	1971	481	522	18			
1	F	392	Total	C	N	O	S	0	0	0
			2914	1922	465	510	17			

There are 90 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	37	HIS	ASP	engineered mutation	UNP O59010
A	40	HIS	LYS	engineered mutation	UNP O59010
A	125	HIS	LYS	engineered mutation	UNP O59010
A	132	HIS	LYS	engineered mutation	UNP O59010
A	198	CYS	VAL	engineered mutation	UNP O59010
A	223	HIS	LYS	engineered mutation	UNP O59010
A	264	HIS	LYS	engineered mutation	UNP O59010
A	321	ALA	CYS	engineered mutation	UNP O59010
A	368	HIS	GLU	engineered mutation	UNP O59010
A	380	CYS	ALA	engineered mutation	UNP O59010
A	418	THR	-	EXPRESSION TAG	UNP O59010
A	419	LEU	-	EXPRESSION TAG	UNP O59010
A	420	VAL	-	EXPRESSION TAG	UNP O59010
A	421	PRO	-	EXPRESSION TAG	UNP O59010
A	422	ARG	-	EXPRESSION TAG	UNP O59010
B	37	HIS	ASP	engineered mutation	UNP O59010
B	40	HIS	LYS	engineered mutation	UNP O59010

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	125	HIS	LYS	engineered mutation	UNP O59010
B	132	HIS	LYS	engineered mutation	UNP O59010
B	198	CYS	VAL	engineered mutation	UNP O59010
B	223	HIS	LYS	engineered mutation	UNP O59010
B	264	HIS	LYS	engineered mutation	UNP O59010
B	321	ALA	CYS	engineered mutation	UNP O59010
B	368	HIS	GLU	engineered mutation	UNP O59010
B	380	CYS	ALA	engineered mutation	UNP O59010
B	418	THR	-	EXPRESSION TAG	UNP O59010
B	419	LEU	-	EXPRESSION TAG	UNP O59010
B	420	VAL	-	EXPRESSION TAG	UNP O59010
B	421	PRO	-	EXPRESSION TAG	UNP O59010
B	422	ARG	-	EXPRESSION TAG	UNP O59010
C	37	HIS	ASP	engineered mutation	UNP O59010
C	40	HIS	LYS	engineered mutation	UNP O59010
C	125	HIS	LYS	engineered mutation	UNP O59010
C	132	HIS	LYS	engineered mutation	UNP O59010
C	198	CYS	VAL	engineered mutation	UNP O59010
C	223	HIS	LYS	engineered mutation	UNP O59010
C	264	HIS	LYS	engineered mutation	UNP O59010
C	321	ALA	CYS	engineered mutation	UNP O59010
C	368	HIS	GLU	engineered mutation	UNP O59010
C	380	CYS	ALA	engineered mutation	UNP O59010
C	418	THR	-	EXPRESSION TAG	UNP O59010
C	419	LEU	-	EXPRESSION TAG	UNP O59010
C	420	VAL	-	EXPRESSION TAG	UNP O59010
C	421	PRO	-	EXPRESSION TAG	UNP O59010
C	422	ARG	-	EXPRESSION TAG	UNP O59010
D	37	HIS	ASP	engineered mutation	UNP O59010
D	40	HIS	LYS	engineered mutation	UNP O59010
D	125	HIS	LYS	engineered mutation	UNP O59010
D	132	HIS	LYS	engineered mutation	UNP O59010
D	198	CYS	VAL	engineered mutation	UNP O59010
D	223	HIS	LYS	engineered mutation	UNP O59010
D	264	HIS	LYS	engineered mutation	UNP O59010
D	321	ALA	CYS	engineered mutation	UNP O59010
D	368	HIS	GLU	engineered mutation	UNP O59010
D	380	CYS	ALA	engineered mutation	UNP O59010
D	418	THR	-	EXPRESSION TAG	UNP O59010
D	419	LEU	-	EXPRESSION TAG	UNP O59010
D	420	VAL	-	EXPRESSION TAG	UNP O59010
D	421	PRO	-	EXPRESSION TAG	UNP O59010

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	422	ARG	-	EXPRESSION TAG	UNP O59010
E	37	HIS	ASP	engineered mutation	UNP O59010
E	40	HIS	LYS	engineered mutation	UNP O59010
E	125	HIS	LYS	engineered mutation	UNP O59010
E	132	HIS	LYS	engineered mutation	UNP O59010
E	198	CYS	VAL	engineered mutation	UNP O59010
E	223	HIS	LYS	engineered mutation	UNP O59010
E	264	HIS	LYS	engineered mutation	UNP O59010
E	321	ALA	CYS	engineered mutation	UNP O59010
E	368	HIS	GLU	engineered mutation	UNP O59010
E	380	CYS	ALA	engineered mutation	UNP O59010
E	418	THR	-	EXPRESSION TAG	UNP O59010
E	419	LEU	-	EXPRESSION TAG	UNP O59010
E	420	VAL	-	EXPRESSION TAG	UNP O59010
E	421	PRO	-	EXPRESSION TAG	UNP O59010
E	422	ARG	-	EXPRESSION TAG	UNP O59010
F	37	HIS	ASP	engineered mutation	UNP O59010
F	40	HIS	LYS	engineered mutation	UNP O59010
F	125	HIS	LYS	engineered mutation	UNP O59010
F	132	HIS	LYS	engineered mutation	UNP O59010
F	198	CYS	VAL	engineered mutation	UNP O59010
F	223	HIS	LYS	engineered mutation	UNP O59010
F	264	HIS	LYS	engineered mutation	UNP O59010
F	321	ALA	CYS	engineered mutation	UNP O59010
F	368	HIS	GLU	engineered mutation	UNP O59010
F	380	CYS	ALA	engineered mutation	UNP O59010
F	418	THR	-	EXPRESSION TAG	UNP O59010
F	419	LEU	-	EXPRESSION TAG	UNP O59010
F	420	VAL	-	EXPRESSION TAG	UNP O59010
F	421	PRO	-	EXPRESSION TAG	UNP O59010
F	422	ARG	-	EXPRESSION TAG	UNP O59010

- Molecule 2 is ASPARTIC ACID (three-letter code: ASP) (formula: C₄H₇NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			9	4	1	4		
2	B	1	Total	C	N	O	0	0
			9	4	1	4		
2	C	1	Total	C	N	O	0	0
			9	4	1	4		
2	D	1	Total	C	N	O	0	0
			9	4	1	4		
2	E	1	Total	C	N	O	0	0
			9	4	1	4		
2	F	1	Total	C	N	O	0	0
			9	4	1	4		


- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	2	Total	Na	0	0
			2	2		
3	E	2	Total	Na	0	0
			2	2		
3	B	2	Total	Na	0	0
			2	2		
3	C	2	Total	Na	0	0
			2	2		
3	A	2	Total	Na	0	0
			2	2		
3	F	2	Total	Na	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: sodium-coupled L-aspartate transporter




- Molecule 1: sodium-coupled L-aspartate transporter







• Molecule 1: sodium-coupled L-aspartate transporter



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	203.69Å 119.63Å 223.41Å 90.00° 113.77° 90.00°	Depositor
Resolution (Å)	15.00 – 4.66 68.15 – 4.66	Depositor EDS
% Data completeness (in resolution range)	73.1 (15.00-4.66) 66.2 (68.15-4.66)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.67 (at 4.65Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.255 , 0.294 0.291 , 0.311	Depositor DCC
R_{free} test set	1028 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	255.4	Xtriage
Anisotropy	0.256	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 436.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	18038	wwPDB-VP
Average B, all atoms (Å ²)	283.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/3110	0.54	0/4241
1	B	0.37	0/3059	0.53	0/4171
1	C	0.41	0/3020	0.61	0/4116
1	D	0.37	0/3110	0.53	0/4241
1	E	0.37	0/3050	0.53	0/4158
1	F	0.40	0/2968	0.60	0/4045
All	All	0.38	0/18317	0.56	0/24972

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2
1	F	0	2
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	223	HIS	Peptide
1	C	224	VAL	Peptide
1	F	223	HIS	Peptide
1	F	224	VAL	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3050	0	3223	81	0
1	B	3001	0	3177	79	0
1	C	2965	0	3147	129	0
1	D	3050	0	3223	82	0
1	E	2992	0	3170	74	0
1	F	2914	0	3093	121	0
2	A	9	0	3	0	0
2	B	9	0	3	0	0
2	C	9	0	3	0	0
2	D	9	0	3	0	0
2	E	9	0	3	0	0
2	F	9	0	3	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
All	All	18038	0	19051	532	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 532 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:243:ILE:HA	1:D:247:TYR:HD2	1.30	0.96
1:E:243:ILE:HA	1:E:247:TYR:HD2	1.30	0.95
1:B:243:ILE:HA	1:B:247:TYR:HD2	1.29	0.95
1:A:243:ILE:HA	1:A:247:TYR:HD2	1.31	0.92
1:C:134:LEU:O	1:C:137:ILE:HG12	1.71	0.90

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/422 (97%)	357 (87%)	44 (11%)	8 (2%)	9	48
1	B	400/422 (95%)	350 (88%)	44 (11%)	6 (2%)	12	54
1	C	396/422 (94%)	302 (76%)	75 (19%)	19 (5%)	2	29
1	D	409/422 (97%)	356 (87%)	44 (11%)	9 (2%)	8	46
1	E	399/422 (94%)	350 (88%)	42 (10%)	7 (2%)	10	50
1	F	388/422 (92%)	296 (76%)	73 (19%)	19 (5%)	2	28
All	All	2401/2532 (95%)	2011 (84%)	322 (13%)	68 (3%)	6	41

5 of 68 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	358	ALA
1	A	370	VAL
1	B	358	ALA
1	B	370	VAL
1	C	75	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/330 (97%)	309 (96%)	12 (4%)	39	69
1	B	317/330 (96%)	305 (96%)	12 (4%)	38	68
1	C	312/330 (94%)	281 (90%)	31 (10%)	9	36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	321/330 (97%)	308 (96%)	13 (4%)	36	66
1	E	316/330 (96%)	304 (96%)	12 (4%)	38	68
1	F	308/330 (93%)	279 (91%)	29 (9%)	10	39
All	All	1895/1980 (96%)	1786 (94%)	109 (6%)	24	57

5 of 109 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	342	VAL
1	D	178	LYS
1	F	317	TYR
1	C	343	LEU
1	D	12	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	108	ASN
1	D	337	GLN
1	F	40	HIS
1	C	337	GLN
1	F	108	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 18 ligands modelled in this entry, 12 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ASP	A	501	-	1,8,8	0.20	0	1,10,10	0.54	0
2	ASP	B	501	-	1,8,8	0.26	0	1,10,10	0.62	0
2	ASP	C	501	-	1,8,8	0.17	0	1,10,10	0.58	0
2	ASP	D	501	-	1,8,8	0.23	0	1,10,10	0.55	0
2	ASP	E	501	-	1,8,8	0.24	0	1,10,10	0.56	0
2	ASP	F	501	-	1,8,8	0.17	0	1,10,10	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ASP	A	501	-	-	0/2/8/8	0/0/0/0
2	ASP	B	501	-	-	0/2/8/8	0/0/0/0
2	ASP	C	501	-	-	0/2/8/8	0/0/0/0
2	ASP	D	501	-	-	0/2/8/8	0/0/0/0
2	ASP	E	501	-	-	0/2/8/8	0/0/0/0
2	ASP	F	501	-	-	0/2/8/8	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	411/422 (97%)	0.65	58 (14%) 3 6	216, 306, 404, 449	0
1	B	404/422 (95%)	0.48	48 (11%) 5 8	198, 281, 385, 436	0
1	C	400/422 (94%)	0.36	30 (7%) 15 14	192, 246, 347, 443	0
1	D	411/422 (97%)	0.46	50 (12%) 5 7	213, 290, 399, 435	0
1	E	403/422 (95%)	0.42	41 (10%) 7 9	208, 264, 369, 402	0
1	F	392/422 (92%)	0.26	26 (6%) 19 17	200, 246, 348, 452	0
All	All	2421/2532 (95%)	0.44	253 (10%) 7 8	192, 274, 386, 452	0

The worst 5 of 253 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	256	ILE	16.3
1	C	36	ALA	9.6
1	B	124	PRO	9.6
1	A	36	ALA	9.3
1	A	257	ASP	8.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ASP	A	501	9/9	0.81	0.18	-0.41	263,269,278,280	0
2	ASP	D	501	9/9	0.84	0.14	-0.55	258,265,274,278	0
2	ASP	F	501	9/9	0.49	0.24	-0.59	209,210,213,215	0
3	NA	D	503	1/1	0.97	0.17	-0.65	269,269,269,269	0
2	ASP	E	501	9/9	0.91	0.14	-0.66	220,226,234,237	0
2	ASP	C	501	9/9	0.93	0.18	-0.69	203,205,206,210	0
3	NA	A	503	1/1	0.91	0.34	-0.72	270,270,270,270	0
3	NA	D	502	1/1	0.92	0.13	-0.75	281,281,281,281	0
3	NA	B	503	1/1	0.85	0.18	-0.85	253,253,253,253	0
3	NA	A	502	1/1	0.73	0.10	-0.86	283,283,283,283	0
2	ASP	B	501	9/9	0.56	0.20	-0.90	244,250,260,262	0
3	NA	C	503	1/1	0.97	0.21	-1.01	195,195,195,195	0
3	NA	E	502	1/1	0.86	0.07	-1.05	245,245,245,245	0
3	NA	B	502	1/1	0.81	0.13	-1.08	264,264,264,264	0
3	NA	E	503	1/1	0.85	0.14	-1.16	241,241,241,241	0
3	NA	C	502	1/1	0.93	0.13	-1.19	193,193,193,193	0
3	NA	F	502	1/1	0.99	0.13	-1.23	206,206,206,206	0
3	NA	F	503	1/1	0.96	0.16	-1.62	203,203,203,203	0

6.5 Other polymers [i](#)

There are no such residues in this entry.