



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 11:41 pm GMT

PDB ID : 2VC0  
Title : FEAST OR FAMINE REGULATORY PROTEIN (RV3291C)FROM M. TUBERCULOSIS COMPLEXED WITH L-LEUCINE  
Authors : Shrivastava, T.; Ramachandran, R.  
Deposited on : 2007-09-18  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	150	 2% 68% 19% 11% •
1	B	150	 3% 55% 33% 8% ••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	LEU	A	1151	-	-	-	X

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2293 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTIONAL REGULATORY PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	147	Total	C	N	O	S	0	0	0
			1127	697	209	220	1			
1	B	147	Total	C	N	O	S	0	0	0
			1111	689	203	218	1			

- Molecule 2 is LEUCINE (three-letter code: LEU) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			9	6	1	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	25	Total	O	0	0
			25	25		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	21	Total	O	0	0
			21	21		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: TRANSCRIPTIONAL REGULATORY PROTEIN



#### • Molecule 1: TRANSCRIPTIONAL REGULATORY PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.88Å 100.88Å 99.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.25 – 2.50 24.47 – 2.50	Depositor EDS
% Data completeness (in resolution range)	95.0 (71.25-2.50) 95.1 (24.47-2.50)	Depositor EDS
$R_{merge}$	0.95	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.10 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.212 , 0.257 0.191 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	44.1	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 47.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.021 for -h,-l,-k 0.014 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2293	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.36	6/1141 (0.5%)	1.35	18/1552 (1.2%)
1	B	1.26	5/1125 (0.4%)	1.26	7/1533 (0.5%)
All	All	1.31	11/2266 (0.5%)	1.31	25/3085 (0.8%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	146	ARG	CD-NE	-7.98	1.32	1.46
1	A	146	ARG	CB-CG	-6.21	1.35	1.52
1	B	98	TYR	CD1-CE1	6.13	1.48	1.39
1	B	134	ARG	CG-CD	6.04	1.67	1.51
1	A	98	TYR	CD1-CE1	5.80	1.48	1.39
1	A	143	TYR	CD2-CE2	5.48	1.47	1.39
1	B	55	SER	CA-CB	5.21	1.60	1.52
1	A	114	SER	CB-OG	5.19	1.49	1.42
1	B	86	ALA	CA-CB	5.08	1.63	1.52
1	A	41	ARG	CG-CD	-5.06	1.39	1.51
1	B	98	TYR	CE1-CZ	5.03	1.45	1.38

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	ARG	CB-CA-C	7.68	125.77	110.40
1	B	100	VAL	CG1-CB-CG2	7.33	122.63	110.90
1	B	146	ARG	NE-CZ-NH1	7.28	123.94	120.30
1	B	130	ASN	CB-CA-C	7.22	124.83	110.40
1	A	110	VAL	CG1-CB-CG2	6.83	121.83	110.90
1	A	111	ARG	NE-CZ-NH2	-6.54	117.03	120.30
1	B	129	ALA	C-N-CA	-6.54	105.36	121.70
1	A	48	ARG	C-N-CA	-6.53	108.58	122.30
1	A	100	VAL	CG1-CB-CG2	6.29	120.97	110.90
1	A	111	ARG	NE-CZ-NH1	6.21	123.41	120.30
1	B	77	ASP	CB-CG-OD2	-6.13	112.78	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	77	ASP	CB-CG-OD2	-6.12	112.79	118.30
1	A	131	VAL	CG1-CB-CG2	5.70	120.02	110.90
1	A	146	ARG	CB-CG-CD	-5.70	96.78	111.60
1	A	146	ARG	NE-CZ-NH2	-5.69	117.46	120.30
1	A	146	ARG	N-CA-CB	-5.62	100.49	110.60
1	A	146	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	B	146	ARG	NE-CZ-NH2	-5.60	117.50	120.30
1	A	131	VAL	N-CA-CB	-5.60	99.19	111.50
1	B	21	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	A	120	ASP	CB-CG-OD1	-5.51	113.34	118.30
1	A	120	ASP	CB-CG-OD2	5.21	122.99	118.30
1	A	126	ARG	NE-CZ-NH1	5.19	122.90	120.30
1	A	14	ARG	NE-CZ-NH1	-5.10	117.75	120.30
1	A	146	ARG	CA-CB-CG	5.00	124.41	113.40

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1127	0	1130	25	0
1	B	1111	0	1104	48	0
2	A	9	0	10	1	0
3	A	25	0	0	3	0
3	B	21	0	0	1	0
All	All	2293	0	2244	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:ASP:HB3	3:A:2024:HOH:O	1.47	1.14

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:44:ARG:HB3	1:B:44:ARG:HH11	1.16	1.04
1:B:72:ALA:HB1	1:B:104:GLU:HG3	1.51	0.91
1:B:104:GLU:HG2	1:B:134:ARG:NH2	1.85	0.89
1:B:44:ARG:HB3	1:B:44:ARG:NH1	1.89	0.87
1:A:124:ARG:O	1:A:128:THR:HG23	1.81	0.80
1:B:33:LEU:HD22	1:B:37:ALA:CB	2.14	0.78
1:B:91:ILE:HD12	1:B:93:GLU:OE1	1.87	0.75
1:B:6:ASP:O	1:B:9:ASP:HB2	1.90	0.71
1:A:146:ARG:NH2	1:B:89:GLU:O	2.24	0.70
1:B:63:VAL:HG12	1:B:63:VAL:O	1.95	0.67
1:A:94:VAL:O	1:B:146:ARG:NH1	2.29	0.66
1:B:124:ARG:O	1:B:128:THR:HG23	1.95	0.65
1:B:42:VAL:O	1:B:46:GLU:HG3	1.98	0.63
1:B:8:ILE:HG21	1:B:41:ARG:HH21	1.63	0.63
1:B:104:GLU:HG2	1:B:134:ARG:HH21	1.62	0.62
1:B:91:ILE:HD11	1:B:93:GLU:HG2	1.82	0.62
1:B:91:ILE:CD1	1:B:93:GLU:OE1	2.49	0.61
1:A:24:LEU:HB3	1:A:35:VAL:HG13	1.83	0.60
1:B:72:ALA:CB	1:B:104:GLU:HG3	2.31	0.60
1:B:44:ARG:CB	1:B:44:ARG:HH11	2.04	0.59
1:A:146:ARG:NH1	1:B:94:VAL:O	2.35	0.59
1:B:124:ARG:O	1:B:128:THR:CG2	2.50	0.59
1:B:28:ALA:HA	1:B:38:VAL:HG21	1.85	0.59
1:A:4:ALA:N	3:A:2001:HOH:O	2.36	0.57
1:B:104:GLU:CG	1:B:134:ARG:NH2	2.65	0.57
1:B:33:LEU:HD22	1:B:37:ALA:HB3	1.85	0.56
1:B:6:ASP:OD2	1:B:41:ARG:NH1	2.39	0.56
1:B:33:LEU:HD22	1:B:37:ALA:HB1	1.86	0.56
1:B:7:ASP:HA	1:B:10:ARG:HG3	1.87	0.56
1:B:45:LEU:O	1:B:48:ARG:O	2.23	0.55
1:B:59:ASN:HD22	1:B:62:ALA:HB2	1.70	0.55
1:A:89:GLU:O	1:B:146:ARG:NH2	2.40	0.55
1:A:83:ASP:OD2	1:A:83:ASP:N	2.37	0.53
1:A:13:VAL:HB	1:B:58:ILE:HD12	1.91	0.53
1:A:25:SER:O	1:A:29:THR:HG23	2.08	0.53
1:B:34:SER:C	1:B:36:SER:H	2.12	0.52
1:A:80:GLN:HE21	1:A:80:GLN:CA	2.22	0.52
1:A:93:GLU:HG2	1:A:121:LEU:HD13	1.92	0.51
1:B:74:THR:OG1	1:B:104:GLU:OE2	2.21	0.50
1:B:8:ILE:HG21	1:B:41:ARG:NH2	2.27	0.50
1:A:150:PRO:HD2	1:B:63:VAL:HG12	1.94	0.49

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:LEU:O	1:A:126:ARG:HG3	2.12	0.49
1:B:146:ARG:HH11	1:B:146:ARG:CG	2.26	0.48
1:B:28:ALA:CA	1:B:38:VAL:HG21	2.43	0.47
1:A:150:PRO:HD2	1:B:63:VAL:CG1	2.44	0.47
1:B:83:ASP:N	1:B:83:ASP:OD2	2.38	0.46
1:A:14:ARG:HD2	3:A:2003:HOH:O	2.16	0.46
1:B:124:ARG:HD3	3:B:2013:HOH:O	2.17	0.45
1:A:95:GLU:CG	1:B:145:ASP:HA	2.48	0.44
1:B:24:LEU:HG	1:B:39:GLN:HG3	2.00	0.44
1:A:95:GLU:HG3	1:B:145:ASP:HA	1.99	0.43
1:A:146:ARG:NH2	1:B:89:GLU:HG3	2.33	0.43
1:B:14:ARG:HA	1:B:149:ILE:HD13	1.99	0.43
1:A:74:THR:HA	1:A:75:PRO:HD3	1.93	0.42
1:A:124:ARG:O	1:A:128:THR:CG2	2.58	0.42
1:B:21:ARG:O	1:B:22:ALA:C	2.57	0.42
1:A:76:LEU:HD23	1:A:76:LEU:HA	1.72	0.42
1:A:83:ASP:O	1:A:87:ARG:HG3	2.20	0.42
1:B:34:SER:O	1:B:36:SER:N	2.53	0.41
1:B:84:ALA:N	1:B:85:PRO:CD	2.83	0.41
1:A:105:SER:HB3	2:A:1151:LEU:HB3	2.03	0.40
1:B:73:ILE:HG22	1:B:133:THR:HG22	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/150 (97%)	142 (98%)	3 (2%)	0	100	100
1	B	145/150 (97%)	124 (86%)	16 (11%)	5 (3%)	4	5
All	All	290/300 (97%)	266 (92%)	19 (7%)	5 (2%)	11	18

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	28	ALA
1	B	145	ASP
1	B	30	ARG
1	B	35	VAL
1	B	29	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/126 (96%)	100 (83%)	21 (17%)	2	4
1	B	118/126 (94%)	97 (82%)	21 (18%)	2	3
All	All	239/252 (95%)	197 (82%)	42 (18%)	2	3

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	7	ASP
1	A	24	LEU
1	A	29	THR
1	A	44	ARG
1	A	45	LEU
1	A	67	LEU
1	A	71	VAL
1	A	74	THR
1	A	75	PRO
1	A	80	GLN
1	A	83	ASP
1	A	87	ARG
1	A	93	GLU
1	A	94	VAL
1	A	100	VAL
1	A	110	VAL
1	A	112	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	127	THR
1	A	128	THR
1	A	131	VAL
1	B	5	LEU
1	B	6	ASP
1	B	11	ILE
1	B	23	THR
1	B	24	LEU
1	B	36	SER
1	B	41	ARG
1	B	51	VAL
1	B	61	GLU
1	B	67	LEU
1	B	87	ARG
1	B	91	ILE
1	B	94	VAL
1	B	100	VAL
1	B	112	VAL
1	B	120	ASP
1	B	128	THR
1	B	130	ASN
1	B	132	ARG
1	B	145	ASP
1	B	146	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	GLN
1	A	80	GLN
1	B	59	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	LEU	A	1151	-	4,8,8	36.98	1 (25%)	5,10,10	6.36	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LEU	A	1151	-	-	0/4/8/8	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1151	LEU	CB-CA	73.95	3.62	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1151	LEU	CG-CB-CA	-11.59	45.36	116.28
2	A	1151	LEU	CB-CA-N	-8.17	92.20	110.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1151	LEU	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	147/150 (98%)	-0.39	3 (2%) 65 67	19, 38, 64, 82	0
1	B	147/150 (98%)	-0.11	5 (3%) 46 48	19, 43, 92, 97	0
All	All	294/300 (98%)	-0.25	8 (2%) 55 58	19, 40, 91, 97	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	35	VAL	3.9
1	B	36	SER	3.7
1	A	80	GLN	3.0
1	B	4	ALA	3.0
1	A	81	PRO	2.8
1	A	79	SER	2.6
1	B	25	SER	2.2
1	B	31	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	LEU	A	1151	9/9	0.90	0.24	2.57	68,71,73,75	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.